certainly have helped to fashion actually has considerable success in reconciling professional, scientific objectives with administrative requirements and realities. I refer to the Public Health Laboratory Service (PHLS) Board which I had the honour and pleasure to serve from 1963 to 1973. In 1961 this board took over the management of the service from the Medical Research Council, which from 1939 had directed the Emergency PHLS and later the PHLS for the Ministry of Health. The composition and functioning of the board, and its relations with the minister, followed essentially the three requirements set out in Sir Harold's paper. The board combined political, administrative, and professional skills. The board determined all major issues of policy, but left the day-to-day running of the PHLS to the medical director and the secretary to the Board. The service could often have done with more money, but the administration was skilled at helping the laboratory directors to find enough to fulfil a good many of their most important professional requirements. As a result, although there were quite numerous problems for the service, there were not insoluble problems in the service. Nor were there intolerable delays in reaching decisions and communicating them to those who sought answers. Visits, telephone calls, and personal discussion were the order of the day—not passing the problems on to someone else or even failing to answer petitioners. There were easy and fruitful relationships within the service and between the service and regional health boards, the Department of Health, the Medical Research Council, the veterinary service, the universities, the royal colleges, and indeed with all concerned with epidemiological microbiology.

I think that Sir Harold and the many other distinguished people who worked out the pattern for the PHLS from 1938 onwards were indeed successful in bestowing on the service "... at its centre an in-house capability, for policy formation and executive action" just as Sir Harold prescribes, as may be seen in detail from Landsborough Thomson's admirable record.¹ I think that the board and its headquarters functioned and function effectively, that it is a splendid example of the English (I mean *English*) genius for devising workable solutions to actual problems, and that the success of the PHLS may well point to ways of achieving the kind of integration which Sir Harold so rightly identifies as necessary for the National Health Service as a whole.

JAMES HOWIE

Edinburgh EH13 0BU

¹ Thomson AL, *Half a century of medical research*. Vol 2. London: HMSO, 1975:255-70.

Breast cancer trials—a new initiative

SIR,—Professor M Baum and others use some surprising statements and outdated concepts in attempting to justify their adjuvant breast cancer study using cyclophosphamide (13 September, p 742).

Firstly, the rationale for aggressive adjuvant combination chemotherapy is *not* based on "guesswork extrapolated from the behaviour of advanced disease." There is very strong experimental, theoretical, and clinical evidence¹⁻⁴ to show that the following principles should be applied if adjuvant chemotherapy in breast cancer is to be effective: (1) Treatment should be started as soon as possible—that is,

within a few days. (2) Combination chemotherapy will be far more effective than single-drug treatment (as mentioned in the National Institutes of Health consensus statement in the same issue of the *BMJ* (p 724). (3) Intervals between treatments should be the minimum consistent with clinical tolerance. (4) Reduction of dose or the use of single agents will reduce the probability of cure. The proposed trial ignores every one of these principles except the first.

Secondly, the implication that an inadequate trial should be carried out because surgeons and radiotherapists in the United Kingdom are too busy to carry out an adequate one is illogical. Is it really the case that many surgeons and radiotherapists are "naturally reluctant to take part in studies involving additional unrewarding effort" when such effort might lead to a significant improvement in the prognosis for breast cancer?

Thirdly, the old-fashioned assumption that aggressive chemotherapy is automatically associated with increased toxicity is not true. Over the past 11 years we have demonstrated over several thousand treatment cycles that intensive chemotherapy can be given safely by using simple principles derived from stem-cell kinetic studies.^{5,6} This approach has been confirmed by several other groups for many different tumour types, including breast cancer. Specifically, in an adjuvant study based on these principles using five drugs in combination it has been shown that there is no significant myelosuppression and minimal interference with the quality of the patient's life.⁷

Fourthly, why is the cyclophosphamide given daily for six days? This merely increases the likelihood of leucopenia without any reasonable prospect of improved therapeutic effect. Presumably this protocol is based on the relatively ineffective Nissen-Meyer study,⁸ the validity of which has been challenged.⁹ Therefore what is presented as a new initiative for the 1980s is really a rehash of an old initiative for the 1960s.

It is particularly unfortunate that, at a time when all the necessary requirements for a significant improvement in the survival rate in patients with stage II breast cancer can be met, the Cancer Research Campaign should support a 10-year study using cyclophosphamide alone. As Professor Baum and others admit, this trial is unlikely to have a major effect on the overall survival rates. It even violates the statement by the Cancer Research Campaign's own working party (4 October, p 918), which says that "to justify the expense, the question asked by the trial must be of critical relevance to the future management of the disease under study." Apart from confirming the conservative and outdated approach of many British oncologists to one of the major cancer problems of our time, the proposed study raises a major ethical question—namely, what are the patients going to be told? Will they know, for example, that the adjuvant treatment that they may be given is on present evidence less likely to prolong their survival than the protocol of Cooper?⁴

On logical and ethical grounds, I urge all clinicians interested in this trial to read the most recent literature before they allocate patients to this study.

L A PRICE

London W1N 1AA

¹ Burchenal JH. *Cancer* 1976;37:46-56.

² Hill BT, Price LA. In: Stoll B, ed. *Secondary spread in breast cancer*, vol 3. London: Heinemann, 1977: 193-211.

³ Goldie JH, Goldman A. *Cancer Treatment Reports* 1979;63:1727-31.

⁴ Cooper RG, Holland JF, Glidewell O. *Cancer* 1979; 44:793-8.

⁵ Price LA, Goldie JH. *Br Med J* 1971;iv:336-9.

⁶ Goldie JH, Price LA. *Br Med J* 1977;ii:1064-5.

⁷ Morrison JM. *Cancer Treatment Reviews* 1979;6, suppl:87-90.

⁸ Nissen-Meyer R, Kjellgren K, Malmio K, Mansson B, Norin T. *Cancer* 1978;41:2088-98.

⁹ Carter SK. *Cancer Chemother Pharmacol* 1980;4: 147-63.

*.*We sent this letter to the authors, and Professor Baum replies below.—Ed, *BMJ*.

SIR,—Dr Price is making two fundamental errors of scientific philosophy.

In the first he is confusing the building of a hypothesis with its testing. He is thus guilty of the worst kind of conceptual rationalism. It is one thing to propose that complex chemotherapeutic schedules developed from experiments on animal models might have a significant impact on mortality for patients treated for early breast cancer, but it is altogether a different matter to confirm this effect with a properly designed clinical trial. Such experiments are in fact already in progress both in the United States of America and under the good auspices of the West Midlands oncology group, but it will be many years before we know whether the results corroborate the hypothesis proposed by Dr Price.

The other philosophical pitfall concerns the decision about what constitutes a *major* effect on overall survival rates for breast cancer. It may be true that in the long run the Price/Hill kinetic approach¹ in the hands of expert medical oncologist will have a profound effect on breast cancer mortality among patients treated in specialist centres, but at the moment very few centres exist in the world at large capable of delivering these complex and potentially lethal regimens. Incidentally it is quite untrue that the subjective toxicity is minimal as judged by a very recent report in the *BMJ*.² In the meantime, if very simple adjuvant techniques—as proposed, for example, in the new Cancer Research Campaign trial—can be shown to have an effect, albeit of a modest nature, then because of their very ease of administration they would be immediately acceptable by clinicians practising not only in the United Kingdom, but throughout the developing world. I leave the statisticians to calculate whether a 30% improvement in 10-year survival among patients treated in five cancer centres in the United Kingdom, each of which may see 300 cases a year, will have a more *major* effect on overall survival rates than a 10% improvement in survival seen among the 20 000 new cases of breast cancer that present in the United Kingdom each year, or the hundreds of thousands of patients presenting with the disease in the developing world.

In contradistinction to the anticipated benefits of the complex kinetic approach, there is some direct evidence, based on actual (not actuarial) 10-year survival figures from the Toronto and the Scandinavian trials,^{3,4} that supports the use of either endocrine therapy or a short course of cyclophosphamide. The only reason why such adjuvant therapy is not routinely applied throughout the world at present has been modesty on the part of the authors of these studies combined with criticism of the experimental methods used. It is therefore essential to repeat these experiments as quickly as possible before we should even contemplate the necessity of total reorganisation of oncology services that would be necessary to deliver the "adequate" therapy proposed by Dr Price.

To deal with some of the individual points that Dr Price raises, it should first of all be emphasised that no concept is "outdated" if corroborated by evidence. Thus, although we are now repeatedly told that single short-term chemotherapy *should* not be effective, none the less there remains evidence from two prospective randomised trials^{4,5} and one uncontrolled series,⁶ that such an approach is followed by benefits in survival. Secondly, we are assured that combination therapy is more