- 36 Lansky S, Cairns N, Stevenson L, Narin G. CNS prophylaxis: studies showing impairment in verbal skills and academic achievement *Am J Ped Hematol Oncol* 1984; 6:183.
- 37 Moss H, Nannis ED. The effects of prophylactic treatment of the CNS on the intellectual functioning of children with ALL. *Am J Med* 1981;71:47–52.
- 38 MacLean WE, Noll RB, Stebbens JA, Kaleita TA. Neuropsychological effects of cranial irradiation in young children with acute lymphoblastic leukemia nine months after diagnosis. *Arch Neurol* 1995;52:156–60.
- 39 Waber DP, Tarbell NJ. Toxicity of CNS prophylaxis for childhood leukemia. *Oncology (Huntingt)* 1997;11:259–64.
- 40 Hill JM, Kornblith AB, Jones D, *et al.* A comparative study of the long-term psychosocial functioning of childhood acute lymphoblastic leukemia. Survivors treated by intrathecal methotrexate with or without cranial irradiation. *Cancer* 1998;82:208–18.
- 41 Brown RT, Madan-Swain A, Pais R, Lambert RG. Cognitive status of children treated with central nervous system prophylactic chemotherapy of acute lymphocytic leukaemia. *Archives of Clinical Neuropsychology* 1992;7:481–97.

Ethology and cortical visual impairment

In industrialised countries there has been a switch in recent decades from ocular to non-ocular causes of visual impairment. The term cortical visual impairment (CVI) is applied to visual loss arising from defects in the geniculostriate pathways although, by default, other neurological defects may be included. The ophthalmological investigation of CVI is difficult and researchers in Utrecht have enlisted help from their local department of ethology (G Porro and colleagues. *British Journal of Ophthalmology* 1998;82:1231–5. See also editorial, pages 1225–6). [The Greek word *ethos* is variously translated as character, manners, habits, or behaviour. Ethology is, therefore, the study of behaviour in animals and people.]

They studied six girls and three boys aged 19–116 months, all of whom had cerebral palsy, lacked visual contact, had no stable visual fixation, and had defied attempts at visual function assessment. Each was observed by an ethologist for 15 minutes, and responses to visual stimuli were observed and recorded on video. As a result of their observations these workers defined six direct signs (indicative of visual perception) and six indirect signs (regarded as supportive or confirmatory). The direct signs were: intermittent fixation towards the stimulus, looking past it, head movement towards it, following, reaching out, and looking away while reaching. The indirect signs were: avoiding behaviour, posture change, smiling, blinking, facial expression change, and stereotypic behaviour.

More detailed observational studies may improve the understanding of behavioural adaptations to CVI in these children.

ARCHIVIST