- 6 Oakley GA, Muir T, Ray M, Kennedy R, Girdwood RWA, Donaldson MDC. Increased incidence of congenital anomalies in infants with transient TSH elevation detected by newborn screening. *J Pediatr* 1998;132:726-30.
- 7 Lobo E de H, Khan M, Tew J. Community study of hypothyroidism in Down's syndrome. *BMJ* 1980;24:1253.
- 8 Zori RT, Schatz DA, Ostrer H, *et al.* Relationship of autoimmunity to thyroid dysfunction in children and adults with Down syndrome. *Am J Med Genet Suppl* 1990;7:238-41.
- 9 Kinnell HG, Gibbs N, Teale JD, *et al.* Thyroid dysfunction in institutionalised Down syndrome adults. *Psychol Med* 1987;17:387-92.
- 10 Murdoch JC, Ratcliffe WA, McLarty DG, Rodger JC, Ratcliffe JC. Thyroid function in adults with Down's syndrome. *J Clin Endocrinol Metab* 1977;44:453-8.
- 11 Sare Z, Ruvalcaba RHA, Kelley VC. Prevalence of thyroid disorder in Down syndrome. *Clin Genet* 1978;14:154-8.
- 12 Pozzan GB, Rigon F, Girelli ME, *et al.* Thyroid function in patients with Down syndrome: preliminary results from non-institutionalised patients in the Veneto region. *Am J Med Genet Suppl* 1990;7:57-8.
- 13 Mani C. Hypothyroidism in Down syndrome. *Br J Psychiatry* 1988;153:102-4.
- 14 Karlsson B, Gustafsson J, Hedov G, Ivarsson S-A, Anneren G. Thyroid dysfunction in Down's syndrome: relation to age and thyroid autoimmunity. *Arch Dis Child* 1998;79:242-5.
- 15 Selikowitz M. A five year longitudinal study of thyroid function in children with Down syndrome. *Dev Med Child Neurol* 1993;35:396-401.
- 16 Cronk C, Crocker AC, Pueschel SM, *et al.* Growth charts for children with Down's syndrome; 1 month to 18 years of age. *Pediatrics* 1998;81:102-10.
- 17 Findlay C, Donaldson M. How often should we screen for hypothyroidism in girls with Turner's syndrome [letter]? *Arch Dis Child* 1998;78:500-1.

Cost of chickenpox

The incidence of chickenpox is said to be approximately equal to the number of births. Over 95% of cases are in children under 15 with the highest incidence in the 5-9 year age range. Details of the financial cost of the disease have been reported from Canada.

In two papers (Barbara Law and colleagues. *Pediatrics* 1999;104:1-6 and 7-14) costs are calculated from studies of 179 otherwise healthy 1-9 year olds with uncomplicated chickenpox treated at home, 160 otherwise healthy children with complicated (treated in hospital) chickenpox, and 40 children with leukaemia admitted to hospital with chickenpox. For uncomplicated cases the total cost to society per case was estimated at around CDN\$370 for younger children and CDN\$240 for children aged 5-9. Medical costs were about 10% of the total. The total national cost was estimated at CDN\$110 million annually.

For children admitted to hospital the total cost to society per case was around CDN\$8000 for otherwise healthy children and for children with leukaemia the medical cost of an admission with chickenpox was over CDN\$7000. This brought the estimated total national cost of chickenpox to CDN\$122 million, with CDN\$24 million of that cost being borne by the health ministry. Added to that there were considerable quality of life costs for children and parents. In the USA it has been calculated that vaccination against chickenpox would result in a total saving to society of US\$66 per child vaccinated. (CDN\$1 approximately US\$0.68.)

ARCHIVIST