- 9 Krizova P, Musilek M. Changing epidemiology of meningococcal invasive disease in the Czech Republic caused by new clone *Neisseria meningitidis* C:2a:P1.2(P1.5), ET-15/37. *Cent Eur J Public Health* 1995;3:189-94.
- 10 Kriz P, Vlkova J, Bobak M. Targeted vaccination with meningococcal polysaccharide vaccine in one district of the Czech Republic. *Epidemiol Infect* 1995;115:411-18.
- 11 Clayton D, Hills M. *Statistical models in epidemiology*. Oxford: Oxford University Press, 1993.
- 12 Krizova-Kuzemenska P, Benes C. Predbezna informace o nove zjistovane zavazne epidemiologicke situaci invazivnich meningokokovych onemocneni v Ceske republice [Preliminary information on a new epidemiological situation regarding invasive meningococcal disease in the Czech Republic]. *Zpravy Centra Epidemiol Mikrobiol* 1993;2:15.
- 13 Bobak M, Kriz B, Leon DA, Danova J, Marmot M. Socioeconomic factors and height of preschool children in the Czech Republic. *Am J Public Health* 1994;84:1167-70.
- 14 Pikhart H, Prikazsky V, Bobak M, Kriz B, Celko M, Danova J, Pryl K, Pretel J. Association between ambient air concentrations of nitrogen dioxide and respiratory symptoms in children in Prague, Czech Republic. Preliminary results from the Czech part of the SAVIAH study. *Cent Eur J Public Health* 1997;5:82-5.
- 15 Cope GF, Heatley RV. Cigarette smoking and intestinal defence. *Gut* 1992;33:721-3.
- 16 Barton JR, Riad MA, Gaze MN, Maran AGD, Ferguson A. Mucosal immunodeficiency in smokers, and patients with epithelial head and neck tumours. *Gut* 1990;31:378-82.
- 17 Dye JA, Adler KB. Effects of cigarette smoke on epithelial cells of the respiratory tract. *Thorax* 1994;49:825-34.
- 18 Green GM, Carolin D. The depressant effect of cigarette smoke on the in vitro antibacterial activity of alveolar macrophages. *N Engl J Med* 1967;276:421-7.
- 19 El Ahmer OR, Essery SD, Saadi AT, Raza MW, Ogilvie MM, Weir DM, Blackwell CC. The effect of cigarette smoke on adherence of respiratory pathogens to buccal epithelial cells. *FEMS Immunol Med Microbiol* 1999;23:27-36.
- 20 Cartwright K. Meningococcal carriage and disease. In: Cartwright K, ed. *Meningococcal disease*. Chichester: John Wiley & Sons, 1995:115-46.
- 21 Colley JRT, Holland WW, Corkhill RT. Influence of passive smoking and parental phlegm on pneumonia and bronchitis in early childhood. *Lancet* 1974;ii:1031-4.
- 22 Bonham GS, Wilson RW. Children's health in families with cigarette smokers. *Am J Public Health* 1981;71:290-3.
- 23 Dybing E, Sanner T. Passive smoking, sudden infant death syndrome (SIDS) and childhood infections. *Hum Exp Toxicol* 1999;18:202-5.
- 24 Koupilova I, Vagero D, Leon DA, Pikhart H, Prikazsky V, Holcik J, Bobak M. Social variation in size at birth and preterm delivery in the Czech Republic and Sweden, 1989-1991. *Paediatr Perinatal Epidemiol* 1998;12:7-24.
- 25 Koupilova I, Bobak M, Holcik J, Pikhart H, Leon D. Increasing social variation in birth outcome in the Czech Republic after 1989. *Am J Public Health* 1998;88:1343-7.
- 26 Bobak M, Hertzman C, Skodova Z, Marmot M. Socioeconomic status and cardiovascular risk factors in the Czech Republic. *Int J Epidemiol* 1999;28:46-52.
- 27 United Nations Children's Fund. *Children at risk in Central and Eastern Europe: perils and promises. Central and Eastern Europe in transition. Public policy and social conditions*. Regional Monitoring Report No. 4. Florence: UNICEF, 1997.
- 28 Jones IR, Urwin G, Feldman RA, Banatvala N. Social deprivation and bacterial meningitis in North East Thames region: three year study using small area statistics. *BMJ* 1997;314:794-5.

Neutralising enterotoxin

The verotoxin (Shiga toxin) of Shiga toxin producing organisms—such as *Escherichia coli* 0157:H7, binds to specific carbohydrate containing receptors on intestinal epithelial cells and on renal and vascular endothelial cells. Treatment aimed at preventing this binding should be effective. Work in California (Adrienne W Paton *et al.* *Nature Medicine* 2000;6:265-70; see also commentary *Ibid*: 257-8) may have made this possible.

Theoretically, the enterotoxin could be bound by administering synthetic carbohydrate which mimics the natural enterotoxin receptor, but the problem with this approach is that such synthetic carbohydrate binds only small amounts of enterotoxin and very large doses would be needed. The Californian workers have apparently overcome this difficulty in an ingenious way. They took a strain of *E coli* which lacked the enzymes responsible for lipopolysaccharide production (glycosyl transferases) and inserted into it two new galactosyl transferase genes, one from a meningococcus and one from a gonococcus. The *E coli* then produced lots of lipopolysaccharide with the same end units as the enterotoxin receptor. The dried bacteria were 10 000 times more effective at binding enterotoxin than was the synthetic carbohydrate previously available, and a live or dead preparation of the transformed *E coli* given orally to mice protected them against a lethal dose of verotoxin.

Another recent report, in *Nature* (PI Kitov *et al.* 2000;403:669-72) described the stereochemical synthesis of a complex trisaccharide containing molecule (called STARFISH because of its configuration) with high affinity for verotoxin. It is suggested that oral administration of the bacterial preparation might bind verotoxin in the gut and intravenous administration of the STARFISH complex might deal with systemic verotoxin.

The hope, of course, is that these new preparations can be given at the right time and in sufficient quantity to influence the course or development of the haemolytic uraemic syndrome.

ARCHIVIST