LETTER TO THE EDITOR

Stress protein response in human oesophageal epithelium may be influenced by the ex vivo culture technique

EDITOR,—In their recent article Hopwood *et al* (*Gut* 1997;41:156-63) conclude that only freshly acquired specimens cultured for less than six hours are suitable for studying stress responses ex vivo. This statement arises from the observation that cells from unheated oesophageal biopsy samples cultured for 22 hours (in unsupplemented HAMS-F10 media in an incubator containing room air with 5% CO₂ at 37°C) exhibited a "stressed" state. The maintenance ex vivo of gastrointestinal biopsy samples has posed formidable problems in the past; however, successful organ culture of oesophageal biopsy specimens for up to 48 hours has been reported by Browning and Trier using Trowell's technique,^{1,2} and in some cases oesophageal tissue has been cultured for up to seven months.³ For successful organ culture the conditions are critical, particularly the addition of calf serum and insulin to the media, and incubation in a gassed mixture of 95% O₂ and 5% CO₂.¹ Under these conditions, ultrastructural integrity, proliferative activity and protein synthesis are maintained. In Hopwood *et al*'s study, standard tissue culture rather than organ culture conditions were used and this may account for the finding that prolonged culture mimicked thermal stress. Although the tissue may appear intact by light microscopic examination, the integrity of the cultures can be readily assessed by lactate dehydrogenase release into the media and by the electron microscopic appearance of the tissue.⁴

It has recently been shown that acid has a dynamic effect on the cell phenotype of Barrett's oesophagus depending on the pattern of acid exposure. However, these phenotypic effects were only apparent after 24 hours of organ culture.⁴ Therefore, more prolonged organ culture may be necessary in order to investigate fully the oesophageal stress protein response to environmental factors such as oesophago-gastroduodenal refluxate. To conclude that analysis of stress responses using human oesophageal tissue is only feasible following short term culture using the methods described in this study is premature.

R C FITZGERALD

Lecturer in Gastroenterology,
Oldchurch Hospital, Havering NHS Trust,
and St Bartholomew's and the Royal London
School of Medicine and Dentistry,
Digestive Disease Research Centre,
2 Turner Street,
Whitechapel, London E1 2AD, UK

- 1 Trier JS. Organ culture of the mucosa of human small intestine. *Methods Cell Biol* 1980; 21B:365-84.
- 2 Howdle PD. Organ culture of the gastrointestinal mucosa — a technique for the study of the human gastrointestinal tract. *J Postgrad Med* 1984;69:645-52.
- 3 Hillman E. Human esophageal organ culture studies. *Methods Cell Biol* 1980;21B:332-48.
- 4 Fitzgerald RC, Omary MB, Triadafilopoulos G. Dynamic effects of acid on Barrett's esophagus: an ex vivo differentiation and proliferation model. *J Clin Invest* 1996;98: 2120-8.

Reply

EDITOR,—Stress proteins comprise a wide class of proteins including drug metabolising enzymes, protein kinases/phosphatases, molecular chaperones, metabolic enzymes, DNA repair enzymes, and transcription factors, all of which coordinately respond to subtle or dramatic changes in the extracellular milieu.¹ Most studies in the field have centred on regulation of the stress proteins using microbes, invertebrates, and human tumour cell lines, and the data acquired do not necessarily reflect in vivo regulation of the same pathways in mammalian tissue. Developing ex vivo cell type specific models will be required to obtain a more physiologically relevant understanding of the effects of stress protein function on normal human tissue integrity and repair.

However, ex vivo organ culture studies will, by virtue of removing the cells from their normal environment, acutely stress and alter some pathways being investigated. Rapid changes in primary/secondary metabolites and signal transduction cascades are well documented following cellular perturbation.²⁻⁵ Such perturbations can stem, in part, from transiently lowered oxygen concentrations during tissue extraction and/or from protein/lipid/DNA damage resulting from oxidative stress.⁶ Activation of these transduction cascades changing gene expression may prevent the physiological study of a stress response system ex vivo.

For example, the AMP activated protein kinase cascade is rapidly perturbed in rat liver ex vivo and the tissue needs to be snap frozen in situ prior to animal death to prevent such hypoxic induced changes in metabolism from altering the system.^{7,8} In addition, protein kinase C activity in rat brain has a half-life of minutes ex vivo and very rapid manipulation is required to preserve enzyme activity.⁹ Induction of acute stress can be prolonged; during short term ischaemia and reperfusion in cardiac tissue, HSP70 induction occurs for many hours after injury.¹⁰ As such, before setting up an ex vivo system to study the heat shock response in normal human oesophageal epithelium, I suspected that metabolic stress would be imposed on the tissue after biopsy and we first set out to determine whether there was, in fact, a biochemical stress imposed by the act of sample collection. Ex vivo culture methods used were those previously optimised to show differences in endocytosis between normal and inflamed oesophageal tissue.¹¹ Using these methods to allow direct comparison between the two pathways, we demonstrated that even before organ culture, changes in the steady state level of protein synthesis could occur, compared with rapidly frozen tissue (*Gut* 1997;41:156-63; fig 1), thus any conclusions we formed on ex vivo heat shock may be compromised by the underlying stress to the cells.

Despite these caveats, successful analysis of the heat shock response ex vivo has been fortuitously possible and we detected two major stress proteins synthesised after heat shock. We have since purified and generated antibodies to one of the heat shock proteins and showed, by immunohistochemistry, unique localisation in vivo (Vojtesek B, Moitra S, Johnston D, *et al*, manuscript submitted). Nevertheless, we feel that further basic and clinical research in this area will be required to determine whether these and other stress pathways are similarly regulated in vivo. One

approach, based on the use and success of in vivo NMR to study metabolic flux in a non-invasive manner,¹² overcomes the biochemical stresses that are imposed on cells when studying biological processes ex vivo, but this method obviously has its limitations for many enzymatic pathways.

It is my opinion that freshly acquired biopsy samples are preferred to study stress responsive systems and, despite the well documented fact that addition of serum and growth factors can control the survival of cells ex vivo,¹³ the relevance of ex vivo metabolic state to in vivo metabolism is not yet clear. Before premature claims are made that ex vivo organ culture does not induce metabolic stress, I suggest sensitive assays are developed to detect possible perturbations, such as changes in metabolites (such as ATP concentrations), changes in intracellular pH, or activation of well recognised regulators of cell stress, such as JNK or MAP kinases^{3,14} and NF- κ B or p53 transcription factors.^{15,16} Markers of stress, including morphological integrity, proliferative activity, protein synthesis, and lactate dehydrogenase release, are not sufficiently sensitive nor specific to detect subtle changes in metabolism. In addition, the use of radiolabelled thymidine to measure proliferative capacity and the use of radiolabelled methionine to measure protein synthesis can activate cellular damage response pathways^{17,18} and are therefore inherently perturbing to the system under study. As such, it would not be accurate to claim that ex vivo organ culture of oesophageal epithelium does not activate interfering stress responses until more sensitive assays are developed.

T R HUPP

Department of Molecular and Cellular Pathology,
University of Dundee,
Dundee DD1 9SY, UK

- 1 Minowada G, Welch WJ. Clinical implications of the stress responses. *J Clin Invest* 1995;95:2-12.
- 2 Schlesinger MJ, Ryan C, Chi MM, *et al*. Metabolite changes associated with heat shocked avian fibroblast mitochondria. *Cell Stress Chaperones* 1997;2:25-30.
- 3 Bradham CA, Stachlewitz RF, Gao W, *et al*. Reperfusion after liver transplantation in rats differentially activates the mitogen-activated protein kinases. *Hepatology* 1997;25:1128-35.
- 4 Van Veldhoven PP, Bell RM. Effect of harvesting methods, growth conditions and growth phase on diacylglycerol levels in cultured human adherent cells. *Biochim Biophys Acta* 1988;959:185-96.
- 5 Delhotal B, Lemonnier F, Couturier M, *et al*. Influence of two cell harvesting methods on intracellular ATP and amino acid concentrations in human fibroblast cultures. *Biochimie* 1983;65:121-5.
- 6 Jaeschke H. Mechanisms of oxidant stress-induced acute tissue injury. *Proc Exp Biol Med* 1995;209:104-11.
- 7 Corton JM, Hardie DG. Regulation of lipid biosynthesis by the AMP-activated protein kinase and its role in the hepatocellular response to stress. *Prog Liver Dis* 1996;14:69-99.
- 8 Hardie DG, Carling D. The AMP-activated protein kinase—fuel gauge of the mammalian cell? *Eur J Biochem* 1997;246:259-73.
- 9 Go M, Koumoto J, Kikkawa U, *et al*. Protein kinase C purification by high-performance liquid chromatography: an improved method. *Methods Enzymol* 1987;141:424-8.
- 10 Currie RW, Tabguay RM, Kingma JG. Heat shock response and limitations of tissue necrosis during occlusion/reperfusion in rabbit hearts. *Circulation* 1993;87:963-71.
- 11 Hopwood D, Spiers EM, Ross PE, *et al*. Endocytosis of fluorescent microspheres by human oesophageal epithelial cells: comparison between normal and inflamed tissue. *Gut* 1995;37:598-602.
- 12 Radda GK. Control of energy metabolism during muscle contraction. *Diabetes* 1996;45:S88-92.

- 13 Evan GI, Wyllie AH, Gilbert CS, *et al.* Induction of apoptosis in fibroblasts by c-Myc protein. *Cell* 1992;69:119–28.
- 14 Hu Y, Metzler B, Xu Q. Discordant activation of stress-activated protein kinases or c-Jun NH₂-terminal protein kinases in tissues of heat-stressed mice. *J Biol Chem* 1997;272:9113–19.
- 15 Pinkus R, Weiner LM, Daniel V. Role of oxidants and antioxidants in the induction of AP-1, NF-kappaB, and glutathione S-transferase gene expression. *J Biol Chem* 1996;271:13422–9.
- 16 Hupp TR, Lane DP. Two distinct signalling pathways are required to activate p53 in a casein kinase II-independent manner. *J Biol Chem* 1995;270:18165–74.
- 17 Dover R, Jayaram Y, Patel K, Chinery R. p53 expression in cultured cells following radioisotope labelling. *J Cell Sci* 1994;107:1181–4.
- 18 Yeargin J, Haas M. Elevated levels of wild type p53 induced by radiolabeling of cells leads to apoptosis or sustained growth arrest. *Curr Biol* 1995;5:423–31.

BOOK REVIEWS

Current Trends in Digestive Ultrasonography. Edited by Gandolfi L, Fukuda M. (Pp 394; illustrated; \$298.25.) Basel, Switzerland: Karger, 1997. ISBN 3-8055-6374-4.

The editors of volume 24 of *Frontiers in Gastrointestinal Research* have enlisted the help of some of the leading international experts in ultrasound to provide a timely review of the leading edges of ultrasound applications in gastroenterology.

The ultrasound stethoscope is regarded by many as the ultimate progression of this technology into gastroenterological practice. In the UK, digestive ultrasonography has largely remained the province of radiologists, although recently the Royal College of Radiologists (RCR) has recognised that there is a legitimate desire by some medical non-radiologists, such as gastroenterologists, to undertake ultrasound examinations as a direct extension of their clinical examination. The RCR has offered guidance for their training in ultrasound, and this book would be essential reading for any gastroenterologist embarking upon such a training programme. However, ultrasound always has been (and still is) highly operator dependent, and whether gastroenterologists and surgeons will ever put the requisite time and effort into acquiring sufficient practical expertise remains open to question.

Most of the chapters in this book are reviews of current ultrasound practice, and are well illustrated and comprehensively referenced. The roles of ultrasound in the diagnosis of hepatobiliary, pancreatic and gastrointestinal tracts are covered, with particular emphasis on the newer problems such as HIV infection and the assessment of liver transplants. There is a fascinating review of ultrasound in tropical medicine, and this is an area where ultrasound equipment scores over other imaging modalities such as computed tomography and magnetic resonance imaging because of the obvious advantages of cost, reliability and portability.

Ultrasound guided therapy for liver tumours, intra-operative ultrasound, and the applications of Doppler ultrasound in gastroenterology are discussed and these are topics

with which many gastroenterologists may not be familiar, whilst endoscopic ultrasound and the use of ultrasound for functional studies in the gastrointestinal system may be of particular interest to readers of this journal.

The editors have succeeded in their stated intention to put together a volume which will be of interest to radiologists, gastroenterologists, internists, and surgeons. Texts such as this are needed, both to inform established specialists of recent developments in gastroenterological ultrasound, and also to stimulate those in training to explore further the newer applications described. The state of the art is described, and the contributors have clearly demonstrated how ultrasound can permit quick, reproducible and non-invasive diagnosis as well as allowing for accurate image guided interventions. This volume will not only inform and encourage those who are not yet ultrasound aficionados to take more of an interest in the technique, but may also stimulate others to use ultrasound in future research endeavours.

H IRVING

Deglutition and its disorders. Anatomy, physiology, clinical diagnosis, and management. Edited by Perlman AL, Schulze-Delrieu KS. (Pp 522; illustrated; £49.95.) Singular Publishing Group, 1997. ISBN 9781565 936218.

The act of swallowing has been described as the most complex reflex which can be reproduced by the stimulation of a peripheral nerve. Such complexity is also encountered in the aetiology, pathogenesis, investigation, and management of deglutition disorders. The senior editor of this comprehensive multi author review, Adrienne Perlman, is a speech and language therapist, now based in Champaign, Illinois, with an outstanding personal record of innovative research in the field of swallowing. Her contributors include most of the principal prominent names in North American dysphagia research, whether internists, neurologists, radiologists or, of course, speech and language therapists. Few readers are likely to approach this volume with important questions about dysphagia which would not, at least in part, be answered by in this comprehensive text. As a reference book, it has very few competitors. Thus any library which is required to carry the journal *Dysphagia* will find this text frequently consulted. Overall, the line diagrams are well chosen and the radiographic images clear. Some of the endoscopic photographs inevitable lose a little in black and white reproduction.

For many practising speech and language therapists the benchmark chapter will be the one on videoendoscopic assessment of swallowing. It is written by the North American leaders in the field, but has certain limitations for therapists practising in countries, such as the United Kingdom, where the technique has yet to be fully integrated into the repertoire of the speech and language therapist. The interpretative section is outstanding, but the equipment section rather suggests that a very substantial investment is required to set up a good system. What many Europeans need is a message that the method can be initiated with very little new capital outlay, and the more sophisticated set-ups, with videotimers, videoprinters and the like graduated to over a period of time.

Some of the individual chapters cover topics which are not often identified in a pure

dysphagia context—for example, the chapter on swallowing and feeding in paediatric patients. Others reflect Dr Perlman's special interests—for example, the use of electromyography in the diagnosis of deglutition disorders. The inclusion of these superspecialist topics is, however, entirely justified because it is only someone with this superspecialist interest in dysphagia who is likely to make a personal purchase of this volume.

The great range of health care professionals interested in dysphagia (reflected in the multidisciplinary authorship list) will come to this text with different expertise. This justifies the inclusion of basic information within each section included in the text. For example, those who are radiologists will find the radiology sections oversimplified but the non-radiologists may find the introductory information invaluable. Four to six multiple choice questions are included at the end of each of the 16 chapters. A comprehensive glossary is also included. *Deglutition and its disorders* therefore attempts to offer a complete dysphagia education, based on minimal prior knowledge of the relevant concepts. Such a goal may well find a niche market in the United Kingdom, where, at the present time, dysphagia work load comprises a huge proportion of clinical speech and language therapy time. Most speech therapy degree courses do not include a significant amount of dysphagia instruction at the undergraduate level and postgraduate experience is often based on practical experience supplemented by ad hoc clinical courses. Thus, this book may well improve the understanding of clinical practice of the group of professionals who bear the brunt of clinical decision making in dysphagic patients in the United Kingdom today. Every speech and language therapy department seeing dysphagic patients should have a copy of this volume on its shelf.

J A WILSON

NOTES

Colorectal Disease in 1998

The 9th Annual Colorectal Disease in 1998: An International Exchange of Medical and Surgical Concepts will be held at Marriott's Harbor Beach Resort, Fort Lauderdale, Florida, USA, from 19 to 21 February 1998. Further information from: Cleveland Clinic Florida, Department of Education, 2950 West Cypress Creek Road, Fort Lauderdale, FL 33309-1743, USA. Fax: 954 978 5539; Other: 800 359 5101, ext 5056; Local/international: 954 978 5056; email: jagels@cesmtp.ccf.org.

6th Southeast European Congress of Paediatric Surgery: Short Bowel Syndrome

The 6th Southeast European Congress of Paediatric Surgery: Short Bowel Syndrome will be held in Graz, Austria, on 22–23 May 1998. Further information from: Dr Günther Schimpl, Department of Paediatric Surgery, Auenbruggerplatz 34, A-8036 LKH-Graz, Austria. Tel: +43 316 385 3762; Fax: +43 316 385 3775.