Are complications of endoscopic sphincterotomy age related?

EDITOR,—Deans *et al* (*Gut* 1997;41:545–8) have supported the use of endoscopic sphincterotomy for bile duct stones in young people in preference to laparoscopic bile duct exploration. They do this because sphincterotomy causes no more complications in young than in old people. Any comparison between the two methods must also take into account the complications, particularly pancreatitis, that occurred in the remaining two-thirds of the patients undergoing endoscopic retrograde cholangiopancreatography (ERCP) without sphincterotomy. Laparoscopic duct exploration is still a relatively new technique. Rhodes *et al* reported 129 explorations with one case of postoperative pancreatitis.¹ We have had no pancreatitis in 53 explorations. Until larger series are reported judgement is impossible.

For the treatment of bile duct stones a comparison of therapeutic success and the manoeuvres required to achieve it are required on an intention-to-treat basis.

The consequences of sphincterotomy and reflux in the very long term are not known. Most bile duct stones are predominantly composed of pigment, thought to be caused by reflux through an incompetent sphincter.² They are seen in about 10% of patients after sphincterotomy but these data include only small numbers of young patients with long term follow up. Data for the events of 20–30 years after sphincterotomy are not available, yet a quarter of the patients in the study of Deans *et al* might be expected to live this length of time. Apart from stone formation, long term duodenobiliary reflux may have other adverse effects. Kurumado *et al* reported dysplastic change in rat mucosa after formation of biliary enteric fistulae.³ Biliary malignancy is seen in anomalous pancreatico-biliary junctions, postulated to be caused by unusual reflux.⁴ The application of these reports to the effect of sphincterotomy needs to be cautious but they act as a warning that sphincterotomy might not be entirely benign in the long term.

Laparoscopic bile duct exploration offers the possibility of safe choledocholithotomy without sphincter destruction and it merits further study.

M H THOMPSON
Department of Surgery,
Southmead Hospital,
Bristol BS10 5NB, UK

- 1 Rhodes M, Nathanson L, O'Rourke N, *et al*. Laparoscopic exploration of the common bile duct: lessons learned from 129 consecutive cases. *Br J Surg* 1995;82:666–8.
- 2 Lotveit T, Osnes M, Aune S, *et al*. Studies of the choledochoduodenal sphincter in patients with and without juxta-papillary duodenal diverticula. *Scand J Gastroenterol* 1980;15:875–80.
- 3 Kurumado K, Nagai T, Kondo Y, *et al*. Long term observations on morphological changes of choledochal epithelium after choledochocenterostomy in rats. *Dig Dis Sci* 1994;39:809–820.
- 4 Tanaka K, Nishimura A, Yamada K, *et al*. Cancer of the gallbladder associated with anomalous junction of the pancreaticobiliary duct system without bile duct dilatation. *Br J Surg* 1993;80:622–4.

Reply

EDITOR,—This study was conceived because we had heard clinicians at several national and international meetings stating that young patients with stones in the common bile duct are better treated by surgical exploration than ERCP sphincterotomy because the latter has

a 1% mortality. These clinicians were advocating open exploration of the common bile duct and to our knowledge none of them were performing laparoscopic bile duct exploration. Their statements were based on the global mortality from ERCP, irrespective of age and we were unable to find any scientific evidence relating the risks of ERCP sphincterotomy with age.

Our study is a prospective, multicentre audit of 1000 endoscopic sphincterotomies. No patient underwent ERCP without sphincterotomy and there was no comparison with any other technique such as laparoscopic common bile duct exploration. Mr Thompson's comments about a comparison with laparoscopic bile duct exploration and about "the remaining two-thirds of the patients undergoing ERCP without sphincterotomy" cannot therefore be directly answered by our report. We would like to emphasise that we use a selective policy for performing ERCP in patients undergoing laparoscopic cholecystectomy.^{1,2} This policy has become more selective since we began using intravenous infusion cholangiography in patients with possible bile duct stones, a technique we have shown to be as effective as ERCP but with fewer complications and less cost.³

We accept that "data for the events of 20–30 years after sphincterotomy are not available." Such statements are true for the long term consequences of all recent innovations, including laparoscopic bile duct exploration. Concerns about the prolonged effect of duodenobiliary reflux in animals have not been substantiated by the few reports looking at the long term effects of ERCP sphincterotomy in humans. In a study of 100 patients undergoing ERCP sphincterotomy followed for a median of 15 years, late complications, such as recurrent ascending cholangitis or malignant degeneration, were not observed.⁴ Recurrent bile duct stones, however, did develop in 24% of patients. The authors concluded that stone recurrence remains the most important long term problem after ERCP sphincterotomy, but that these stones can in general be managed by further ERCP.

Like Mr Thompson, we believe that laparoscopic bile duct exploration merits further study. In the introduction we state that "laparoscopic bile duct exploration is a well established technique but the time and skill required are likely to prevent it becoming universally available to patients." The experience of most surgeons with this technique is limited as supported by the relatively low numbers of patients reported in most series.⁵ The dilemma in many busy NHS hospitals is that unexpected bile duct exploration, either laparoscopic or open, is time consuming and interferes with the running of an already tight operating schedule. We therefore stand by our statement that currently "many clinicians therefore use preoperative ERCP and sphincterotomy." We are, however, enthusiastic supporters of minimally invasive therapy, particularly laparoscopic surgery, and would be interested in being involved in a comparative study of the various treatment options available.

G T DEANS
W A BROUGH
Stockport Unit for Minimally Invasive Therapy,
Stepping Hill Hospital,
Stockport SK2 7JE, UK

P SEDMAN
C M S ROYSTON
Hull Royal Infirmary, Hull, UK

D F MARTIN
South Manchester University Hospital,
Manchester, UK

C K LOEW
W E G THOMAS
Royal Hallamshire Hospital,
Sheffield, UK

Correspondence to: Mr Deans.

- 1 Welbourn CR, Mehta D, Armstrong CP, *et al*. Selective preoperative endoscopic retrograde cholangiography with sphincterotomy avoids bile duct exploration during laparoscopic cholecystectomy. *Gut* 1995;37:576–9.
- 2 Bonatsos G, Leandros E, Polydorou A, *et al*. ERCP in association with laparoscopic cholecystectomy. A strategy to minimize the number of unnecessary ERCPs. *Surg Endosc* 1996;10:37–40.
- 3 Bloom IT, Gibbs SL, Keeling-Roberts CS, *et al*. Intravenous infusion cholangiography for investigation of the bile duct: a direct comparison with endoscopic retrograde cholangiopancreatography. *Br J Surg* 1996;83:755–7.
- 4 Bergman JJ, van der Mey S, Rauws EA, *et al*. Long-term follow-up after endoscopic sphincterotomy for bile duct stones in patients younger than 60 years of age. *Gastrointest Endosc* 1996;44:643–9.
- 5 Drouard F, Passone-Szerzyńska N, Berthou JC. Laparoscopic treatment of common bile duct stones. *Hepatogastroenterology* 1997;44:16–21.

History of dyspepsia in patients with gastric cancer

EDITOR,—We read with interest the study by Suvakovic *et al* (*Gut* 1997;41:308–13) reporting that 63% of the patients with early gastric cancer had a previous history of upper gastrointestinal symptoms. These authors concluded that guidelines for the appropriateness of gastroscopy in patients with dyspepsia need to reflect the importance of not starting treatment prior to gastroscopy. General practitioners need to be made more aware of the potential for antisecretory drugs to mask early gastric cancer, which will then have a major detrimental impact on patient management. We have also studied whether patients with gastric cancer had serious and persistent dyspeptic symptoms necessitating consultation with their general practitioner (Laheij *et al*, unpublished observations). The results of our study, however, differ from Suvakovic *et al*'s.

We identified 46 patients with diagnostically verified early and late stage gastric cancer, using the Continuous Morbidity Register. This database is used to study the epidemiological aspects of diseases in primary care. The primary care population of approximately 12 000 patients has been followed since 1971. The recorded data have passed stringent quality controls and are consistent over the years of registration.¹ Controls were selected and matched for sex, social class, practice, observation period, and age.

The mean observation period between the date of registration and the first diagnosis of gastric malignancy was 12 years. In this period, 18 (39%) patients had dyspeptic symptoms necessitating consultation with the general practitioner compared with 20 of the controls (odds ratio 0.8, 95% confidence interval 0.3 to 2.0). Patients with gastric cancer had no more visits to their general practitioner for gastrointestinal symptoms than controls. Therefore, consultation rates for dyspeptic symptoms cannot be considered a warning for gastric cancer.

Every patient with gastric cancer develops gastrointestinal symptoms and there will always be a delay in diagnosis.² Martin *et al*

found a median delay of 17 weeks from the onset of symptoms to a definitive histological diagnosis in patients with gastric cancer. Use of an open access endoscopy service reduced this delay. There was no correlation between the delay in diagnosis and tumour stage or the success of potentially curative resection. The prognosis for gastric cancer is poor. To detect early gastric cancer, patients should be examined before the onset of dyspeptic symptoms. Fortunately, gastric cancer is rare in the Western world and epidemiological studies have shown that the incidence of and mortality from gastric cancer have decreased considerably. Screening people for gastric cancer before the onset of symptoms is therefore not feasible. Early detection of gastric cancer is not yet possible, and focusing attention on patients with a history of serious gastrointestinal symptoms may be of little value.

R J F LAHEIJ
J B M J JANSEN
Department of Gastroenterology,
University Hospital Nijmegen

E H VAN DE LISDONK
Department of General Practice and Social Medicine,
University of Nijmegen

A L M VERBEEK
Department of Medical Informatics, Epidemiology and
Statistics,
University of Nijmegen

Correspondence to: Dr R Laheij, MIES (152), PO Box 9101, 6500 HB Nijmegen, The Netherlands.

1 van Weel C. Validating long term morbidity recording. *J Epidemiol Community Health* 1995;49(suppl 1):29-32.

2 Martin IG, Young S, Sue-Ling H, et al. Delays in the diagnosis of oesophagogastric cancer: a consecutive case series. *BMJ* 1997;314:467-71.

Reply

EDITOR,—The points raised by Laheij and colleagues are relevant to the debate regarding the late diagnosis of gastric cancer in Western Europe. Their figure of 39% consulting their general practitioner over a mean time period of 12 years is not far away from our figure of 30% investigated for symptoms in the 13 years leading up to diagnosis. We are not suggesting that all of these patients had gastric cancer at the initial time of presentation, but many almost certainly did (most of those investigated in the last four years—a "miss" rate of 1 in 6). There are two important points to note. Firstly, general practitioners may not think of the correct diagnosis because of previous (reassuring) investigations with the result that the diagnosis is delayed by years not months. Secondly, and related to the first point, general practitioners will prescribe powerful acid suppression for patients with a previous ulcer history which will result in resolution of symptoms and a further delay in diagnosis. This raises the question when do general practitioners re-refer a patient for gastroscopy when previous investigations showed no pathology or benign disease? It is also important to remember that early gastric cancer has "benign" symptoms and so it is effectively too late when the patient has lost weight or become anaemic.

We disagree that gastric cancer cannot be diagnosed earlier and progress will only be made if we alter current practice to include investigating patients "at risk" before starting

antisecretory therapy and lowering the threshold for re-investigating patients who may have a history of dyspepsia. Screening is not feasible and we are not proposing such a policy. However within current practice we feel that there is room for improvement in the diagnosis of this disease which is not uncommon, representing the fourth commonest cancer in men in the UK.

Z SUVAKOVIC
M G BRAMBLE
Department of Gastroenterology,
Endoscopy Centre,
South Cleveland Hospital,
Marton Road,
Middlesbrough TS5 5AZ, UK

The genetics of ulcerative colitis based on one Swede

EDITOR,—I was interested to read the review on the role of inherited factors in inflammatory bowel disease by Satsangi *et al* (*Gut* 1997;40:572-4), but dismayed by the evidence provided. They claim that twin pairs provided strong evidence for genetic predisposition and that the most pertinent was that reported by Tysk *et al* using data from the Swedish Twin Registry.¹ However, the claim for ulcerative colitis rests on a single twin of the 50 000 registered. The probands were identified as hospitalised cases in Sweden. The 50 pairs of twins where one twin had suspected ulcerative colitis were whittled down to 36 pairs: 16 monozygotic and 20 dizygotic or of unknown zygosity. No hospitalised pairs were found. Twins of probands were interviewed and the occurrence of any gastrointestinal complaint was thoroughly discussed in order not to miss a mild non-hospitalised case. In spite of this approach only one twin of a proband could be claimed as a case of ulcerative colitis and he was one of the monozygotic group. This pair had only distal ulcerative colitis. From this one twin a heritability coefficient of 0.53 was calculated (with 95% confidence interval 0.24 to 0.82) and not 0.24 to 1.0 as stated by Satsangi *et al*. Thus, the 53% heritability claimed in both papers is based on one man with mild ulcerative colitis. The definition of his disease differed from that in the probands and the authors admit to deficiencies in their selection procedure. Tysk *et al* also admit that the prevalence of ulcerative colitis in the twins differs from that of the general population in Stockholm (mostly singletons). Twin studies cannot inform about disease in singletons if the prevalences are not the same. As the mean observation period was almost 20 years, it is unlikely that many more healthy twins will develop ulcerative colitis.¹ With 94% discordance in monozygotic twins, it is difficult to accept that most ulcerative colitis is inherited in either twins or singletons.

J J HARDING
Nuffield Laboratory of Ophthalmology,
University of Oxford,
Walton Street,
Oxford OX2 6AW, UK

1 Tysk C, Lindberg E, Jarnerot G, et al. Ulcerative colitis and Crohn's disease in an unselected population of monozygotic and dizygotic twins. A study of heritability and the influence of smoking. *Gut* 1988;29:990-6.

Reply

EDITOR,—We are grateful to Dr Harding for his comments, and we share his reservations regarding interpretation of twin studies in common diseases. Indeed, we have pointed out on a number of occasions that the genetic contribution in ulcerative colitis is less strong than that in Crohn's disease. Nevertheless, the model of disease inheritance which is most consistent with the epidemiological and molecular genetic data currently available is that Crohn's disease and ulcerative colitis are related polygenic disorders. The twin data are supported by epidemiological studies of familial prevalence from Scandinavia, the UK, and the USA. In recent years these epidemiological data have been supplemented by molecular genetic studies from a number of independent investigators. These studies have provided evidence for the presence of a number of genetic determinants of susceptibility and disease behaviour in ulcerative colitis. There is great optimism that these studies will result in an advance in the understanding of the pathogenesis of these common disorders. We strongly agree that both environmental as well as genetic factors are important in the pathogenesis of ulcerative colitis.

J SATSANGI
D P JEWELL
Gastroenterology Unit,
John Radcliffe Hospital,
Level 2,
Headington,
Oxford OX3 9DU, UK

Iron reduction therapy in hepatitis C

EDITOR,—Three papers in the July issue report on the possible association between iron and tissue damage in conditions other than haemochromatosis. Tan *et al* (*Gut* 1997;41:14-18) found that gastric cancer cells are more susceptible to photodynamic therapy when iron is removed. Boucher *et al* (*Gut* 1997;41:115-20) found that treatment of hepatitis C with interferon leads to a decrease in liver iron content. Bacon (*Gut* 1997;41:127-8) briefly commented on the association between iron and hepatitis C, including some evidence that iron depletion may be beneficial in patients who fail to respond to interferon α .

Shortly after Bacon *et al*'s pioneering report in 1993 on iron reduction therapy in hepatitis C,¹ and mainly because of a lack of any other option at the time, we started applying this form of therapy in our growing population of patients who had failed to respond to interferon and who had the unfavourable 1b genotype.

The simplest and cheapest way to reduce body iron stores is repeated drawing of one unit of whole blood (as for haemochromatosis). However, we encountered several unexpected difficulties in our attempts to implement a phlebotomy programme for patients with chronic hepatitis C. A small group of patients are reluctant to undergo repeated phlebotomy because of ethnic or psychological reasons (there is a belief that blood equals life and therefore that blood loss depletes the body of life giving power). A second group cannot tolerate blood loss because of recurrent episodes of faintness and presyncope. These patients need several hours' observation, a chaperone, and sometimes intravenous fluid before they can be discharged home. In a third group venous access is a problem. The