LETTERS TO THE EDITOR

Gall bladder motility after endoscopic sphincterotomy

EDITOR,—We read with interest the article by Sugiyama and Atomi (*Gut* 1996;39:856-9) on the effect of endoscopic sphincterotomy (ES) on gall bladder motility. These authors have proved conclusively that ES causes a significant improvement in gall bladder motility in humans. We have assessed gall bladder motility in patients who underwent ES for common bile duct (CBD) stones and compared these with patients with gall bladder calculi but without CBD stones. We found significantly lower fasting and residual gall bladder volumes in the patients who had undergone ES, indicating reduced filling after ES.¹ However, the ejection fraction and the rate constant of gall bladder emptying in these patients was increased significantly, suggesting increased gall bladder emptying. In another study² we found a significant decrease in fasting volume (mean (SD) 18.3 (8.5) v 13.9 (7.3) ml), residual volume (12.0 (8.0) v 4.4 (3.2) ml) and an increase in the ejection fraction (54.3 (9.8) v 83.5 (5.4)) after ES, suggesting decreased stasis and increased gall bladder emptying.

In Sugiyama and Atomi's study, all patients in whom gall bladder motility was assessed before ES had CBD stones; all stones were extracted before the motility studies were repeated. The major problem in this study is in differentiating between the effects of the CBD stones and the sphincterotomy itself. Fasting and residual volumes and maximum contraction were significantly different before and after ES. However, when one looks at these parameters in normal controls, it is clear that abnormal values before ES tend to normalise after ES, especially in patients with primary CBD stones. This strongly suggests that CBD stones adversely affect gall bladder motility. Common bile duct stones hinder gall bladder emptying by obstructing the lower CBD. In our study the patients had dilated CBD and some had multiple/large CBD stones, indicating significant obstruction. We circumvented this problem by including patients without significant CBD obstruction as reflected by normal serum bilirubin, a normal CBD diameter on cholangiography, and single/small CBD stones.³ Therefore, we believe that in this study improvement in gall bladder motility could be due to both ES and removal of the stones.

In Sugiyama and Atomi's study healthy volunteers were used as controls rather than patients with gallstones but without CBD stones. If gall bladder volumes and maximum contraction in patients with CBD stones were the same as in those with gallstones but without CBD stones, then one could conclude that the presence of CBD stones did not affect gall bladder emptying.

Patients with gallstones have small, contracted gall bladders because of associated chronic cholecystitis. However, in Sugiyama and Atomi's study fasting gall bladder volume was the same in patients with both gall- and CBD stones as in those with primary CBD

stones. Abnormal motility is not expected in the latter. Of the patients with both gall- and CBD stones, six of 15 patients had pigmented stones and would probably have had normal gall bladder motility. Behar and colleagues have shown that gall bladder muscle contractility is normal in patients with pigmented stones in comparison with patients with cholesterol stones who have reduced gall bladder muscle contractility.⁴ It is not clear whether Sugiyama and Atomi assessed gall bladder motility in a blinded fashion.

Sugiyama and Atomi mention that thickening of gall bladder wall after ES is due to reduced fasting volume. How a reduction in fasting volume can lead to thickening of the gall bladder wall is not clear. After ES, bactobilia occurs and duodenal contents can enter the gall bladder, which may lead to regurgitation cholecystitis and thickening of the gall bladder wall.⁵

Improvement in gall bladder motility after ES is due to abolition of papillary resistance. With division of the papilla, resistance is lost immediately and CBD pressure equalises to the duodenal pressure.⁶ This leads to a rapid improvement in gall bladder emptying as demonstrated by Sugiyama and Atomi on day 7 after ES. However why gall bladder emptying continued to improve for three months after ES was not explained. Despite a significant reduction in the size of the ES over five years, there was no evidence of a gradual decline after the initial improvement in gall bladder motility.

B C SHARMA
K SINGH

Department of Gastroenterology,
Postgraduate Institute of Medical Education and
Research,
Chandigarh 160 012, India

R K DHIMAN
Department of Hepatology

Correspondence to: Dr B C Sharma, 591, Sector 2, Panchkula, Haryana, India.

- Sharma BC, Singh K. Endoscopic sphincterotomy and gallbladder functions. *Scand J Gastroenterol* 1997;32:95-6.
- Dhiman RK, Phanish MK, Chawla YK, et al. Gallbladder motility and lithogenicity of bile in patients with choledocholithiasis after endoscopic sphincterotomy. *J Hepatol* 1997;26:1302-5.
- Agarwal DK, Sharma BC, Dhiman RK, et al. The effect of endoscopic sphincterotomy on gallbladder motility. *Dig Dis Sci* 1997;42:1495-500.
- Behar J, Lee KY, Thompson WR, et al. Gallbladder contraction in patients with pigment and cholesterol stones. *Gastroenterology* 1989;97:1479-84.
- Cohn MS, Schwartz SI, Faloon WW, et al. Effect of sphincteroplasty on gallbladder function and bile composition. *Ann Surg* 1979;189:317-21.
- Jensen PF, Csendes A, Kruse A, et al. Common bile duct and Oddi sphincter pressure before and after endoscopic papillotomy in patients with common bile duct stones. *Ann Surg* 1979;189:176-8.

Reply

EDITOR,—We appreciate the comments raised by Sharma *et al* about our paper. We agree that improvement in gall bladder motility after endoscopic sphincterotomy could be due to both decreased resistance of the terminal common bile duct and clearance of CBD stones, as we described in the discussion. We have found that patients with cholecystocholedocholithiasis have a lower maximum contraction than those with gallstones but without CBD stones (unpublished data). Therefore, the presence of the CBD stones

seems to affect gall bladder emptying. In our study, gall bladder function tests were carried out in a non-blinded manner.

Patients with severe cholecystitis have a small gall bladder. However, the patients had no or mild cholecystitis and a larger fasting gall bladder volume than normal controls.¹ Some patients with primary CBD stones may have mild cholecystitis or bile stasis, which may affect gall bladder motility. The brown pigmented stones in our study were composed of calcium bilirubinate and were different from black pigmented stones which mainly contain unconjugated bilirubin.^{2,3} Brown pigmented stones cause cholecystitis and gall bladder dysfunction.⁴

The gall bladder wall thickens as the gall bladder empties. Therefore, mild thickening of the gall bladder wall after sphincterotomy seems to be caused partly by the reduced fasting volume of the gall bladder. Regurgitation cholecystitis may also cause wall thickening, as Sharma *et al* state.

Improvement in gall bladder motility reached a plateau three months after sphincterotomy. Initial improvement seems to have been caused by decreased resistance of the terminal duct and stone clearance, as mentioned earlier. Further improvement after three months cannot be fully explained but may result partly from changes in bile composition.⁵

M SUGIYAMA
Y ATOMI

The First Department of Surgery,
Kyorin University School of Medicine,
6-20-2 Shinkawa,
Mitaka, Tokyo 181, Japan

Correspondence to: Dr Sugiyama.

- Spengler U, Sackmann M, Sauerbruch T, et al. Gallbladder motility before and after extracorporeal shock-wave lithotripsy. *Gastroenterology* 1989;96:860-3.
- Maki T. Pathogenesis of calcium bilirubinate gallstones; role of E. coli, beta-glucuronidase and coagulation by inorganic ions, polyelectrolytes and agitation. *Ann Surg* 1966;164:90-100.
- Trotman BW, Soloway RD. Pigment vs cholesterol cholelithiasis: clinical and epidemiological aspects. *Am J Dig Dis* 1975;20:735-40.
- Moriyasu A, Ise H, Kitayama O, et al. Evaluation of gallbladder contraction; influence of bile composition. *Japanese Journal of Biliopancreatic Physiology* 1995;11:29-33.
- Sauerbruch T, Stellaard F, Paumgartner G. Effect of endoscopic sphincterotomy on bile acid pool size and bile lipid composition in man. *Digestion* 1983;27:87-92.

Specificity of small bowel biopsy findings in coeliac disease

EDITOR,—I read Dr Metcalf's recent commentary (*Gut* 1998;42:9-10) on the association between coeliac disease and primary biliary cirrhosis, with much enthusiasm. Her editorial was both succinct and insightful. My concern relates to the author's suggestion that for coeliac disease, "endoscopic small bowel biopsy is specific." Unfortunately, this statement is incorrect. Specifically, a mucosal lesion identical to or closely resembling coeliac disease may be seen in patients with tropical sprue, lymphoma of the small bowel, Zollinger-Ellison syndrome, Crohn's disease, bacterial overgrowth in the small intestine, as well as multiple other disorders.¹ It is for this reason that it has been stressed that the histological appearance of the mucosa in coeliac disease is not specific, and alone cannot be considered diagnostic of this disorder. Importantly, in addition to a biopsy specimen

showing the "typical" coeliac sprue lesion, a clinical response to gluten withdrawal is required to diagnose the disorder unequivocally.

J C YARZE

Gastroenterology Associates of Northern New York,
5 Irongate Center,
Glens Falls, New York, 12801, USA

- 1 Trier JS. Celiac sprue and refractory sprue. In: Feldman M, Scharschmidt BF, Sleisenger MH, eds. *Gastrointestinal and liver disease. Pathophysiology, diagnosis, management*. Philadelphia: WB Saunders, 1998:1566–7.

Reply

EDITOR,—I thank Dr Yarze for his interest and kind comments. He is of course correct that small bowel biopsy is not 100% specific for coeliac disease taken in isolation. (Very few tests reach 100% specificity.) However, I hope he would agree that, in conjunction with a positive anti-gliadin antibody used as the initial screening test, the specificity is high. Another way of looking at this would be the positive predictive value of a small bowel biopsy, which depends upon the prevalence of the disease in the population being studied. As coeliac disease is relatively common among the Caucasian population affected by primary biliary cirrhosis, whereas the other cases of villous atrophy are rare among this group, small bowel biopsy would have a high positive predictive value.

J METCALF

Gloucestershire Royal Hospital,
Great Western Road,
Gloucester GL1 3NN, UK

Prevalence of the factor V Leiden mutation in portal and hepatic vein thrombosis

EDITOR,—We read with great interest the article by Mahmoud *et al* (*Gut* 1997;40:798–800) on the prevalence of factor V Leiden (FVL) mutation in hepatic and portal vein thrombosis. In the past few years the FVL mutation has emerged as the commonest genetic risk factor for venous thrombosis.¹ The FVL mutation seems to be most prevalent in Europe, extending to northern India in the east and Saudi Arabia in the south.² The two striking features of the population genetics of the FVL mutation are the confined racial distribution and the high prevalence in European peoples (allele frequencies ranging from 1.4 to 7%). We have found an allele frequency of 1.9% for the FVL mutation in 130 unrelated subjects from northern India who had no predisposing cause or family history of thrombosis.³

In India portal vein thrombosis (PVT) and hepatic vein thrombosis (Budd-Chiari syndrome) are frequently encountered hepatic disorders causing portal hypertension. In a series of patients with PVT from our institute the cause of disease could be elucidated in only 17 of the 213 cases.⁴ Umbilical sepsis led to PVT in 13 patients and congenital heart disease was responsible in four. Budd-Chiari syndrome represents a spectrum of disease primarily caused by a hypercoagulable state. Analysis of our cases of Budd-Chiari syndrome revealed the aetiology in 65 of the 177 patients.⁵ In four patients Budd-Chiari syndrome was associated with polycythaemia vera and in three with paroxysmal nocturnal haemoglobinuria. It has been well recognised that the clinical course, outcome and aetio-

pathogenic factors leading to PVT and Budd-Chiari syndrome are substantially different in the East and West.

As the FVL mutation has recently been detected in our population we attempted to study cases of PVT and Budd-Chiari syndrome to evaluate the role of this mutation in the aetiopathogenesis of these two conditions. Twenty three cases of PVT and nine cases of Budd-Chiari syndrome, including the five reported in our previous study,³ constituted the study group. The FVL mutation was detected as described by Bertina and colleagues.⁶ Genomic DNA was extracted from leucocytes by chloroform-phenol extraction, amplified using the polymerase chain reaction using specific primers, and the product was digested with the MnlI restriction enzyme. One of the 23 patients with PVT and two of the nine patients with Budd-Chiari syndrome were heterozygous for the FVL mutation. None of the patients was homozygous for the mutation. Another case of Budd-Chiari syndrome was associated with polycythaemia vera.

The prevalence of FVL mutation in our small series seems to be higher in patients with Budd-Chiari syndrome than in those with PVT or in controls. Therefore we concur with Mahmoud *et al*'s observation that screening for the FVL mutation is important in patients with Budd-Chiari syndrome. However, in cases of idiopathic PVT this mutation does not seem to have a major pathogenic role.

R DAS

G GAREWAL

Department of Haematology,
Postgraduate Institute of Medical Education
and Research,
Chandigarh 160 012, India

Y CHAWLA

R K DHIMAN

Department of Hepatology

Correspondence to: Professor G Garewal.

- 1 Dahlback B, Carlsson M, Svensson PJ. Familial thrombophilia due to a poor unrecognised mechanism characterized by poor anticoagulant response to activated protein C: prediction of a cofactor to activated protein C. *Proc Natl Acad Sci USA* 1993;90:1004–8.
- 2 Rees DC. The population genetics of factor V Leiden (Arg 506 Gln). *Br J Haematol* 1996;95:579–86.
- 3 Garewal G, Das R, Trehan U. Factor V Leiden: prevalence in the indigenous population and cases of thrombosis in north India. *Br J Haematol* 1997;97:940.
- 4 Dilawari JB, Chawla YK. Extrahepatic portal venous obstruction. *Gut* 1988;29:554–5.
- 5 Dilawari JB, Bamberg P, Chawla YK, *et al*. Hepatic outflow obstruction (Budd-Chiari syndrome): experience with 177 patients and a review of the literature. *Medicine* 1994;73:21–36.
- 6 Bertina RM, Koeleman BPC, Koster T, *et al*. Mutation in blood coagulation factor V associated with resistance to activated protein C. *Nature* 1994;369:64–7.

Steatorrhoea and pancreatic disease in HIV infected patients

EDITOR,—The study by Carbonnel *et al* (*Gut* 1997;41:805–10) on macronutrient intake and steatorrhoea in HIV infected patients tackles one important clinical feature of HIV infection. On admission to hospital up to 60% of patients with AIDS are underweight (>10% weight loss).¹ Even asymptomatic patients at an early stage of disease have changes in body composition as shown by a reduction in body cell mass, which is associated with an increased mortality and

morbidity.² HIV associated malnutrition has a multifactorial origin but as shown by Carbonnel *et al* and other authors steatorrhoea is a common problem in HIV infected patients, especially in those with opportunistic infections of the gastrointestinal tract.³ Inadequate energy intake is a major determinant of malnutrition in these patients, which is confirmed by the fact that enteral refeeding can improve their nutritional status.⁴

Unfortunately Carbonnel *et al*'s study missed one important aspect of steatorrhoea—next to decreased fat absorption, exocrine pancreatic insufficiency may also result in lipid malabsorption. Raised pancreatic serum enzyme concentrations are frequently seen in patients with AIDS and also to a lower extent in asymptomatic HIV infected patients, suggesting the presence of pancreatic inflammation.⁵ A recently published study reported that 54% of patients with suspected AIDS related cholangitis who underwent endoscopic retrograde cholangiopancreatography (ERCP) had a pathological pancreaticography (I°, 26%; II°, 47%; III°, 27% classified according to the Cambridge classification).⁶ Pathological pancreatographies were associated with opportunistic infections (candida, cytomegalovirus, cryptosporidia, microsporidia, and mycobacteria) and a CD4 count of less than 60/mm³. In the study by Carbonnel *et al* 60% of the patients with diarrhoea were infected with cryptosporidia or microsporidia and the mean CD4 count was 16/mm³, therefore these patients have an increased risk of pancreatic involvement. To our knowledge no studies have been published on exocrine pancreatic function in HIV infected adult patients. In a small preliminary study in HIV infected children five of 17 patients had pancreatic dysfunction and three of five had significant steatorrhoea.⁷

Further investigations of lipid malabsorption and weight loss in HIV infected patients, especially in those with advanced disease and cryptosporidial or microsporidial infection, should include a diagnostic work up of the exocrine pancreas. This is important because these patients may benefit from pancreatic enzyme supplementation.

J OCKENGA

M GÖKE

M P MANN

Department of Gastroenterology and Hepatology,
Medical School Hannover,
30623 Hannover, Germany

Correspondence to: Dr J Ockenga
(email: ocken@aol.com).

- 1 Nahlen BL, Chu SY, Nwanyanwu C, *et al*. HIV wasting syndrome in the United States. *AIDS* 1993;7:183–8.
- 2 Suttman U, Ockenga J, Selberg O, *et al*. Incidence and prognostic value of malnutrition and wasting in human immunodeficiency virus-infected outpatients. *J Acquir Immune Defic Syndr* 1995;8:239–46.
- 3 Sharpstone D, Gazzard B. Gastrointestinal manifestations of HIV infection. *Lancet* 1996;348:379–83.
- 4 Ockenga J, Suttman U, Selberg O, *et al*. Percutaneous endoscopic gastrostomy in AIDS and control patients: risk and outcome. *Am J Gastroenterol* 1996;91:1817–22.
- 5 Dutta SK, Ting CD, Lai LL. Study of prevalence, severity, and etiological factors associated with acute pancreatitis in patients infected with human immunodeficiency virus. *Am J Gastroenterol* 1997;92:2044–8.
- 6 Barthet M, Chauveau E, Bonnet E, *et al*. Pancreatic ductal changes in HIV-infected patients. *Gastrointest Endosc* 1997;45:59–63.
- 7 Carroccio A, Fontana M, Guarino A, *et al*. Exocrine pancreatic function in HIV-infected children: evaluation with non-invasive tests. *Digestion* 1997;58(suppl 2):45.