this study. The data are retrospective and incomplete and based on the study of only 25 patients (the under 55 group) out of 319 with gastric cancer. There is no comparison of this group with the remaining 296 patients over 55 with regard to pattern of presentation or symptoms.

Perhaps more importantly there is no mention of the number of patients who actually presented for endoscopy or were picked up on the open access service. Surely the only way to set protocols for this service would be to analyse the data from it. In contrast this study seems merely to describe 25 patients below the age of 55 presenting in the Gloucester region, drawn from a pathology database, most of whom had advanced gastric cancers.

We would hope that better awareness of the importance of early referral and increased use of diagnostic endoscopic facilities should result in more patients of all ages presenting with early disease. Based on our experience¹ and that of centres such as Leeds,² we would advocate open access endoscopy in anyone with new dyspeptic symptoms over the age of 40.

Finally and perhaps most worryingly the authors state in their discussion that early detection does not necessarily mean improved survival. Those of us involved in the treatment of gastric cancer realise that the only significant factor that is going to improve survival from this disease is early detection and treatment. There is overwhelming and irrefutable evidence to support this. In the UK the detection and treatment of early gastric cancers has led to a five year survival rate of over 90% in these patients.³ In Japan, where mass radiological screening of the over 40s, ready access to endoscopy and population awareness of the disease has meant detection of early gastric cancer in more than half of all gastric cancer cases and again a five year survival of over 90% in these patients.⁴ Early detection *does* mean improved survival.

Until we have adequate prospective data from a large open access endoscopy unit we cannot agree with the interpretation and findings of this study and urge other centres to continue to endoscope symptomatic patients under the age of 55.

D KARAT
J WAYMAN
N HAYES
S A RAIMES
S M GRIFFIN

Northern Oesophago-gastric Cancer Unit,
Royal Victoria Infirmary,
Queen Victoria Road,
Newcastle upon Tyne NE1 4LB, UK

- 1 O'Hanlon DM, Karat D, Hayes N, *et al*. Open access endoscopy facilitates early diagnosis of malignant and pre-malignant lesions of the stomach and oesophagus [abstract]. *Gut* 1995;37(suppl 2):A35.
- 2 Sue-Ling HM, Johnston D, Martin IG, *et al*. Gastric cancer: a curable disease in Britain. *BMJ* 1993;307:591-6.
- 3 Sue-Ling HM, Martin I, Griffiths J, *et al*. Early gastric cancer; 46 patients treated in one surgical department. *Gut* 1992;33:1318-22.
- 4 Hisamichi S. Screening for gastric cancer. *World J Surg* 1989;13:31-7.

Reply

EDITOR,—Most of the arguments raised by Karat and colleagues were covered in our paper. However, because we consider that the epidemiological principles underlying our paper are important, we think it is worth reiterating them.

The data are retrospective but there are no comparable prospective data. Unlike other

studies ours was of a defined population based on postcodes. The importance of this for determining presentation characteristics and natural history of disease cannot be over-emphasised. Most other studies emanate from referral centres that receive selected patients. The resulting referral bias invariably influences the type of patient seen and interpretations made. Thus we believe our study is a better reflection of the real world.

We went to great lengths to ensure completeness (e.g. searching several databases) of patient capture. Our incidences are comparable to OPCS data and therefore we think it unlikely that many, if any, patients were missed. The small number (25 in seven years) of patients aged less than 55 just emphasises how rare the disease is in this age group.

Karat *et al* advocate better awareness of the importance of early referral to improve the chance of finding early disease. A point we stressed in our paper was that referral by general practitioners and subsequent investigation were not significantly delayed.

We did not determine the source of patients (clinic or open access) because general practitioners use clinic referrals rather than open access endoscopy for all sorts of reasons unrelated to symptoms. There would be too many confounders to make a meaningful exploration of differences in these groups. The point about the open access service in Gloucester is that it has been in operation for 20 years. The local doctors are relatively experienced in its use. The experience of its effects in the **past** 10 years in our district is relevant to experience in the **next** 10 years in other districts (the majority) that have introduced an open access service more recently.

In the UK there is no "overwhelming and irrefutable evidence" that early detection improves survival from gastric cancer. Only a randomised controlled trial can give this level of certainty and none has been done in the UK. The case series quoted from Leeds was not based on a geographically defined population and it is subject to the biases of all screening and early detection studies: lead and length bias.

We can sympathise with the sentiments of Karat *et al*: watching patients with gastric cancer die when intuitively you believe an earlier diagnosis would have helped is not easy. However, we would urge them to consider the following questions. Is there evidence that patients with curable gastric cancer are more likely to have uncomplicated dyspepsia than age and sex matched controls? If not, why investigate on the basis of symptoms? What opportunities are missed (e.g. early detection of colorectal cancer) if a huge amount of diagnostic resource is directed towards detecting the small number (<200 per annum) of gastric cancers in the under 55s. Could we use our limited financial resource in a more effective way?

R VALORI
N A SHEPHERD
Gloucestershire Royal Hospital,
Great Western Road,
Gloucester GL1 3NN, UK

Activation of NFκB in inflammatory bowel disease

EDITOR,—In their recent paper (*Gut* 1998;42:477-84) Schreiber and coworkers demonstrated that RelA (p65) is present in nuclear extracts of biopsy specimens or

lamina propria mononuclear cells from patients with active inflammatory bowel disease (IBD). Furthermore, they show NFκB binding activity and a corresponding decrease in IκBα in lamina propria mononuclear cells treated with lipopolysaccharide (LPS). In contrast, treatment with dexamethasone prevented LPS induced nuclear translocation of NFκB due to persistence of IκBα. The authors conclude from these data that corticosteroids inhibit NFκB activation *in vitro* by stabilising the cytosolic inhibitor IκBα against activation induced degradation.

Firstly, this conclusion cannot be drawn from the data presented in the paper. Secondly, their conclusion contradicts a number of previously published observations.

Two models of corticosteroid mediated NFκB inhibition have been proposed. The first proposes that down modulation of κB driven genes results from a physical interaction between the glucocorticoid receptor and the RelA (p65) subunit. Negative cross-talk between the glucocorticoid receptor and RelA is due to direct interaction via the Rel homology domain of RelA and the DNA binding domain of the glucocorticoid receptor in combination with interference by the transactivation domain of RelA with the transcriptional activity of the glucocorticoid receptor.¹⁻⁵

The second model proposes that the inhibitory effect of glucocorticoids is mediated by the induction of the IκBα protein, which traps activated NFκB in inactive cytoplasmic complexes. It has been shown that dexamethasone induces transcriptional upregulation of the IκBα gene, although a glucocorticoid responsive element has not been identified in the IκBα promoter. Thus, in the presence of dexamethasone, released NFκB quickly reassociates with newly synthesised IκBα which results in notably reduced amounts of NFκB that translocates to the nucleus.⁶⁻⁷

So far there is no evidence that steroids stabilise IκBα. However, we have recently shown that sulphasalazine interferes with IκBα phosphorylation and proteasome dependent degradation of IκBα *in vitro*.⁸ Therefore, the available data point towards different levels of interference with NFκB activation by sulphasalazine and corticosteroids.

R M SCHMID
G ADLER
Department of Internal Medicine I,
University of Ulm, Ulm, Germany
S LIPTAY
Department of Paediatrics,
University of Ulm, Ulm, Germany

Correspondence to: Correspondence to: Dr Schmid, Department of Internal Medicine I, University of Ulm, Robert-Koch-Strasse 8, 89081 Ulm, Germany.

- 1 Caldenhoven E, Liden J, Wissink S, *et al*. Negative cross-talk between RelA and the glucocorticoid receptor: a possible mechanism for the anti-inflammatory action of glucocorticoids. *Mol Endocrinol* 1995;9:401-12.
- 2 Ray A, Prefontaine KE. Physical association and functional antagonism between the p65 subunit of transcription factor NF-κappa B and the glucocorticoid receptor. *Proc Natl Acad Sci USA* 1994;91:752-6.
- 3 Scheinman RI, Gualberto A, Jewell CM, *et al*. Characterization of mechanisms involved in transrepression of NF-κappa B by activated glucocorticoid receptors. *Mol Cell Biol* 1995;15:943-53.
- 4 Wissink S, van Heerde EC, Schmitz ML, *et al*. Distinct domains of the RelA NF-κB subunit are required for negative cross-talk and direct

- interaction with the glucocorticoid receptor. *J Biol Chem* 1997;272:22278–84.
- 5 De Bosscher K, Schmitz ML, van den Berghe W, *et al.* Glucocorticoid-mediated repression of nuclear factor- κ B-dependent transcription involves direct interference with transactivation. *Proc Natl Acad Sci USA* 1997;94:13504–9.
 - 6 Scheinman RI, Cogswell PC, Lofquist AK, *et al.* Role of transcriptional activation of I κ B α in mediation of immunosuppression by glucocorticoids. *Science* 1995;270:283–6.
 - 7 Auphan N, DiDonato JA, Rosette C, *et al.* Immunosuppression by glucocorticoids: inhibition of NF κ B activity through induction of I κ B synthesis. *Science* 1995;270:286–90.
 - 8 Wahl C, Liptay S, Adler G, *et al.* Sulfasalazine: a potent and specific inhibitor of nuclear factor kappa B. *J Clin Invest* 1998;101:1163–74.

Reply

EDITOR.—We appreciate the comments by Schmid and colleagues which address important aspects of the cellular mechanisms involved in the in vitro regulation of NF κ B activation. Our sequential publications in *Gut* have tackled the clinical importance of the in vivo activation of the proinflammatory transcription factor NF κ B in inflammatory bowel disease (IBD). We have also shown the importance of human granulocytes in primary cultures as specific contributors to this process.^{1,2} These studies were not designed to investigate the cellular mechanism of steroid action in the NF κ B system, but rather to look at important clinical aspects of NF κ B activation and steroid treatment in Crohn's disease.

An important anti-inflammatory mechanism in steroid action is the inhibition of NF κ B activation.^{3–5} It has been suggested that this mechanism involves the induction of I κ B α expression, the stabilisation of I κ B α (resulting in sequestration of NF κ B in the cytoplasm) and protein–protein interactions between NF κ B and the glucocorticoid receptor, leading to inhibition of transactivation.⁵

The exact mechanisms leading to the steroid induced persistence of I κ B α protein are not established. Although some experiments have shown upregulation of the expression of I κ B α by steroids, as those cited by Schmid *et al.*, these findings have not been reproduced in more recent studies.^{6–8} In addition, the glucocorticoid-like immunosuppressive effect of oestrogen as well as other steroid hormones is also attributed to stabilisation of the I κ B α protein as a result of inhibition of I κ B α degradation.⁹ The kinetics of I κ B α degradation in human granulocytes are very rapid and steroid induced protection of the I κ B α protein is seen within minutes. One may therefore speculate how much induction of expression and how much stabilisation against degradation may contribute to the rapid time course in granulocytes. The relative contribution of these mechanisms to the clinical efficacy of steroids in inhibiting activation of NF κ B needs to be evaluated in future studies.

Taken together, the understanding of the role of I κ B α in steroid induced suppression of NF κ B activation is hampered by conflicting findings which may partly result from the stimulatory conditions chosen. The choice of the cell system (i.e. epithelial carcinoma cell lines which often have an altered I κ B α /NF κ B system versus primary cells¹⁰) may also confound these data.

Schmid *et al.* also discuss the mechanism of the anti-inflammatory action of sulphasalazine.¹¹ They propose that treatment with sulphasalazine results in the inhibition of I κ B α degradation, whereas aminosaliclates (5-ASA) or sulphapyridin have no effect on

NF κ B activation.¹¹ The authors' findings, which have been obtained in colonic carcinoma (SW620) and Jurkat T cell lines, add an important aspect to the mechanisms of action of anti-inflammatory drugs in IBD but also underline the importance of the experimental system investigated as discussed earlier. In other systems (e.g. primary monocyte cultures) aminosaliclates also seemed to inhibit the activation of NF κ B via inhibition of I κ B α degradation.^{12–18}

Activation of NF κ B may be a central element in the pathophysiology of Crohn's disease. Inhibition of NF κ B activation offers an attractive hypothesis for the action of numerous drugs with clinically relevant effects in IBD. The exact mechanisms involved have yet to be determined in the relevant cell systems and in further ex vivo studies using affected tissues of diseased patients.

S SCHREIBER
S NIKOLAUS
J HAMPE

First Medical Department,
Christian Albrechts University,
Schittenhelmstrasse 12,
24105 Kiel, Germany
email: s.schreiber@mucosa.de

- 1 Schreiber S, Nikolaus S, Hampe J. Activation of nuclear factor κ B in inflammatory bowel disease. *Gut* 1998;42:477–84.
- 2 Nikolaus S, Bauditz J, Gionchetti P, *et al.* Increased secretion of pro-inflammatory cytokines by circulating polymorphonuclear neutrophils and regulation by interleukin 10 during intestinal inflammation. *Gut* 1998;42:470–6.
- 3 Ray A, Prefontaine KE. Physical association and functional antagonism between the p65 subunit of transcription factor NF- κ B and the glucocorticoid receptor. *Proc Natl Acad Sci USA* 1994;91:752–6.
- 4 Scheinman RI, Cogswell PC, Lofquist AK, *et al.* Role of transcriptional activation by I κ B in mediation of immunosuppression by glucocorticoids. *Science* 1995;270:283–6.
- 5 Wissink S, van Heerde EC, van der Burg B, *et al.* A dual mechanism mediates repression of NF- κ B activity by glucocorticoids. *Mol Endocrinol* 1998;12:355–63.
- 6 Brostjan C, Anrather J, Cizmadi V, *et al.* Glucocorticoid-mediated repression of NF κ B activity in endothelial cells does not involve induction of I κ B synthesis. *J Biol Chem* 1996;271:19612–16.
- 7 De Bosscher K, Schmitz ML, van den Berghe W, *et al.* Glucocorticoid-mediated repression of nuclear factor- κ B-dependent transcription involves direct interference with transactivation. *Proc Natl Acad Sci USA* 1997;94:13504–9.
- 8 Heck S, Bender K, Kullmann M, *et al.* I κ B α -independent downregulation of NF- κ B activity by the glucocorticoid receptor. *EMBO J* 1997;16:4698–707.
- 9 Sun WH, Keller ST, Stebler BS, *et al.* Estrogen inhibits phorbol ester-induced I κ B α transcription and protein degradation. *Biochem Biophys Res Commun* 1998;244:691–5.
- 10 Jobin C, Haskill S, Mayer L, *et al.* Evidence for altered regulation of I κ B α degradation in human colonic epithelial cells. *J Immunol* 1997;158:226–34.
- 11 Wahl C, Liptay S, Adler G, *et al.* Sulfasalazine: a potent and specific inhibitor of nuclear factor kappa B. *J Clin Invest* 1998;101:1163–74.
- 12 Kopp E, Ghosh S. Inhibition of NF- κ B by sodium salicylate and aspirin. *Science* 1994;265:956–9.
- 13 Bitko V, Velazquez A, Yang L, *et al.* Transcriptional induction of multiple cytokines by human respiratory syncytial virus requires activation of NF- κ B and is inhibited by sodium salicylate and aspirin. *Virology* 1997;232:369–78.
- 14 Stange C, Ramirez I, Gomez I, *et al.* Phosphorylation of nuclear proteins directs binding to salicylic acid responsive elements. *Plant J* 1997;11:1315–24.
- 15 Oeth P, Mackman N. Salicylates inhibit lipopolysaccharide-induced transcriptional activation of the tissue factor gene in human monocytic cells. *Blood* 1995;86:4144–52.
- 16 Takashiba S, Van Dyke TE, Shapira L, *et al.* Lipopolysaccharide-inducible and salicylate,

sensitive nuclear factor (s) on human tumor necrosis factor alpha promoter. *Infect Immun* 1995;63:1529–34.

- 17 Pierce JW, Read MA, Ding H, *et al.* Salicylates inhibit I κ B α phosphorylation, endothelial-leukocyte adhesion molecule expression, and neutrophil transmigration. *J Immunol* 1996;156:3961–9.
- 18 Schwenger P, Alpert D, Slotnik EY, *et al.* Activation of p38 mitogen-activated protein kinase by sodium salicylate leads to inhibition of tumor necrosis factor-induced I κ B α phosphorylation and degradation. *Mol Cell Biol* 1998;18:78–84.

BOOK REVIEWS

Pancreatitis. Lankisch PG, Banks PA. (Pp 377; illustrated; £57.00.) Berlin: Springer-Verlag, 1998. ISBN 3-540-61726-4.

This book will be a bench mark by which others on the subject of pancreatitis will be judged. This is no ordinary volume assembled by two hard-pressed editors from an array of international experts with differing styles and experience but a text written solely by two eminent authors. The gestation period was 10 years, the evidence base is 1900 citations and the bias is distinctly clinical—that is, this is a reference book for the working gastroenterologist (surgeon or physician), not a text littered with incomprehensible molecular biology.

The two diseases, acute and chronic pancreatitis, are dealt with in a similar format, beginning with classification, aetiology, pathology, and epidemiology, progressing to the clinicians' holy grail: diagnosis and treatment. Herein lies the book's strength. There is no artificial segregation into separate sections by a radiologist, an endoscopist or a surgeon, but an integrated and detailed overview giving practical guidance. For two disease processes whose pathogenesis is poorly understood and for which there are few specific therapies, it is obviously necessary for guidance to be more than anecdotal. Thus, where there is controversy the literature has been carefully sifted through, debated and a bottom line recommendation reached. The approach is not nihilistic, but thoroughly positive in its analysis of international efforts to solve the conundrum of pancreatitis.

So what gems can be extracted to whet your appetite? Firstly a riveting discussion of scoring systems and prognostic signs in acute pancreatitis, a topic which often switches off the relative newcomer to this area but which holds the key to stratification for current or new treatment options. Ranson, Imrie and APACHE II take on a new life, liberally illustrated with explanatory figures and tables. Secondly, there is a clear and detailed discussion of the merits and practical value of the various pancreatic function tests which can be used in chronic pancreatitis. When to use each one, sensitivity and specificity, and practical tips bring clarity to an often muddled area. And on the back of this come recommendations for intervention and treatment. In my opinion the chapters on treatment cannot be bettered. Finally, although the authors are both physicians the surgical options available are debated sensibly and concisely and placed in a realistic perspective.