moner in age/sex matched alcoholics with either of the haemochromatosis gene mutations. However, although not quite reaching statistical significance, in this population twice as many alcoholics with one copy of the C282Y mutation did have grade >2 hepatocyte iron compared with those without this mutation (36% v 18%) and twice as many alcoholics with one copy of the C282Y mutation had any degree of histological iron compared with those without the C282Y mutation (46% v 24%).

In order to eliminate the effect of cirrhosis per se on hepatic iron loading it would be interesting to know whether there was any greater association between the *HFE* gene mutations and hepatic iron stores in the 59 patients with precirrhotic alcoholic liver disease. Perhaps more importantly it would be interesting to know in these 59 patients whether there was any relation between hepatic iron stores and the degree of hepatic fibrosis.

The fact that the *HFE* gene mutations were not significantly commoner in patients with fibrotic/cirrhotic alcoholic liver disease than controls does seem to suggest that these mutations do not have a major role in the progression of alcoholic liver disease. However, it is likely that any effect that *HFE* might have on the progression of chronic liver disease is mediated through raising iron stores, as we have shown in non-alcoholic steatohepatitis,³ rather than a direct effect of *HFE* on fibrogenesis. Only a minority of heterozygotes show some phenotypic expression of iron overload,⁴ other causes of iron overload clearly occur^{5,6} and many factors may be involved in the progression of alcoholic liver disease. Therefore a much larger fibrotic alcoholic population may be required to investigate definitively whether or not the *HFE* gene mutations are enriched in this population.

We agree with Grove *et al* that further research is required to investigate the factors that determine liver iron content in alcoholic liver disease and the prognostic significance of this iron. We do not think, however, that the effect of the haemochromatosis gene mutations on liver iron content and disease progression can yet be completely discounted.

D K GEORGE
Gartnavel General Hospital,
1053 Great Western Road,
Glasgow G11 6NT, UK

L W POWELL
Clinical Sciences Unit,
Queensland Institute of Medical Research,
300 Herston Road,
Brisbane, Queensland, Australia

Correspondence to: Dr George.

- 1 Ludwig J, Hashimoto E, Porayko MK, *et al*. Hemosiderosis in cirrhosis: A study of 447 native livers. *Gastroenterology* 1997;112:882–8.
- 2 Deugnier YM, Olivier L, Turlin B, *et al*. Liver pathology in genetic hemochromatosis: a review of 135 homozygous cases and their biochemical correlations. *Gastroenterology* 1992;102:2050–9.
- 3 George DK, Goldworn S, Macdonald GA, *et al*. Increased hepatic iron stores in non alcoholic steatohepatitis are associated with increased liver fibrosis. *Gastroenterology* 1998;114:311–19.
- 4 Bulaj ZL, Griffen LM, Jorde LB, *et al*. Clinical and biochemical abnormalities in people heterozygous for hemochromatosis. *N Engl J Med* 1996;335:1799–805.
- 5 Moirand M, Mortaji AM, Loreal O, *et al*. A new syndrome of liver iron overload with normal transferrin saturation. *Lancet* 1997;349:95–7.

6 The UK Haemochromatosis Consortium. A simple genetic test identifies 90% of UK patients with haemochromatosis. *Gut* 1997;41:841–4.

Reply

EDITOR,—We thank Drs George and Powell for their comments on our paper. We agree that confining our iron studies to the assessment of iron staining in hepatocytes of cirrhotic patients may have led to underestimation of the effect of the *HFE* genotype on iron content in alcoholic liver disease. However, as they point out, our results are fully consistent with the findings in non-alcoholic C282Y heterozygotes—only a minority of whom show evidence of iron overload—and any small effect of C282Y heterozygosity on liver iron content is likely to be swamped by the other causes of iron overload present in heavy drinkers. On the question of whether or not heavy drinkers with the C282Y mutation have more liver iron than drinkers without the mutation, in contrast to George and Powell, we believe our results of *HFE* genotyping do largely exclude a role for this mutation in determining the risk of fibrotic alcoholic liver disease. This is based not only on the similar frequency of the mutation in patients and controls, but also the identical mean age of presentation (and therefore cumulative alcohol intake) in patients with and without the mutation. Moreover, the lack of any increased frequency of the C282Y mutation in patients with alcoholic liver disease compared with controls has been reported by others.¹

With respect to the relation between iron stores and the degree/risk of fibrosis in heavy drinkers, we have no data on iron staining in the patients with fibrosis. However, we have reported recently that the liver iron content of patients with alcoholic fatty liver does not predict the risk of progression to fibrosis or cirrhosis.² If we accept that the C282Y mutation increases liver iron content but not the risk of alcoholic liver disease, taken together, these results would appear largely to exclude a role for excess liver iron in the pathogenesis of alcohol related liver disease.

C P DAY
Department of Medicine,
The Medical School,
Framlington Place,
Newcastle upon Tyne NE2 4HH, UK

- 1 Aldersley MA, Howdle PD, Wyatt JL, *et al*. Haemochromatosis gene mutation in liver disease patients. *Lancet* 1997;349:1025–6.
- 2 Day CP, James OFW, Guzail M, *et al*. Iron content of alcoholic fatty liver does not predict progression to alcoholic liver disease. *J Hepatol* 1998;28(suppl 1):131.

Ileal pouch-anal anastomosis for Crohn's disease

EDITOR,—Mr Phillips (*Gut* 1998;43:303–4) enjoys stirring the pot, but we cannot support all of his views on ileo-anal pouch surgery for Crohn's disease. Firstly, we do not think ulcerative colitis and Crohn's disease are necessarily "apples and oranges" (though familial adenomatous polyposis is). We believe that there is often a spectrum of disease and that in some cases only the long term natural history enables clinicians to know whether they are dealing with one or other disorder.

Also, we do not think it is appropriate to compare the outcome of Kock pouch surgery against pelvic pouch surgery for Crohn's disease. If there is a stricture in or above a pouch

which is out of the pelvis, the chances of successful reconstruction without pouch excision is much greater than for pouches in the pelvis (as our own experience confirms).

To confuse matters further, Mr Phillips compares colectomy and ileo-rectal anastomosis with pouch-anal anastomosis for Crohn's disease. This certainly introduces an "apples and oranges" scenario as ileo-rectal anastomosis is selective because of rectal sparing and, furthermore, even quiescent perianal disease or ileal involvement is not generally a contraindication. By contrast, ileo-anal pouches are not advised if there is perianal or small bowel disease.

Much of today's literature highlights a 10% change in diagnosis from ulcerative colitis to Crohn's disease over 10 years.¹ In some cases, as in our earlier experience,² the initial pathological scrutiny was not sufficiently rigorous and the patient had Crohn's disease from the start, whereas in others, small bowel disease develops after pouch construction and the diagnosis has to be revised. To advise pouch construction for known Crohn's proctocolitis when there is no anal or small bowel disease was heretical in the past and, still, is rarely advised. If a patient does not want a stoma a lot of counselling is needed if a surgeon proceeds with the patient's wishes. Admittedly, in our experience, just under half have a reasonable functional result over 10 years. However, for those that fail, morbidity is high with many more hospital admissions, a much greater length of small bowel sacrificed and a far higher risk of perineal sinus than in patients having proctectomy after a failed ileo-rectal anastomosis.

M R B KEIGHLEY
R N ALLAN
D S A SANDERS
Queen Elizabeth Hospital,
Edgbaston,
Birmingham B15 2TH, UK

- 1 Marcello P, Schoetz D, Roberts P, *et al*. Evolutionary changes in pathologic diagnosis after the ileoanal pouch procedure. *Dis Colon Rectum* 1997;40:263–9.
- 2 Grobler S, Hosie K, Affie E, *et al*. Outcome of restorative proctocolectomy when the diagnosis is suggestive of Crohn's disease. *Gut* 1993;34:1384–8.

Reply

EDITOR,—I think the readership will need to decide whether "apples and oranges" have been compared in the past. The leading article simply tried to point out the huge differences between ulcerative colitis and Crohn's disease and queried whether comparisons were fair, or even reasonable. After all, ulcerative colitis is curable by surgery, it has a reasonably low surgical morbidity and good prospects. Conversely, Crohn's disease is not cured by surgery, there is a recognised high surgical morbidity, all anastomoses are prone to recurrent disease, and reoperation is frequent.

As far as disease spectrum is concerned, Keighley *et al* may be right, although another possibility, partly supported by their own final paragraph, is that Crohn's disease may arise in patients with ulcerative colitis just as arthritis, heart disease or asthma may arise in these patients.

The rationale of alluding to Kock pouches was not to indicate that they are exactly equivalent to an ileo-anal pouch, but rather to show that when Kock pouches were made in

Crohn's disease, the results were nothing like as bad as many people currently assume.

The purpose of introducing colectomy and ileo-rectal anastomosis (IRA) was to make the point that, by comparison with proctocolectomy and ileostomy, reoperation is frequent, lengths of bowel are lost and short bowel syndrome is rare and yet it is an acceptable treatment modality. Keighley *et al* assist this comparison by indicating that, despite rectal sparing, patients undergoing IRA still do badly when compared with patients undergoing total excision and an ileostomy, the latter generally having worse disease. Obviously the issue here is that all anastomoses in Crohn's disease are prone to recurrence and yet the medical community accepts the recurrences as a reasonable trade off when the alternative is a permanent stoma.

In the leading article I tried to indicate that excluding ileo-anal pouch construction as one choice in carefully selected and counselled patients with Crohn's disease who have neither anal nor small bowel disease was neither logical nor consistent with the published results of pouches made, usually inadvertently, in Crohn's disease.

R S K PHILLIPS
Consultant Surgeon and Dean,
St Mark's Academic Institute,
Harrow HA1 3UJ, UK

Excess nitric oxide in ulcerative colitis may be generated by nitric oxide synthase independent pathways

EDITOR.—Kimura *et al* (*Gut* 1998;42:180–7) provide further data in support of the proposal that production of nitric oxide (NO) in ulcerative colitis (UC) is ascribable primarily to the action of inducible NO synthase (NOS) on L-arginine.^{1–3}

This reaction relies on molecular oxygen as a substrate (fig 1).⁴ However, the P_{O_2} of gas in the normal colon is low (sometimes only 1–2 mm Hg), not only as a result of bacterial utilisation of oxygen, but also reflecting mucosal hypoxia as a result of cellular metabolism and counter current diffusion of oxygen between mucosal capillaries.⁵ In active UC, mucosal ischaemia and conversion of oxygen to reactive oxygen metabolites⁶ may further reduce colonic mucosal P_{O_2} . Under such conditions, it seems likely that NOS independent pathways contribute to NO generation.

In vivo rectal dialysis has shown excessive luminal accumulation of nitrite in active UC.⁷

Using homogenates of colonoscopic biopsy specimens from 10 patients with UC, we have found that, when incubated anaerobically with nitrite and NADH, inflamed mucosa produces NO (median NO generation 0.01, range 0.0047–0.033 $\mu\text{M}/\text{mg}$ protein/min), detected directly with a chemiluminescence NO analyser.

Although it is possible that, in vivo, luminal bacterial nitrate/nitrite reductases catalyse this reaction, another candidate is the structurally similar mucosal enzyme, xanthine oxidase.⁸ This ubiquitous metalloflavoprotein is abundant in normal liver and small bowel, and catalyses the oxidation of NADH under hypoxic conditions to form NO with concomitant reduction of nitrite (fig 1). The activity of xanthine oxidase is increased in ischaemia-reperfusion injury and by activated neutrophils and a variety of inflammatory mediators present in inflamed gut

Figure 1 Production of nitric oxide (NO) from NO synthase (NOS) is oxygen dependent, but from xanthine oxidase (XO) occurs in hypoxic conditions. Reagents and buffers were pre-equilibrated with nitrogen. NOS (100 μM) was incubated with 1 mM L-arginine in buffer containing 50 mM HEPES (pH 7.4), 1 mM NADPH, 1 mM EDTA, 1.25 mM CaCl_2 , 1 mM dithiothreitol, and 10 $\mu\text{g}/\text{ml}$ calmodulin (1). XO (70 μM) was incubated with 1 mM NADH and 1 mM nitrite (2). As controls, XO was incubated with nitrite (3), NADH with nitrite (4), and XO with NADH (5). Reactions were carried out in a 60 ml sealed flask with a reaction mixture volume of 500 μl . After incubation at 37°C for 20 hours, head-space gas was aspirated and injected directly into an NO analyser (TEA-502, Thermo Electron Corp, USA). Aerobic experiments were performed in a 25 ml flask linked to a NO-free air source, and hypoxic experiments were performed in an anaerobic cabinet (MDH, Hampshire) with aqueous O_2 concentration < 1%. The mean of three experiments is shown.

mucosa. Using non-dissociated polyacrylamide gel electrophoresis with NADH activity staining, we have confirmed the presence of xanthine oxidase in homogenates of colonic biopsy samples from patients with UC.

As the mucosal production of hydrogen peroxide and reactive oxygen metabolites is notably increased in the inflamed bowel,⁹ the recently described non-enzymatic pathway involving the reaction of hydrogen peroxide with both L- and D-arginine⁹ may be a third route by which NO is produced in UC.

Conventionally, NO generation in UC has been attributed predominantly to the activity of NO synthase.^{1–3} We suggest that, in the inflamed and hypoxic large bowel wall, important alternative sources include the reduction of nitrite by xanthine oxidase and bacterial nitrate/nitrite reductases, and the non-enzymatic reaction of hydrogen peroxide with arginine.

Z ZHANG
D P NAUGHTON
E CARTY
D S RAMPTON
MRC Collaborative Centre,
1–3 Burtonhole Lane,
Mill Hill, London NW7 1AD, UK and
Inflammation Research Group and
Digestive Diseases Research Centre,
St Bartholomew's and Royal London
School of Medicine and Dentistry,
London E1 2AD, UK

- 1 Warner TD, McCartney SA. NOS inhibitors in colitis: a suitable case for treatment [commentary]? *Gut* 1998;42:152.
- 2 Lundberg JON, Lundberg JM, Alving K, *et al*. Nitric oxide and inflammation: the answer is blowing in the wind. *Nat Med* 1997;3:30–1.
- 3 Boughton-Smith NK, Evans SM, Hawkey CJ, *et al*. Nitric oxide synthase activity in ulcerative colitis and Crohn's disease. *Lancet* 1993;342:338–40.
- 4 Leone AM, Palmer RM, Knowles RG, *et al*. Constitutive and inducible nitric oxide synthases incorporate molecular oxygen into both nitric oxide and citrulline. *J Biol Chem* 1991;266:23790–5.
- 5 Wrong OM, Edmonds CJ, Chadwick VS. *The large intestine. Its role in mammalian nutrition and homeostasis*. Lancaster: MTP Press Ltd, 1981:168–9.
- 6 Simmonds NJ, Rampton DS. Inflammatory bowel disease—a radical view. *Gut* 1993;34:865–8.
- 7 Roediger WEW, Lawson NJ, Nance SH, *et al*. Detectable colonic nitrite levels in inflammatory bowel disease: mucosal or bacterial malfunction? *Digestion* 1986;35:199–204.
- 8 Zhang Z, Naughton DP, Blake DR, *et al*. Human xanthine oxidase converts nitrite ions into nitric oxide. *Biochem Soc Trans* 1997;25:524S.
- 9 Nagase S, Takemura K, Ueda A, *et al*. A novel non enzymatic pathway for the generation of nitric oxide by the reaction of hydrogen peroxide and D- or L-arginine. *Biochem Biophys Res Commun* 1997;233:150–3.

BOOK REVIEWS

Surgical Diseases of the Pancreas.

Howard J, Idezuki Y, Ihse I, Prinz R. (Pp 868; illustrated; £115.00.) Williams and Wilkins, 1998. ISBN 0-683-18019-3.

This book is a delight to own. It has already become my first choice for seeking out detailed information about obscure topics in pancreatic disease. The scope is defined by its title, and in a field where there are two three other similar volumes, this book stands ahead of its rivals because it contains in one volume almost everything the pancreatic surgeon will need to know about his specialty. The pages are packed with information although the small font size sometimes makes heavy going.

The production of the book is first class, and the quality of the illustrations, particularly the radiographs and photomicrographs, is excellent. The authors of the chapters are predominantly (by a ratio of 5 to 4) from North America, and the remainder from Europe and Japan. This gives a truly international feel.

The chapters vary from all embracing and heavily referenced (e.g. Causes of Pancreatitis has over 900 references, Pathophysiology and Pancreatitis deals with all the possible complications and covers medical treatment, with nearly 300 references) to chapters closely focused on specific areas covered in a few pages (Tumour Markers, Epidemiology of Cancer).

There is even detailed description of operative technique, rare in a book of this type, but welcome as an insight into how other experienced surgeons have approached difficult problems.

Some of the editing is understandably light handed in a book of this size. There is some variation in style, and it was surprising to see detailed description of multiple factor scoring systems in two consecutive chapters.