ELECTRONIC LETTER

Partial hexasomy 15pter→15q13 including *SNRPN* and D15S10: first molecular cytogenetically proven case report

A Nietzel, B Albrecht, H Starke, A Heller, G Gillesen-Kaesbach, U Claussen, T Liehr

J Med Genet 2003;40:e28(<http://www.jmedgenet.com/cgi/content/full/40/3/e28>)

Isodicentric chromosomes 15 (idic(15) or inv dup(15) chromosomes) are the most common supernumerary marker chromosome (SMC) in humans.¹ Buckton *et al*² found that 0.24% of newborns had a SMC, of which 50% could be described as idic(15). The presence of such chromosomes is usually associated with mental retardation, developmental delay, epilepsy, and behavioural problems like autism.³ It is possible that the clinical severity depends on the size of the SMC and its euchromatic material, for example, almost no clinical signs were detectable in a case with idic(15)(pter→q12),⁴ while a severe phenotype including all typical clinical signs was present in a patient with the karyotype 47,+idic(15)(pter→q13).^{5,6} However, there are also reports which show an inconsistent relationship between marker size, gene dosage, and severity of the phenotype.^{7,8} In the study of Mignon *et al*,⁸ in all 16 cases studied the derivative chromosomes 15 were of maternal origin with an identical methylation profile, and neither imprinting nor methylation could explain the phenotypic variability. Partial trisomy of 15q11-q14 resulting from intrachromosomal duplication of chromosome 15q11-q14 has also been reported.^{9,10}

Here we describe the first molecular cytogenetically proven case of a girl with the karyotype 48,XX,+2 idic(15)(pter→15q13).ish 15q13(SNRPN×6, D15S10×6). The clinical details and the results of the molecular and molecular cytogenetic studies are presented and discussed.

MATERIALS AND METHODS

Clinical report

The girl was born after 41 weeks of an uneventful pregnancy to a non-consanguineous German couple. At birth, the mother was 27 years and the father 31 years old. Anthropometric measurements at birth were within the normal range (length 50 cm, weight 3520 g, head circumference 34 cm). The girl had postaxial polydactyly of both hands (fig 1A(a)) and the right foot (fig 1A(b)), a haemangioma on the back (5 × 5 cm), and muscular hypotonia. The heart showed patent ductus arteriosus and an open foramen ovale. From the age of 3 days the girl developed seizures that are still difficult to prevent. At the age of 3¼ years she presented with severe developmental delay and microcephaly (44 cm, -4 SD) (fig 1A(c)). Height and weight were within the normal range for her age. In addition, she has strabismus, hyperopia, and coxa vara on both sides. MRI scan of the skull showed pachygyria.

Cytogenetics and molecular cytogenetics

Cytogenetic and molecular cytogenetic studies were performed on chromosomes derived from peripheral blood. Chromosome preparations and GTG banding were performed according to standard techniques¹¹ (fig 1B). Fluorescence in situ hybridisation (FISH) including RNase and pepsin pretreatment, denaturation of the slides, and addition of the probe to the sample were performed according to standard

Key points

- We report on a case of a partial hexasomy 15pter→15q13 including the loci *SNRPN* and D15S10 as shown by different fluorescence in situ hybridisation (FISH) methods.
- The newborn girl presented with postaxial polydactyly of both hands and one foot, muscular hypotonia, and a heart defect. She developed seizures and at the age of 3 years she had severe developmental delay, microcephaly, and pachygyria.
- The karyotype was described as 48,XX,+2mar after GTG banding. The origin of the two additional identical supernumerary marker chromosomes (SMC) was characterised using a centromere specific multicolour FISH (cenM FISH) approach. Multicolour banding (MCB) and two probes specific for the Prader-Willi-syndrome chromosome region in 15q13 defined the derivative chromosomes as idic(15)(pter→q13).
- PCR based methylation analysis at the *SNRP* locus suggested that the isodicentric chromosomes 15 were of maternal origin.

protocols.¹² CenM FISH was performed as described previously¹³ with the modification that as 24th probe a probe specific for the short arms of all acrocentric chromosomes was added to the cenM FISH probe mix (fig 1C(a)); the probe called midi54 is described in Mrasek *et al*.¹⁴ The cenM54 FISH results (fig 1C(b)) have been verified using single colour FISH with two commercially available centromeric probes for chromosome 15 (alpha satellite probe and satellite III DNA probe, Vysis (fig 1D). Additionally, the *SNRPN* and the D15S10 specific probes in 15q13 (Vysis, fig 1D) and the multicolour banding (MCB) probe set for chromosome 15^{15,16} were applied (fig 1E). The results were evaluated on a fluorescence microscope equipped with a CCD camera and an image analysis system (MetaSystems, Altlussheim, Germany).

Methylation analysis

Methylation analysis for the *SNRP* exon 1 region was performed using methylation specific PCR.¹⁷ The PCR products were analysed on a 2% agarose gel (fig 1F).

RESULTS

Cytogenetic analysis in the severely retarded girl (fig 1A(c)) showed a karyotype 48,XY,+2mar (fig 1B). The small markers were present in all lymphocyte metaphase spreads analysed. The markers could not be detected in either parent.

A new variant of the centromere specific multicolour FISH (cenM FISH), cenM54 FISH, applied to metaphases of the

child showed in one single hybridisation that a partial hexasomy of 15pter→15q12-14 was present, as the SMC were characterised as dicentric derivatives of chromosome 15 including two p arms (fig 1C). FISH with a commercially available alphoid probe and satellite III probe for chromosome 15 (Vysis) confirmed the result (fig 1(D)). Additionally, the *SNRPN* specific probe in 15q13 (Vysis⁴), the probe D15S10 (data not shown), and the multicolour banding (MCB) probe set for chromosome 15^{15 16} (fig 1E) showed that the SMC could be described as *idic(15)(pter→15q13).ish 15q13 (SNRPN×6,D15S10×6)*.

The methylation analysis at the *SNRPN* locus allows determination of the ratio of the PCR product of the methylated maternal allele compared to the PCR product of the unmethylated paternal allele. The overexpression of the methylated maternal PCR product suggests that the derivative chromosomes 15 are of maternal origin (fig 1(F)).

DISCUSSION

To the best of our knowledge this is the first report of a case with a karyotype 48,XX,+2 *idic(15)(pter→15q13).ish 15q13 (SNRPN×6,D15S10×6)*. The two isodicentric chromosomes 15 were characterised by GTG banding (fig 1B), by *cenM54* FISH (fig 1C(b)), by multicolour banding (MCB),^{15 16} (fig 1E) and by locus specific probes for 15q13 (fig 1D). Apart from the common cases with one *idic(15)*,^{1 3 5 6} corresponding to a partial trisomy or tetrasomy of the Prader-Willi and Angelman syndrome critical region (PWACR), one mosaic case with two *idic(15)* with one copy of *SNRPN* and *GABRB3* each, leading to a partial tetrasomy of the PWACR,¹⁸ and two non-mosaic cases have been described so far with partial pentasomy of 15q11-q13 (karyotype: 47,XX,dup(15) (q11q13),+*idic(15)(pter→q13)*).^{19 20} Robinson *et al*²¹ described a case with 48,XX,+2,inv dup(15)(q13). However, they could only show the presence of four copies of the PWACR. Cases described in the time before molecular cytogenetic characterisation was possible⁴ might have had six copies of the PWACR as well. However, to prove this such cases would have to be re-examined.

A maternal origin of the two derivative chromosomes 15 of the present case is most likely, as described for all similar clinical cases with tetrasomy 15q13.^{8 18 19 21-23} Mechanisms for *idic(15)* formation have been proposed previously.^{21 24} For the

establishment of a dicentric chromosome, a crossing over event in meiosis I followed by a non-disjunction in meiosis II has been suggested by Martinsson *et al*.²⁴ The presence of two identical isodicentric chromosomes can be explained by a non-disjunction event in early embryogenesis.

In 1993, Robinson *et al*²¹ postulated that the clinical severity of cases with *idic(15)* is associated with the dosage of the PWACR rather than with differences in the extent of the duplicated segment. This suggestion is confirmed by the clinical signs of the present case, which are in accordance with a very severe type of the “inv dup(15) syndrome”.²⁵ However, the polydactyly is unique in our patient and has not been described in patients with SMC(15) so far.

ACKNOWLEDGEMENTS

This work was supported by the Herbert Quandt Stiftung der VARTA AG, the Robert Pflieger-Stiftung, the DFG (PO284/6-1), the Wilhelm Sander-Stiftung (99.105.1), and the EU (ICA2-CT-2000-10012 and QLRT-1999-31590). We thank Bernhard Horsthemke (Essen, Germany) for helpful discussion, Karin Buiting (Essen, Germany) for continuous assistance, and design of fig 1F and C Lich and H Solka for excellent technical assistance. The continuous support of the Carl Zeiss GmbH (Jena, Germany) is gratefully acknowledged.

Authors' affiliations

A Nietzel, H Starke, A Heller, U Claussen, T Liehr, Institut für Humangenetik und Anthropologie, 07740 Jena, Germany
B Albrecht, G Gillesen-Kaesbach, Institut für Humangenetik und Anthropologie, Universitätsklinikum, 45122 Essen, Germany

Correspondence to: Dr T Liehr, Institut für Humangenetik und Anthropologie, Postfach, D-07740 Jena, Germany; i8li@mti.uni-jena.de

REFERENCES

- Webb T. Inv dup(15) supernumerary marker chromosomes. *J Med Genet* 1994;**31**:585-94.
- Buckton KE, Spowart G, Newton MS, Evans HJ. Forty four probands with an additional “marker” chromosome. *Hum Genet* 1985;**69**:353-70.
- Hotopf M, Bolton P. A case of autism associated with partial tetrasomy 15. *J Autism Dev Disord* 1995;**25**:41-9.
- Miny P, Basaran S, Kuwertz E, Holzgreve W, Pawlowski IH. Inv dup (15): prenatal diagnosis and postnatal follow-up. *Prenat Diagn* 1986;**6**:303-6.
- Gorla N, Slavutsky I, Lisanti J, Pedrazzini E, Vanella L, Larrija I. Cytogenetic characterization of an extra structurally abnormal chromosome associated with severe mental retardation: inv dup (15) (q13). *Hereditas* 1998;**129**:1-5.
- Rauch A, Pfeiffer RA, Trautmann U, Liehr T, Rott HD, Ulmer R. A study of ten small supernumerary (marker) chromosomes identified by fluorescence in situ hybridization (FISH). *Clin Genet* 1992;**42**:84-90.
- Webb T, Hardy CA, King M, Watkiss E, Mitchell C, Cole T. A clinical, cytogenetic and molecular study of ten probands with supernumerary inv dup (15) marker chromosomes. *Clin Genet* 1998;**53**:34-43.
- Mignon C, Malzac P, Moncla A, Depetris D, Roessel N, Croquette MF, Mattei MG. Clinical heterogeneity in 16 patients with inv dup 15 chromosome: cytogenetic and molecular studies, search for an imprinting effect. *Eur J Hum Genet* 1996;**4**:88-100.
- Ungaro P, Christian SL, Fantès JA, Mutirangura A, Black S, Reynolds J, Malcolm S, Dobyns WB, Ledbetter DH. Molecular characterisation of four cases of intrachromosomal triplication of chromosome 15q11-q14. *J Med Genet* 2001;**38**:26-34.
- Abeliovich D, Dagan J, Werner M, Lerer I, Shapira Y, Meiner V. Simultaneous formation of inv dup(15) and dup(15q) in a girl with developmental delay: origin of the abnormal chromosomes. *Eur J Hum Genet* 1995;**3**:49-55.
- Verma RS, Babu A. *Human chromosomes - manual of basic techniques*. New York: Pergamon Press, 1989.
- Liehr T, Thoma K, Kammler K, Gehring C, Ekici A, Bathke KD, Grehl H, Rautenstrauss B. Direct preparation of uncultured EDTA-treated or heparinized blood for interphase FISH analysis. *Appl Cytogenet* 1995;**21**:185-8.
- Nietzel A, Rocchi M, Heller A, Starke H, Wlodarska I, Beensen V, Claussen U, Liehr T. A new multicolor-FISH approach for the characterization of marker chromosomes: centromere-specific multicolor-FISH (cenM-FISH). *Hum Genet* 2001;**108**:199-204.
- Mrasek K, Heller A, Rubtsov N, Trifonov V, Starke H, Rocchi M, Claussen U, Liehr T. Reconstruction of the female Gorilla gorilla karyotype by Zoo-FISH using 25-color FISH and multicolor banding (MCB). *Cytogenet Cell Genet* 2001;**93**:242-48.
- Chudoba I, Plesch A, Lorch T, Lemke J, Claussen U, Senger G. High resolution multicolor-banding: a new technique for refined FISH analysis of human chromosomes. *Cytogenet Cell Genet* 1999;**84**:156-60.

Figure 1 (A) The left hand (a) and the right foot (b) of the female patient showing postaxial hexadactyly and (c) the patient's head showing microcephaly. (B) GTG banding results of the two normal and the two derivative chromosomes 15, karyotype: 48,XX,+2mar. (C) The identical two supernumerary marker chromosomes (SMC) could be identified as derived from chromosome 15 *cenM54* FISH (b). As in the five colour overlay the resulting colours in some cases look very similar, below each chromosome pair the corresponding colour combinations are given. They were analysed in the different colour channels using MetaSystems software. The labelling scheme for *cenM* FISH is depicted in (a). The probe *mid54* (m54) is specific for the short arms of all human acrocentric chromosomes; as outlined in Mrasek *et al*⁴ the chromosomal bands 9p12 and 9q13-21.1 are stained by *mid54* too. The SMC are identified as dicentric der(15) including two short arms of the acrocentric chromosomes (arrow-head) applying the *cenM54* probe set. (D) Results of a two colour FISH experiment hybridising simultaneously commercially available centromeric probe for chromosome 15 (Vysis, SpectrumGreen) and the *SNRPN* probe specific for the Prader-Willi and Angelman syndrome region (Vysis). The result of *cenM* FISH was verified by the centromeric probe and the PWS specific probes showed that a hexasomy of this region was present. (E) Multicolour banding (MCB) results of the two chromosomes 15 and the two SMC. Pseudo-colour and fluorescence profiles clearly indicate the origin of the chromosomal material on the *idic(15)* chromosomes. The short arms are pseudo-coloured in grey. (F) Results of the analysis of allelic methylation differences in 15q11-q13. The maternal (mat) and paternal (pat) allele are present in an internal control (C) in equal proportions, while in the patient (P) the maternal allele is clearly over-represented. Thus, maternal origin of one chromosome 15 and the two *idic(15)* is suggested.

- 16 **Liehr T**, Heller A, Starke H, Rubtsov N, Trifonov V, Mrasek K, Weise A, Kuechler A, Claussen U. Microdissection based high resolution multicolor banding (MCB) for all 24 human chromosomes. *Int J Mol Med* 2002;**9**:335-9.
- 17 **Zeschnigk M**, Lich C, Buiting K, Doerfler W, Horsthemke B. A single-tube PCR test for the diagnosis of Angelman and Prader-Willi syndrome based on allelic methylation differences at the SNRPN locus. *Eur J Hum Genet* 1997;**5**:94-8.
- 18 **Crolla JA**, Harvey JF, Sitch FL, Dennis NR. Supernumerary marker 15 chromosomes: a clinical, molecular and FISH approach to diagnosis and prognosis. *Hum Genet* 1995;**95**:161-70.
- 19 **Long FL**, Duckett DP, Billam LJ, Williams DK, Crolla JA. Triplication of 15q11-q13 with inv dup(15) in a female with developmental delay. *J Med Genet* 1998;**35**:425-8.
- 20 **Abeliovich D**, Dagan J, Werner M, Lerer I, Shapira Y, Meiner V. Simultaneous formation of inv dup(15) and dup(15q) in a girl with developmental delay: origin of the abnormal chromosomes. *Eur J Hum Genet* 1995;**3**:49-55.
- 21 **Robinson WP**, Binkert F, Gine R, Vazquez C, Müller W, Rosenkranz W, Schinzel A. Clinical and molecular analysis of five inv dup(15) patients. *Eur J Hum Genet* 1993;**1**:37-50.
- 22 **Cheng SD**, Spinner NB, Zackai EH, Knoll JH. Cytogenetic and molecular characterization of inverted duplicated chromosomes 15 from 11 patients. *Am J Hum Genet* 1994;**55**:753-9.
- 23 **Maraschio P**, Zuffardi O, Bernardi F, Bozzola M, De Paoli C, Fonatsch C, Flatz SD, Ghersini L, Gimelli G, Loi M, Lorini R, Peretti D, Poloni L, Tonetti D, Vanni R, Zamboni G. Preferential maternal derivation in inv dup(15): analysis of eight new cases. *Hum Genet*. 1981;**57**:345-50.
- 24 **Martinsson T**, Johannesson T, Vujic M, Sjostedt A, Steffenburg S, Gillberg C, Wahlstrom J. Maternal origin of inv dup(15) chromosomes in infantile autism. *Eur Child Adolesc Psychiatry* 1996;**5**:185-92.
- 25 **Battaglia A**, Gurrieri F, Bertini E, Bellacosa A, Pomponi MG, Paravatou-Petsotas M, Mazza S, Neri G. The inv dup(15) syndrome: a clinically recognizable syndrome with altered behavior, mental retardation, and epilepsy. *Neurology* 1997;**48**:1081-6.