Family tree.

(X25 at 9q13).² Most patients are homozygous for expansions of a GAA triplet repeat within intron 1 of this gene. Normal alleles have between 7 and 22 repeats, whereas the range in affected patients is 200 to 900 copies. Around 5% of patients carry heterozygous mutations, a GAA expansion on one allele being accompanied by a point mutation on the other. In this report, the clinical features of a family being investigated for presumed autosomal dominant spinocerebellar degeneration, and in which GAA expansions were identified, are reported. The finding raises the possibility that heterozygous carriers of this mutation may manifest clinical symptoms and signs.

The proband (figure; III.1) is currently 28 years of age, and has a 10 year history of insidious onset and progressive unsteadiness when walking. Upper limb ataxia and dysarthria had supervened around 7 years previously. He needs the support of a single person or a rollator Zimmer for walking, and uses a wheelchair for longer distances. A week long course of synthetic gonadotrophin had been received in childhood for delayed puberty. Examination disclosed pes cavus and kyphoscoliosis. He was dysarthric. There was no optic atrophy, but horizontal gaze evoked nystagmus and bidirectional hypermetropic saccades were noted. He had intention tremor and dysmetria, worse on the left. Reflexes were retained and plantars were flexor. Gait was ataxic, with no clinical evidence of lower limb weakness or impairment of joint position sense. The following investigations gave normal or negative results: serum biochemistry, thyroid function, lipids, urinary amino acids, blood film and full blood count, serum vitamin B₁₂, red cell folate, vitamin E, phytanic acid, syphilis serology, autoantibody profile, chest radiography, EEG, MRI of head and cervical spine, CSF cytology, biochemistry, and immunology, antiPurkinje cell antibody assay, and gene tests for spinocerebellar ataxia types 1 and 3. Nerve conduction studies showed an axonal neuropathy affecting sensory fibres. Lower limb somatosensory evoked potentials were unobtainable, and those in the upper limbs were of low amplitude but normal latency. Echocardiography disclosed mildly impaired left ventricular contractility.

The proband's sister (III.4) is currently 24 years of age, and has had poor balance and ataxia since 15 years. Dysarthria has been

present for four years. She is wheelchair bound. She has taken valproate for presumed complex partial seizures since the age of 8. Examination disclosed limb incoordination, worse on the left. Lower limb reflexes were hyperactive, especially on the left, and both plantars were upgoing.

The proband's father (II.4) is currently 47 years of age. He sought help 4 years ago for partial and generalised seizures, and is currently taking lamotrigine. He admitted to heavy alcohol intake. Examination disclosed hepatomegaly and some memory impairment. There was sustained horizontal gaze evoked nystagmus, limb incoordination worse on the left, diminished knee jerks, absent ankle jerks, and flexor plantar responses. He has marked truncal ataxia. Brain MRI was normal.

The proband's daughter (IV.2) was seen at the age of 8 months after becoming floppy, unable to sit, or control her head acutely. She was intermittently irritable and sleepy, and had vomited repeatedly. She was admitted to hospital, where she was afebrile and well perfused. She could not sit unsupported, was unable to reach out for toys, and was hypotonic. She had gaze dependent very fast vertical nystagmus with no failure of upgaze. Ultrasound examination of the head, blood gases, blood and urinary amino acids, liver function tests, and blood ammonia were normal. At review 6 months later, a further episode of acute ataxia in the context of a febrile infection was reported. This had settled within a few hours, and she had not been admitted to hospital. Since then she has been well (currently 3.5 years) with normal development.

Polymerase chain reaction for the GAA triplet repeat in Friedreich's ataxia was carried out using the primers and method described by Campuzano *et al.*² III.1 showed two expanded alleles carrying 320 and 840 GAA repeats. III.4 was homozygous for two expansions of 660 repeats. II.4 and IV.2 were heterozygous for the GAA expansion, the expanded allele bearing a repeat of 840 triplets consistent with their carrier status. II.5 was also heterozygous for the expansion, and III.2 had no expanded alleles.

The presentation of this family with ataxic features in three generations had suggested a form of autosomal dominant cerebellar ataxia. However, the proband and his sister have clinical phenotypes consistent with

Friedreich's ataxia, and this diagnosis has been established by genome analysis. The carrier status of the two other members of this pedigree manifesting ataxic features has been confirmed. This finding raises the possibility that Friedreich's ataxia carriers are at risk of developing ataxia, especially in the context of environmental insults (such as alcohol in II.4 and viral infections in IV.2). In a recent series of 56 pedigrees, at least two heterozygous parents (both fathers) manifesting ataxic features were identified.³ No data are available on possible environmental insults in these members. The current finding lends support to the conclusion of Lamont *et al.*³ that a history of ataxia in preceding (or successive) generations should not preclude a diagnosis of Friedreich's ataxia. Finally, it may be fruitful to investigate those who develop spinocerebellar ataxia secondary to recognised environmental insults for their carrier status of Friedreich's ataxia.

We are grateful to Professor Tetsuo Ashizawa, Dr Pragna Patel, and Dr Sanjay Bidichandani at the Department of Neurology, Baylor College of Medicine, Houston, Texas for assistance with genome analysis and helpful discussion.

RAJITH DE SILVA
RICHARD PETTY

Department of Neurology, Institute of Neurosciences,
Southern General Hospital, 1345 Govan Road,
Glasgow G51 4TF, Scotland, UK

MARY LOUDON

Monklands District General Hospital, Monkscourt
Avenue, Airdrie ML6 0JS, Scotland, UK

CATHERINE FREW

ALEXANDER COOKE

ROSEMARIE DAVIDSON

Duncan Guthrie Institute of Medical Genetics,
Yorkhill, Glasgow G3 8SJ, Scotland, UK

Correspondence to: Dr R de Silva, Old Church
Hospital, Old Church Road, Romford RM7 0BE,
UK.

1 Harding AE. Friedreich's ataxia: a clinical and genetic study of 90 families with an analysis of early diagnostic criteria and intrafamilial clustering of clinical features. *Brain* 1981;104:589-620.

2 Campuzano V, Montermini L, Moltò MD, *et al.* Friedreich's ataxia: autosomal recessive disease caused by an intronic GAA triplet repeat expansion. *Science* 1996;271:1423-7.

3 Lamont PJ, Davis MB, Wood NW. Identification and sizing of the GAA trinucleotide repeat expansion of Friedreich's ataxia in 56 patients. Clinical and genetic correlates. *Brain* 1997;120:673-80.

Intracerebral haemorrhage due to possible venous obstruction in the neck

Multiple concurrent cerebral haemorrhages in the absence of trauma or a bleeding diathesis suggest venous sinus thrombosis, multiple haemorrhagic infarcts, and haemorrhagic metastases. Iatrogenic venous obstruction is another possible cause. Patient 1 was a 55 year old obese woman who underwent a left posterior fossa craniotomy (Jannetta procedure) for trigeminal neuralgia. Examination and a CT of the head were normal. At surgery she was positioned on her right side with her head held in slight lateral flexion to place the left occipital area uppermost. During the operation the anaesthetist reported that on two occasions the pulse rate slowed only to return to normal when the retractor was immediately removed. The patient's head was repositioned to provide increased lateral flexion and because of her short neck and broad shoulders, slight Trendelenburg tilt was applied. A large artery was found indenting

the left fifth nerve. On repositioning the retractor, swelling of the left hemisphere was noted which bulged through the craniectomy. A small cerebellar resection was performed and the head repositioned with upward tilt. The operation was then abandoned.

Postoperatively she failed to regain consciousness. Brain CT showed moderate swelling of the left cerebellar hemisphere and both occipital lobes appeared hypodense. There were also high density haemorrhagic lesions in the left temporal, right frontal and occipital lobes. She died the next day.

At postmortem the brain was swollen and markedly congested with patchy areas of subarachnoid haemorrhage scattered over the cerebral convexities. The venous sinuses and cortical veins were patent. The vessels of the circle of Willis were patent and free of atheroma. Multiple haemorrhagic venous infarcts were present in the right anterior frontal (40×30×20 mm), right posterior frontal (30×20×20 mm), right parietal (15×10×10 mm), right temporal (20×20×10 mm), right occipital (25×10×20 mm) and left posterior frontal (50×35×40 mm) lobes. The overlying subarachnoid space and sulci in the affected areas were widely distended with blood (sulcal haematomas) and adjacent cerebral cortex and underlying subcortical white matter were disrupted by confluent small haemorrhages (haemorrhagic infarction of venous type). Microscopical assessment of the haemorrhagic lesions disclosed venous and capillary congestion and coalescence of multiple microhaemorrhages typical of venous obstruction. The left cerebellum was lacerated and swollen.

In 1978 Jannetta *et al* reviewed 825 cases of posterior fossa surgery.¹ Five were complicated by supratentorial haemorrhage, and in four the cause remained elusive. Two came to postmortem examination disclosing haemorrhage without evidence of underlying neoplasia or vascular malformation but it was not clear whether they were arterial or venous. It was considered that patient position (modified sitting) may have been in some way implicated and subsequent procedures were done in the lateral decubitus position. In 1988 Hanakita and Kondo² reviewed 278 patients who underwent microvascular decompression all in the lateral decubitus position. There were two fatal intracerebral haemorrhages but there was no postmortem. Excessive cerebellar traction and a disturbance of venous return with an increase in blood pressure was assumed to be the cause in one, but no explanation was given for the second.

The haemorrhages which occurred in our case were simultaneous and widespread and pathologically were venous yet the venous sinuses were patent macroscopically.

We postulate then that the venous haemorrhages were due to acute intraoperative venous hypertension secondary to neck positioning, possibly contributed to by the short thick neck. It is not the lateral decubitus position versus the sitting position which so important but the neck positioning.

In support of the hypothesis of mechanical venous obstruction, Gooding and Stimac³ have demonstrated in animal and infant cadaver models that turning the head to one side results in torsion and compression of the ipsilateral jugular vein. They postulate that jugular venous occlusion on the side of the dominant venous drainage can result in severely limited cerebral venous drainage. Emerson and Parker⁴ showed that with obstruction there was a marked reduction in

Figure 1 Brain CT on day 10 showing multiple areas of cerebral haemorrhage.

cerebral blood flow with an increase in cerebral venous pressure to 25 mm Hg. Doppler studies have shown that cranial venous drainage is often asymmetric, being more dominant on the right and with unilateral venous occlusion, efferent contralateral shunting of blood is only possible to the dominant side. It was the right side of the neck which was compressed at surgery in our case.

Case 2 was a 40 year old woman with cryptogenic cirrhosis who underwent orthotopic liver transplantation. On admission for transplantation a central venous line was inserted into the right internal jugular vein. A Swan-Ganz catheter was inserted into the left internal jugular vein intraoperatively.

Postoperatively transient thrombocytopenia was noted, the platelet count falling to 68 000 rising to 100 000 by day 2 and 335 000 by day 9. On day 4 the Swan-Ganz catheter was removed and on day 7 the central venous line was taken out. In addition to cirrhosis of uncertain aetiology the liver showed multifocal well differentiated hepatocellular carcinoma.

On the ninth post-operative day she complained of dizziness, blurred vision, and headache and the next day she was transiently dysphasic. Brain CT disclosed multiple superficial and deep haemorrhages in both hemispheres (fig 1). There were no cutaneous or systemic bleeding manifestations at this or any other time. Brain MRI confirmed the haemorrhagic lesions and disclosed patent venous sinuses. A subsequent ultrasound of the jugular veins 4 days later showed circumferential thrombus in both the right and left internal jugular veins with some venous flow. A repeat CT 1 month later was normal (fig 2). The patient remains well.

Neurological complications after liver transplantation include cerebral haemorrhage but we could find no case report of multiple perioperative intracerebral haemorrhages without a significant bleeding diathesis.

Radiologically there were multiple haemorrhagic lesions without venous sinus thrombosis. Clinically there was no coagulopathy, no vasculitis, or evidence of haemorrhagic metastases. The complete resolution of the haemorrhages without evidence of encephalo-

Figure 2 Brain CT 1 month postdischarge showing complete resolution of the haemorrhagic cerebral lesions.

malacia and the multifocal haemorrhages not conforming to any particular arterial territory supported the hypothesis that they were venous.

We therefore postulate that jugular venous catheterisation induced jugular venous thrombosis and venous hypertension with secondary multiple haemorrhages. There is one case⁵ reported of an infant aged 2 months who died of multiple cerebral haemorrhages thought to be produced by bilateral internal jugular vein thromboses caused by jugular vein catheters.

The concept of venous hypertension causing cerebral haemorrhage is not generally recognised. Neck positioning or traction and jugular vein catheterisation causing extracranial venous obstruction are two possible mechanisms with important management implications.

We thank the Department of Radiology and Department of Media and Illustration, Flinders Medical Centre, for their assistance in the preparation of radiology and pathology illustrations. We also thank Vicky Frangoulis for manuscript preparation.

MICHELLE A KILEY

RICHARD J BURNS

Department of Neurology, Flinders Medical Centre, Bedford Park 5042 South Australia

PETER C BLUMBERGS

Department of Neuropathology, Institute of Medical and Veterinary Science, Adelaide, South Australia

Correspondence to: Dr R J Burns, Department of Medicine, Flinders Medical Centre, Bedford Park 5042, South Australia.

- 1 Haines SJ, Maroon JC, Jannetta PJ. Supratentorial intracerebral haemorrhage following posterior fossa surgery. *J Neurosurg* 1978;49:881-6.
- 2 Hanakita J, Kondo A. Serious complications of microvascular decompression operations for trigeminal neuralgia and hemifacial spasm. *J Neurosurg* 1988;22:348-52.
- 3 Gooding CA, Stimac GK. Jugular vein obstruction caused by turning of the head. *AJNR Am J Neuroradiol* 1983;4:1223-6.
- 4 Emerson TE, Parker JL. Effects of local increase of venous pressure on canine cerebral haemodynamics. In: Langlitt JW, McHenry L, Revrich M, *et al*, eds. *Cerebral circulation and metabolism*. New York: Springer-Verlag, 1975: 10-13.
- 5 Müller KM, Blaesser B. Tödliche thrombotische Komplikationen nach zentralen Venenkatheter. *Dtsch Med Wschr* 1976;101: 411-13.