al with the same monoclonal antibodies and in the same *Journal* (see Flisser² for review). Similarly, the authors lack knowledge on the initial Mexican publications regarding the association of HLA and cysticercosis.²

Many Mexican authors have published long ago the clinical and imaging descriptions of neurocysticercosis as well as its classification.² Furthermore the authors criticise cestocidal treatment in neurocysticercosis and state that "there is no evidence that cysticidal treatment does more good than harm". Publications by Mexican authors²⁻⁶ clearly demonstrate the usefulness of drug treatment, as does the welfare of patients (children and adults) after such treatment seen in routine work by many neurologists in Mexico. The fact that seizures and inflammation are exacerbated during treatment is the main manifestation of a cestocidal effect, as was experimentally described in cysticercotic pigs that received treatment and had an important enhancement of the inflammatory response surrounding the parasites.²⁻⁶ Due to this, anti-inflammatory and anticonvulsive drugs are recommended in an individual based scheme.

The second part of the article—"management of epilepsy in the developing world"—misleads on the problem of cysticercosis, as it deals almost exclusively with epilepsy in children, when it is well known that the association of this syndrome with neurocysticercosis is mainly with late onset epilepsy.²⁻⁶

From these considerations, different conclusions are drawn:

- Epidemiology of neurocysticercosis is quite well known (not only in Mexico but in several countries where it has been studied)
- Imaging and serological diagnosis are useful tools
- Cysticidal treatment is usually beneficial to patients with neurocysticercosis
- Neurocysticercosis is a major cause of acquired epilepsy
- Prevention measures, based on health education, targeted treatment, and swine vaccination, will probably be sufficient to control neurocysticercosis, and even to eradicate the parasite.

A FLISSER

Departamento de Microbiología y Parasitología,
Facultad de Medicina, Universidad Nacional
Autónoma de México, México DF 04510, México

Correspondence to: flisser@servidor.unam.mx

- 1 Pal KD, Carpio A, Sander JWAS. Neurocysticercosis and epilepsy in developing countries. *J Neurol Neurosurg Psychiatry* 2000;68:137-43.
- 2 Flisser A. Taeniasis and cysticercosis due to *Taenia solium*. In: T Sun, ed. *Progress in clinical parasitology*. Florida, USA: CRC Press, 1994: 77-116. (Review article.)
- 3 Sarti E, Schantz PM, Avila G, et al. Mass treatment against human taeniasis for the control of cysticercosis: population-based intervention study. *Trans R Soc Trop Med Hyg* 2000;94:85-9.
- 4 Sarti E, Flisser A, Schantz PM, et al. Development and evaluation of a health education intervention against *Taenia solium* in a rural community in Mexico. *Am J Trop Med Hyg* 1997;56:127-32.
- 5 Plancarte A, Flisser A, Gauci CG, et al. Vaccination against *Taenia solium* cysticercosis in pigs using native and recombinant oncosphere antigens. *Int J Parasitol* 1999;29:643-7.
- 6 Sotelo J, Flisser A. Neurocysticercosis. Practical treatment guidelines. *CNS Drug* 1997;7:17-25. (Review article.)

High psychiatric morbidity in patients with medically unexplained symptoms

Carson *et al*¹ reported that one third of the patients referred by general practitioners to general neurology outpatient clinics had

symptoms that were poorly explained by identifiable organic disease. Similar trends have been reported from a developing country—namely, India. Bagadia *et al*,² for example, have reported that 57% of non-acutely ill patients attending a general hospital medical outpatient department scored high on the general health questionnaire, indicating a likelihood of psychiatric morbidity. The same authors in another study³ have reported that in a general medical outpatient department, 36% of the patients had psychiatric morbidity, of which 24% had pure psychiatric illness and 12% had psychiatric disorders with associated physical illnesses. These figures have an important bearing on the health care delivery system. These patients take up considerable time of a medical specialist for reasons other than those he specialises in and they also account for higher medical expenses.

Other findings of the same study give us important clues for circumventing this problem. For instance, 66% of those patients whose symptoms were not fully explained by "organicity" had diagnosable anxiety and depressive disorders. Referring such patients directly to a psychiatrist would save time of both the consultant neurologists and the patients. One way of doing this would be to "sensitise" the general practitioners about identification of psychiatric problems. An Indian study showed that after a brief period of training in detection of psychiatric morbidity, the sensitivity of rural primary care physicians increased.⁴

We highlight some of the shortcomings of the report of Carson *et al*.¹ In a strict sense, the design does not fulfil the requirements for a "prospective cohort study".⁵ The follow up assessments were based only on the review of "neurological case notes", the details of which are not made explicit in the article.

The degree by which patients' symptoms were considered to be medically explained was rated by five neurologists using a four point Likert scale. This measure is central to the study. Development and standardisation of the scale are not mentioned in the report. For assessing the prevalence rates a reliable and valid scale is needed. The authors have attempted to validate the scale by reviewing the "neurological case notes" after 6 months in consultation with the treating consultant when necessary. Whether the authors reviewing the case notes were blind to the first rating is not clear. The lack of blindness can lead to serious biases. Further, provision of the components of the "initial consultation" (whether it was based only on history and physical examination or it included relevant investigation) would have helped in understanding the use of the Likert scale better.

N MURALI

JAGADISHA

N KAR

S REHMAN

P S V N SHARMA

Department of Psychiatry, Kasturba Medical College,
MANIPAL-576 119, India

Correspondence to: Dr N Murali
navankul@yahoo.com

- 1 Carson AJ, Ringbauer B, Stone J, et al. Do medically unexplained symptoms matter? A prospective cohort study of 300 new referrals to neurology outpatient clinics. *J Neurol Neurosurg Psychiatry* 2000;68:207-10.
- 2 Bagadia VN, Ayyar KS, Lakdawala PD, et al. Value of the general health questionnaire in detecting psychiatric morbidity in a general hospital outpatient population. *Indian Journal of Psychiatry* 1985;27:293-6.

- 3 Bagadia VN, Ayyar KS, Lakdawala PD, et al. Psychiatric morbidity among patients attending medical outpatient department. *Indian Journal of Psychiatry* 1986;28:139-44.
- 4 Isaac MK, Kapur RL, Chandrasekhar CR, et al. Mental health delivery through rural primary care: development and evaluation of a pilot training programme. *Indian Journal of Psychiatry* 1982;24:131-8.
- 5 Regier DA, Burke J. Quantitative and experimental methods in psychiatry. In: Sadock BJ, Sadock VA, eds. *Kaplan and Sadock's comprehensive textbook of psychiatry*, 7th ed. London: Lippincott Williams and Wilkins, 2000:500-45.

The psychoses of epilepsy

The recent editorial by Toone¹ serves to highlight the deficiencies in our understanding of the relation between epilepsy and psychosis. Although some aspects of this association are now better established than they were at the time of Slater's classic report,² insights into the pathophysiological processes underlying these psychoses seem to have advanced very little. A major reason for this, in my opinion, is that the greater focus of research has been on chronic schizophrenia-like psychoses rather than on the episodic psychoses, which are more likely to yield testable hypotheses.

It can be argued that postictal psychoses present the best opportunity to examine some of the pathophysiological mechanisms that underlie the relation. These psychoses occur in patients who have generally been examined in their prepsychotic states, are likely to develop in hospital while the patient is being monitored for epilepsy, and can be closely studied up to their termination and possible recurrence. That they usually occur in relation to partial complex epilepsy that is secondarily generalised suggests the importance of bilateral involvement of limbic structures in their genesis. The fact that postictal psychoses often follow a cluster of seizures must bring into focus the role of neural inhibitory processes in their development. Epileptic conditions are presumed to involve a chronic imbalance of excitatory and inhibitory influences,³ whereas increased inhibition plays an important part in the development and maintenance of the interictal state.⁴ Human temporal lobe epilepsy is associated with enhanced inhibition,⁵ which contrasts it from neocortical epilepsy in which the epileptogenic regions demonstrate reduced afterdischarge thresholds.⁶ The role these inhibitory processes have in the genesis of psychosis has not been adequately examined. Active inhibitory processes may produce focal reversible deficits—such as Todd's paralysis—which may persist for long periods. Is it possible that postictal psychoses are a manifestation of similar processes bilaterally in the limbic cortex? The question lends itself to examination using electrophysiological and neuroimaging techniques, and even in vivo neurotransmitter measurement using microdialysis in patients being monitored with depth electrodes.⁷

The inhibitory processes not only bring about the postictal state but are also necessary for maintaining the interictal state. Postictal and brief interictal psychoses may therefore not be so different from each other as is generally thought. As the development of seizures may be due to either disinhibition or hypersynchrony involving enhanced disinhibition,³ the occurrence of a seizure during psychosis may have different pathogenetic mechanisms and may indicate either disinhibition or increased inhibition. The effect on

psychosis of a seizure could thereby be either an amelioration of symptoms or their exacerbation. This may explain why seizures during postictal psychosis often exacerbate the psychosis whereas those during brief interictal psychosis may improve the psychiatric status. Different patterns of excitation and inhibition may also explain why the EEG may show "forced normalisation" in some cases of interictal psychosis. These speculations can be tested by the longitudinal examination of patients with brief psychoses, using neurophysiological and neuroimaging methods.

The important neurotransmitters involved in the inhibitory processes are GABA, opioids, and adenosine, whereas glutamate is the key excitatory amino acid. Both GABA⁸ and glutamic acid⁹ have been implicated in the development of psychosis. Glutamate is also important for the maintenance of brain plasticity and surges in its concentrations may be responsible for mossy fibre sprouting in the hippocampus in temporal lobe epilepsy. These plastic brain changes associated with epilepsy raise the question whether repeated postictal psychoses eventually lead to chronic psychosis. There is a suggestion that this may indeed be the case in some patients,¹⁰ but few follow up studies have been published.

The above mechanisms may explain some episodic psychosis, but the enigmatic question is: Why is it that only some patients with recurrent partial complex epilepsy develop psychosis? Is it the "right" mix of inhibition and excitation that is responsible? Is it the involvement of some key anatomical structures? Does the development of psychosis require a substratum of a structural abnormality on which the interplay of these processes comes about? Unless we begin to consider these questions empirically, the relation between psychosis and epilepsy will continue to remain mysterious.

P SACHDEV

Neuropsychiatric Institute, Prince of Wales Hospital,
Randwick, NSW 2031, Australia

- 1 Toone BK. The psychoses of epilepsy. *J Neurol Neurosurg Psychiatry* 2000;69:1-3.
- 2 Slater E, Beard AW, Glithero E. The schizophrenia-like psychoses of epilepsy, i-v. *Br J Psychiatry* 1963;109:95-150.
- 3 Engel J Jr. Excitation and inhibition in epilepsy. *Can J Neurol Sci* 1996;23:167-74.
- 4 Swanson TH. The pathophysiology of human mesial temporal lobe epilepsy. *J Clin Neurophysiol* 1995;12:2-21.
- 5 Cherlow DG, Dymond AM, Crandall PH, et al. Evoked response and after-discharge thresholds to electrical stimulation in temporal lobe epileptics. *Arch Neurol* 1977;34:527-531.
- 6 Penfield W, Jasper H. Epilepsy and the functional anatomy of the human brain. Boston: Little Brown, 1954.
- 7 Wilson CL, Maidment NT, Shomer MH, et al. Comparison of seizure related amino acid release in human epileptic hippocampus versus a chronic, kainate rat model of hippocampal epilepsy. *Epilepsy Res* 1996;26:245-54.
- 8 Lewis DA, Pierri JN, Volk DW, et al. Altered GABA neurotransmission and prefrontal cortical dysfunction in schizophrenia. *Biol Psychiatry* 1999;46:616-26.
- 9 Tamminga C. Glutamatergic aspects of schizophrenia. *Br J Psychiatry* 1999;37(suppl):12-15.
- 10 Kanemoto K, Kawasaki J, Kawai I. Postictal psychosis: a comparison with acute interictal and chronic psychoses. *Epilepsia* 1996;37:551-6.

Diagnostic criteria of multiple system atrophy

I think that the correspondence concerning diagnostic criteria for corticobasal degeneration and the specificity of glial cytoplasmic inclusions¹ should not be closed, as the reply

by Xuereb and Hodges contains a somewhat misleading statement concerning the diagnosis of multiple system atrophy (MSA). In fact there was a consensus statement on the diagnosis of MSA,² establishing three diagnostic categories reflecting differing levels of certainty: definite, probable, and possible. Its conclusion is that "the diagnosis of definite MSA can only be made after a neuropathological examination of the CNS disclosing the characteristic density and distribution of glial cytoplasmic inclusions" in association with degenerative changes. Thus the presence of glial cytoplasmic inclusions is not confirmatory evidence, but the hallmark lesion of MSA.

P L LANTOS

Department of Neuropathology, Institute of Psychiatry,
De Crespigny Park, Denmark Hill,
London SE5 8AF, UK

Correspondence to: npath@iop.kcl.ac.uk

- 1 Lantos PL. Diagnostic criteria for corticobasal degeneration. *J Neurol Neurosurg Psychiatry* 2000;69:705-6.
- 2 Gilman S, Low PA, Quinn N, et al. Consensus statement on the diagnosis of multiple system atrophy. *J Neurol Sci* 1999;163:94-8.

BOOK REVIEWS

Normal and pathologic development of the human brain and spinal cord.

By MARIA DAMBSKA and KRYSZYNA E WISNIEWSKI (Pp 212, £45.00). Published by John Libbey, London, 1999. ISBN 0 86196 591 4

Recent years have witnessed a considerable increase in the knowledge about the genetic and molecular mechanisms involved in the programming of normal, and the genetic defects implicated in the abnormal development of the CNS. Therefore the need for good introductory texts, which focus on the morphological development and also provide information about the genetic as well as molecular processes, is paramount. The book discussed here is such a concise, summary text and reflects the experience of two distinguished experts; one (MD) is a developmental neuropathologist and the second (KEW) a paediatric neurologist/neuropathologist. The book comprises two parts each with a separate section for references. The first part covering chapters 1-3, deals with the normal, and the second, comprising chapters 4-9, describes the abnormal development. The first two chapters are on the development of the cellular elements of the CNS and vessels, and the third chapter gives an account of the morphogenesis of CNS structures. This relatively long chapter is especially well supported with photomicrographs, informative drawings, and tabulated information. Chapter 4 gives a short overview of the aetiological factors involved in the abnormal development, and chapter 5 deals with the phakomatoses. Chapter 6 on the CNS malformations is understandably one of the longest. It provides a detailed account of the major abnormalities, such as neural tube defects, midline malformations, cortical malformations and dysplasia, cerebellar abnormalities, microcephaly and megalencephaly, brain stem as well as spinal cord anomalies, hydrocephalus, and also

changes related to vascular, skull bone, and meningeal anomalies. There are short chapters on developmental disturbances due to chromosomal aberrations (chapter 7), late anomalies (chapter 8), and finally on delay of the CNS maturation (chapter 9).

This is a concise and well written book, which makes entertaining reading even for the non-paediatric neuropathologist reviewer. The chapters are well balanced and the illustrations are of good quality. I have no hesitation in recommending this book not only for neuropathologists, especially for trainees, but also for paediatric neurologists requiring a specialist knowledge of both normal and abnormal CNS development.

TAMAS REVESZ

Multiple sclerosis. By GEORGE D PERKIN and JERRY S WOLINSKY. (Pp 68, £12.00.) Published by Health Press, Oxford, 2000. ISBN 1-899541-28-4

This book is written as part of the *Fast Facts* series to serve as guides to clinical practice. It is envisaged by the authors that the book would be of particular value to family physicians but also nurses and paramedical staff involved in the care of patients with multiple sclerosis.

In the first chapter, dealing with epidemiology, pathology, and pathophysiology, an up to date account is given of the immunological aspects of the condition, which should be of particular value to the reader unfamiliar with inflammatory disease of the nervous system. In the following chapter, on classification, presentation, and early stages, it is particularly helpful to have several case histories which illustrate the various symptoms and presentations listed. Of particular value is a section on questions which patients often ask, with sensible and balanced answers in the chapter on the established condition. The criteria applied to the diagnosis of multiple sclerosis are outlined.

The book then considers treatment aspects of the disease and begins with treatment of acute attacks and symptomatic measures. Of particular interest is the section on symptomatic measures including a practical account of management of spasticity and bladder, bowel, and sexual problems. In a chapter on treatment with immunomodulators an overall view of the evidence for β -interferon in relapsing disease and glatiramer acetate copolymer 1 is given and brief reference to immunoglobulin. Clearly such a treatment chapter depends on an evolving data base of results from clinical trials. Paramedical staff and support groups and future trends are the subjects of the last two chapters. The practical guide to the role of the physiotherapist, occupational therapist, speech and swallowing therapist, and psychologist should be of use to the interested physician.

The authors are well respected and experienced clinicians in the field of multiple sclerosis and although the general approach and various views expressed in such a text will necessarily reflect individual or combined opinions, most of the points raised and conclusions drawn would be supported by most neurologists with an interest in multiple sclerosis. The text is not referenced in a formal way although towards the back of the book a list of references is given to selected subject areas and important clinical trials. Overall I think that the book is a useful addition to the multiple sclerosis library and in