successful in bringing about considerable weight reduction and also resolution of symptoms and signs of IIH including reduction of CSF pressure to normal.

How does obesity lead to an increase in intracranial pressure? Hormonal mechanisms have been postulated,⁷⁰ but Sugarman *et al.*⁶² have provided persuasive evidence in morbidly obese people for a simpler mechanism. They showed that their obese patients with IIH had raised intra-abdominal pressures, raised intrathoracic pressures, and raised central venous pressures, supporting a direct cause and effect relation. However, they did not have a satisfactory, concurrent control group, and did not provide an explanation as to why the pulmonary artery pressures were higher in their obese patients with IIH than in equivalently obese patients without IIH. It would be easy to diagnose this patient group as just having a pickwickian syndrome, but only two out of eight morbidly obese patients of Sugarman *et al.*⁶⁰ had evidence of alveolar hypoventilation with reduced blood oxygen and raised blood carbon dioxide concentrations. Obesity is common, whereas IIH is rare, and obesity affects males as well as females, whereas IIH is much commoner in females than males, so obesity cannot be a sole cause of IIH, even though it may be the immediate cause in a suitably predisposed person. The established relation between obesity and IIH raises further questions. Although IIH in obese persons may be rare, headache in people of all shapes and sizes is very common, and the question does arise as to whether a proportion of obese persons with headache but without papilloedema in fact do have IIH. In the classic IIH series, papilloedema was present in 100% cases and was a diagnostic sine qua non.⁷¹⁻⁷⁴ However, it is not necessary to modify the Dandy criteria much further to open up a whole new vista of cases of overweight patients with bad headache and with high CSF pressures. Indeed, only 12 out of 24 patients had papilloedema in the updated series of Sugarman *et al.*⁶³ Several centres have published on IIH without papilloedema.⁷⁵⁻⁷⁷ Disappointingly, the response of the patients to medication aimed at lowering intracranial pressure was poor.

Conclusion

It seems inescapable that the condition currently called IIH is heterogeneous, and indeed Johnston *et al.*⁷⁸ proposed using the term pseudotumour syndrome to encompass this heterogeneity. In some patients there may be just one aetiology operating, such as occult venous sinus thrombosis. Perhaps others have risk factors which combine to precipitate the condition. In particular, in many cases obesity may be a risk factor whereas in extreme cases it may be a sufficient cause. Brain MRI really should be able to provide definitive information about cerebral and CSF volumes in IIH, but as yet the tunnel has not shed much light, and intriguing enigmas remain. Laboratory animal research, if possible, into factors influencing function of the arachnoid villi might well be informative. More effective means of preventing or treating obesity would undoubtedly have an impact on the prevention and treatment of IIH.

R W H WALKER

Department of Neurology, Barts and the London NHS Trust, The Royal London Hospital, Whitechapel, London E1 1BB, UK

- Corbett JJ, Savino PJ, Thompson HS, *et al.* Visual loss in pseudotumor cerebri: follow up of 57 patients from five to 41 years and a profile of 14 patients with permanent severe visual loss. *Arch Neurol* 1982;39:461-74.
- Johnston LN, Krohel GB, Madsen RW, *et al.* The role of weight loss and acetazolamide in the treatment of idiopathic intracranial hypertension (pseudotumor cerebri). *Ophthalmology* 1998;105:2313-7.
- Digre KB. Idiopathic intracranial hypertension. *Current treatment options in neurology* 1999;1:74-81.
- Johnston I, Jacobson E, Besser M. Acquired Chiari malformation and syringomyelia following spinal CSF drainage: a study of incidence and management. *Acta Neurochir (Wien)* 1998;140:417-27.
- Burgett RA, Purvin VA, Kawasaki A. Lumboperitoneal shunting for pseudotumor cerebri [see comments]. *Neurology* 1997;49:734-9.
- Eggenberger ER, Miller NR, Vitale S. Lumboperitoneal shunt for the treatment of pseudotumor cerebri. *Neurology* 1990;46:1524-30.
- Spor TC, McHenry JG. Long-term effectiveness of optic nerve sheath decompression for pseudotumor cerebri. *Arch Ophthalmol* 1993;111:632-5.
- Corbett JJ, Nerad JA, Tse DT, *et al.* Results of optic nerve sheath fenestration for pseudotumor cerebri. *Arch Ophthalmol* 1988;106:1391-7.
- Chumas PD, Armstrong DC, Drake JM, *et al.* Tonsillar herniation: the rule rather than the exception after lumboperitoneal shunting in the pediatric population. *J Neurosurg* 1993;78:568-73.
- Payner TD, Prenger E, Berger TS, *et al.* Acquired Chiari malformations: incidence, diagnosis, and management. *Neurosurgery* 1994;34:429-34.
- Fishman RA. *Cerebrospinal fluid in diseases of the nervous system*. 2nd ed. Philadelphia: WB Saunders, 1992.
- Weisberg LA. Computed tomography in benign intracranial hypertension. *Neurology* 1985;35:1075-8.
- Rothwell PM, Gibson RJ, Sellar RJ. Computed tomographic evidence of cerebral swelling in benign intracranial hypertension. *J Neurol Neurosurg Psychiatry* 1994;57:1407-9.
- Reid AC, Matheson MS, Teasdale G. Volume of the ventricles in benign intracranial hypertension. *Lancet* 1980;ii:7-8.
- Jacobson DM, Karanjia PN, Olson KA, *et al.* Computed tomography ventricular size has no predictive value in diagnosing pseudotumor cerebri. *Neurology* 1990;40:1454-5.
- Moser FG, Hilal SK, Abrams G, *et al.* MR imaging of pseudotumor cerebri. *AJR Am J Roentgenol* 1988;150:903-9.
- Gideon P, Sorensen PS, Thomsen C, *et al.* Increased brain water self-diffusion in patients with idiopathic intracranial hypertension. *Am J Neuroradiol* 1995;16:381-7.
- Brooks DJ, Beaney RP, Leenders KL, *et al.* Regional cerebral oxygen utilization, blood flow, and blood volume in benign intracranial hypertension studied by positron emission tomography. *Neurology* 1985;35:1030-4.
- Saah AL, Joynt RJ. Brain swelling of unknown cause. *Neurology* 1956;6:791-803.
- Wall M, Dollar JD, Sadun AA, *et al.* Idiopathic intracranial hypertension. Lack of histologic evidence for cerebral edema. *Arch Neurol* 1995;52:141-5.
- Gjerris F, Borgesen SE. Current concepts of measurement of cerebrospinal fluid absorption and biomechanics of hydrocephalus. *Adv Tech Stand Neurosurg* 1992;19:145-77.
- Donaldson JO. CSF hypersecretion in pseudotumor cerebri. *Trans Am Neurol Assoc* 1979;104:196-8.
- Gideon P, Sorensen PS, Thomsen C, *et al.* Assessment of CSF dynamics and venous flow in the superior sagittal sinus by MRI in idiopathic intracranial hypertension: a preliminary study. *Neuroradiology* 1994;36:350-4.
- Eisenberg HM, McComb JG, Lorenzo AV. Cerebrospinal fluid overproduction and hydrocephalus associated with choroid plexus papilloma. *J Neurosurg* 1974;40:381-5.
- Milhorat TH, Hammock MK, Davis DA, *et al.* Choroid plexus papilloma I. Proof of cerebrospinal fluid overproduction. *Child's Brain* 1976;2:273-89.
- Kollar CD, Johnston IH. Pseudotumor after arteriovenous malformation embolisation [letter]. *J Neurol Neurosurg Psychiatry* 1999;67:249.
- Fishman RA. The pathophysiology of pseudotumor cerebri: an unsolved puzzle. *Arch Neurol* 1984;41:257-8.
- Rekate HL, Brodkey JA, Chizeck HJ, *et al.* Ventricular volume regulation: a mathematical model and computer simulation. *Pediatr Neurosci* 1988;14:77-84.
- Rekate HL, Erwood S, Brodkey JA, *et al.* Etiology of ventriculomegaly in choroid plexus papilloma. *Pediatr Neurosci* 1986;12:196-201.
- Borgesen SE, Gjerris F. Relationships between intracranial pressure, ventricular size, and resistance to CSF outflow. *J Neurosurg* 1987;67:535-9.
- Malm J, Kristensen B, Markgren P, *et al.* CSF hydrodynamics in idiopathic intracranial hypertension: a long-term study. *Neurology* 1992;42:851-8.
- Martins AN. Resistance to drainage of cerebrospinal fluid: clinical measurement and significance. *J Neurol Neurosurg Psychiatry* 1973;36:313-8.
- Calabrese VP, Selhorst JB, Harbison JW. CSF infusion test in pseudotumor cerebri. *Trans Am Neurol Assoc* 1978;103:146-50.
- Janny P, Chazal J, Colnet G, *et al.* Benign intracranial hypertension and disorders of CSF absorption. *Surg Neurol* 1981;15:168-74.
- Sklar FH, Beyer CW, Ramanathan M, *et al.* Cerebrospinal fluid dynamics in patients with pseudotumor cerebri. *Neurosurgery* 1979;5:208-16.
- Gjerris F, Sorensen PS, Vorstrup S, *et al.* Intracranial pressure, conductance to cerebrospinal fluid outflow, and cerebral blood flow in patients with benign intracranial hypertension. *Ann Neurol* 1985;17:158-62.
- Massicotte EM, Del Bigio MR. Human arachnoid villi response to subarachnoid hemorrhage: possible relationship to chronic hydrocephalus. *J Neurosurg* 1999;91:80-4.
- Johnson RN, Maffeo CJ, Dacey RG, *et al.* Mechanism for intracranial hypertension during experimental subarachnoid hemorrhage: acute malfunction of arachnoid villi by components of plasma. *Trans Am Neurol Assoc* 1979;103:138-42.
- Scheld WM, Dacey RG, Winn HR, *et al.* Cerebrospinal fluid outflow resistance in rabbits with experimental meningitis. Alterations with penicillin and methylprednisolone. *J Clin Invest* 1980;66:243-53.
- Cremer PD, Johnston IH, Halmagyi GM. Pseudotumor cerebri syndrome due to cryptococcal meningitis. *J Neurol Neurosurg Psychiatry* 1997;62:96-8.
- Feldmann E, Bromfield E, Navia B, *et al.* Hydrocephalic dementia and spinal cord tumor. *Arch Neurol* 1986;43:714-8.
- Ropper AH, Marmarou A. Mechanism of pseudotumor in Guillain-Barré syndrome. *Arch Neurol* 1984;41:259-61.
- Prockop LD, Fishman RA. Experimental pneumococcal meningitis. Permeability changes influencing the concentration of sugars and macromolecules in cerebrospinal fluid. *Arch Neurol* 1968;19:449-63.
- Riddsdale L, Moseley I. Thoracolumbar intraspinal tumours presenting features of raised intracranial pressure. *J Neurol Neurosurg Psychiatry* 1978;41:737-45.
- Gutierrez Y, Friede RL, Kaliney WJ. Agnathesis of arachnoid granulations and its relationship to communicating hydrocephalus. *J Neurosurg* 1975;43:553-8.
- Gilles FH, Davidson RI. Communicating hydrocephalus associated with deficient dysplastic parasagittal arachnoid granulations. *J Neurosurg* 1971;35:421-6.
- Hayes KC, McCombs HL, Faherty TP. The fine structure of vitamin A deficiency II. Arachnoid granulations and CSF pressure. *Brain* 1971;94:213-24.

- 48 Johnston PK, Corbett JJ, Maxner CE. Cerebrospinal fluid protein and opening pressure in idiopathic intracranial hypertension. *Neurology* 1991;41:1040–2.
- 49 Chandra V, Bellur SN, Anderson RL. Low CSF protein concentration in idiopathic pseudotumor cerebri. *Ann Neurol* 1986;19:80–2.
- 50 Cognard C, Casasco A, Toevi M, et al. Dural arteriovenous fistulas as a cause of intracranial hypertension due to impairment of cranial venous outflow. *J Neurol Neurosurg Psychiatry* 1998;65:308–16.
- 51 Biousse V, Ameri A, Boussier MG. Isolated intracranial hypertension as the only sign of cerebral venous thrombosis. *Neurology* 1999;53:1537–42.
- 52 Plant G, Donald JJ, Jackowski A, et al. Partial, non-thrombotic, superior sagittal sinus occlusion due to occipital skull tumours. *J Neurol Neurosurg Psychiatry* 1991;54:520–3.
- 53 Leker RR, Steiner I. Features of dural sinus thrombosis simulating pseudotumor cerebri. *Eur J Neurology* 1999;6:601–4.
- 54 King JO, Mitchell PJ, Thomson KR, et al. Cerebral venography and manometry in idiopathic intracranial hypertension. *Neurology* 1995;45:2224–8.
- 55 Cremer PD, Thompson EO, Johnston IH, et al. Pseudotumor cerebri and cerebral venous hypertension. *Neurology* 1996;47:1602–3.
- 56 Sussman J, Leach M, Greaves M, et al. Potentially prothrombotic abnormalities of coagulation in benign intracranial hypertension. *J Neurol Neurosurg Psychiatry* 1997;62:229–33.
- 57 Leker RR, Steiner I. Anticardiolipin antibodies are frequently present in patients with idiopathic intracranial hypertension. *Arch Neurol* 1998;55:817–20.
- 58 Kesler A, Ellis MH, Reshef T, et al. Idiopathic intracranial hypertension and anticardiolipin antibodies. *J Neurol Neurosurg Psychiatry* 2000;68:379–80.
- 59 Karahalios DG, Rekeate HL, Khayata MH, et al. Elevated intracranial venous pressure as a universal mechanism in pseudotumor cerebri of varying etiologies. *Neurology* 1996;46:198–202.
- 60 Sugerman HJ, Felton WL 3rd, Salvant JB, et al. Effects of surgically induced weight loss on idiopathic intracranial hypertension in morbid obesity. *Neurology* 1995;45:1655–9.
- 61 King JO, Mitchell PJ, Thomson KR, et al. Cerebral venography and manometry in idiopathic intracranial hypertension [abstract]. *Neuroophthalmology* 1996;16:293.
- 62 Sugerman HJ, DeMaria EJ, Felton WL 3rd, et al. Increased intra-abdominal pressure and cardiac filling pressures in obesity-associated pseudotumor cerebri. *Neurology* 1997;49:507–11.
- 63 Sugerman HJ, Felton WL 3rd, Sismanis A, et al. Gastric surgery for pseudotumor cerebri associated with severe obesity. *Ann Surg* 1999;229:634–40.
- 64 Friedfeld L, Fishberg AM. The relation of the cerebrospinal and venous pressures in heart failure. *J Clin Invest* 1934;13:495–501.
- 65 Ireland B, Corbett JJ, Wallace RB. The search for causes of idiopathic intracranial hypertension. *Arch Neurol* 1990;47:315–20.
- 66 Giuseffi V, Wall M, Siegel PZ, et al. Symptoms and disease associations in idiopathic intracranial hypertension: a case control study. *Neurology* 1991;41:239–44.
- 67 Radhakrishnan K, Ahlskog JE, Cross SA, et al. Idiopathic intracranial hypertension. Descriptive epidemiology in Rochester, Minn, 1976–90. *Arch Neurol* 1993;50:78–80.
- 68 Corbett JJ, Mehta MP. Cerebrospinal fluid pressure in normal obese subjects and patients with pseudotumor cerebri. *Neurology* 1983;33:1386–8.
- 69 Kupersmith MJ, Gamell L, Turbin R, et al. Effects of weight loss on the course of idiopathic intracranial hypertension in women. *Neurology* 1998;50:1094–8.
- 70 Rowe FJ, Sarkies NJ. Visual outcome in a prospective study of idiopathic intracranial hypertension [letter]. *Arch Ophthalmol* 1999;117:1571.
- 71 Weisberg LA. Benign intracranial hypertension. *Medicine* 1975;54:197–207.
- 72 Johnston I, Paterson A. Benign intracranial hypertension. I. Diagnosis and prognosis. *Brain* 1974;97:289–300.
- 73 Rush JA. Pseudotumor cerebri: clinical profile and visual outcome in 63 patients. *Mayo Clinic Proc* 1980;55:541–6.
- 74 Wall M, George D. Idiopathic intracranial hypertension. A prospective study of 50 patients. *Brain* 1991;114:155–80.
- 75 Marcellis J, Silberstein SD. Idiopathic intracranial hypertension without papilledema. *Arch Neurol* 1991;48:392–9.
- 76 Mathew NT, Ravishankar K, Sanin LC. Coexistence of migraine and idiopathic intracranial hypertension without papilledema. *Neurology* 1996;46:1226–30.
- 77 Wang S, Silberstein SD, Patterson S, et al. Idiopathic intracranial hypertension without papilledema. A case control study in a headache center. *Neurology* 1998;51:245–9.
- 78 Johnston I, Hawke S, Halmagyi GM, et al. The pseudotumor syndrome. Disorders of cerebrospinal fluid circulation causing intracranial hypertension without ventriculomegaly. *Arch Neurol* 1991;48:740–7.
- 79 Sussman J, Sarkies NJ, Pickard JD. Benign intracranial hypertension. *Adv Tech Stand Neurosurg* 1998;24:261–305.

EDITORIAL COMMENTARIES

The volumes of memory

In 1937, Papez described a circuit for the processing of emotions, which has subsequently proved to be critical for memory function.¹ Various pathological entities can affect structures in this circuit, resulting in amnesic syndromes. In this issue of the *Journal* (pp 13–28), two related papers by Colchester *et al*² and Kopelman *et al*³ used volumetric MR to assess the differing patterns of atrophy in patients with amnesia caused by several neurological diseases, and to examine the relation of these MR volumes to cognitive performance.

A great deal of recent work has focused on detecting atrophy in patients with neurodegenerative dementias, in particular Alzheimer's disease. Recent papers have reported significant atrophy in structures within the medial temporal lobe in memory impaired subjects even before the clinical diagnosis of Alzheimer's disease.^{4–6} Relatively few studies have examined volumes of medial temporal lobe and other memory subserving structures in non-degenerative amnesic syndromes.

The report of Colchester *et al*² suggests that atrophy among the components of the circuit of Papez can be reliably quantified using MR volumetric assessment, and that amnesic syndromes of varying aetiology show specific patterns of atrophy. Of particular interest was the consistent finding of thalamic atrophy in the patients with Korsakoff's syndrome.

Complementing this work, Kopelman *et al*³ examined the correlations between MR volumes of multiple brain regions and performance on several cognitive tests in patients with amnesic and other cognitive syndromes. Of

note, the strongest relations were seen between hippocampal volume and anterograde memory measures, particularly evident in a factor analysis of the neuropsychological tests, although the thalamic measures did correlate with several memory tests.

These studies provide additional evidence for the critical role of the hippocampus in memory, and support the contribution of additional structures within the circuit of Papez to memory function. This work suggests that volumetric MR may be useful in elucidating the pattern of injury to neuroanatomical networks in various cognitive impairment syndromes.

R SPERLING

Brigham and Women's Hospital, Division of Cognitive and Behavioural Neurology, 221 Longwood Avenue, Boston MA 02115, USA
reisa@rics.bwh.harvard.edu

- 1 Papez JW. A proposed mechanism of emotion. *Arch Neurol Psychiatry* 1937;38:725–43.
- 2 Colchester A, Kingsley D, Lasserson D, et al. Structural MRI volumetric analysis in patients with organic amnesia. 1: methods and comparative findings across diagnostic groups. *J Neurol Neurosurg Psychiatry* 2001;70:13–22.
- 3 Kopelman MD, Lasserson D, Kingsley D, et al. Structural MRI volumetric analysis in patients with organic amnesia. 2: correlations with anterograde memory and executive tests in 40 patients. *J Neurol Neurosurg Psychiatry* 2001;70:23–28.
- 4 Fox NC, Warrington EK, Rossor MN. Serial magnetic resonance imaging of cerebral atrophy in preclinical Alzheimer's disease. *Lancet* 1999;353:2125.
- 5 Jack CR, Petersen RC, Xu YC, et al. Prediction of AD with MRI-based hippocampal volume in mild cognitive impairment. *Neurology* 1999;52:1397–403.
- 6 Killiany RJ, Gomez-Isla T, Moss M, et al. Use of structural magnetic resonance imaging to predict who will get Alzheimer's disease. *Ann Neurol* 2000;47:430–9.

EDITORIAL COMMENTARIES

Long term effects of locomotor training in spinal humans

A major concern for patients after spinal cord injury is whether or not they will ever walk again. For completely paralysed patients this prospect is slim, but the outlook for those with partial cord lesions is more hopeful. Any programme, such as that suggested in the paper by Wirz *et al* (this issue, pp 93–96), which may increase the likelihood of subsequent ambulation is to be welcomed.¹

Control of locomotion in primates is predominantly supraspinal. In the thalamic macaque monkey electrical stimulation of the posterior subthalamic region or the mid-brain tegmentum just ventral to the inferior colliculi produced stepping movements.² This was abolished in the suspended animal if the cord was completely transected, even when reflexes had returned. However, if pathways in the ventral white matter on one or both sides of the cord were spared, stepping and walking did occur.

Locomotor training of patients with unilateral lower limb paralysis supported over a moving treadmill was found to improve power in the paralysed limb to make walking possible with minimal splintage and this was attributed to enhancing spinal interneuronal locomotor networks.³ However, as the ventral part of the cord was intact, at least in part, this improvement could equally well arise from the training of higher centres, if not also to continuing recovery in spinal neurons, although unlikely in those patients who had started training many years after injury.

When this locomotor training was applied to patients with complete spinal cord injury, only four out of 10 showed an EMG response.⁴ The failure of the remainder to respond was attributed to the drugs they had been taking—the adrenoceptor antagonist prazosin or cannabis. Support for this was obtained by giving intrathecal adrenaline (epinephrine) which increased and clonidine which decreased motor performance. It could also be possible that those who responded may have had some tracts functioning that could not otherwise be detected by clinical or electrophysiological techniques, but nevertheless allowed conduction to supraspinal centres.

The present study shows that in the partially paralysed patients there was an improvement in the amplitude of

gastrocnemius EMG on completion of the treadmill training which was still maintained a mean of 2 years later, especially as all these patients continued to walk during this time. The completely paralysed patients could not walk so the smaller improvement in gastrocnemius EMG was lost when they no longer received treadmill training. The improvement during training never reached functional levels in these patients and advocating such intensive treatment without showing a functional benefit could be psychologically harmful. There may be some benefit if these patients were to proceed to a functional electrical stimulation walking programme after discharge from the treadmill training.

As has been advocated, it is important to compare novel interventions in treatment with conventional techniques.⁵ Many treadmill studies lack adequate spinal cord injury control groups to demonstrate the value of such training and although spinal interneuronal responses may be stimulated, the vital contribution of higher centres cannot be ignored. With some current research focused on spinal neuronal regeneration, such as by the transplant of oligodendroglia, olfactory, or Schwann cells, any technique to increase the functional capacity of the receiving circuits must surely be encouraged.

C G INMAN

Department of Spinal Injuries, Rookwood Hospital, Llandaff, Cardiff, and
Department of Academic Surgery, University of Wales College of
Medicine, Cardiff, Wales
clive.inman@UHW-TR.nhs.uk

- 1 Wirz M, Columbo G, Dietz V. Long term effects of locomotor training in humans. *J Neurol Neurosurg Psychiatry* 2001;71:93–6.
- 2 Eidelberg E, Walden JG, Nguyen LH. Locomotor control in macaque monkeys. *Brain* 1981;104:647–63.
- 3 Wernig A, Müller S. Laufband locomotion with body weight support improved walking in persons with severe spinal cord injuries. *Paraplegia* 1992;30:229–38.
- 4 Dietz V, Colombo G, Jensen L, *et al*. Locomotor capacity of spinal cord in paraplegic patients. *Ann Neurol* 1995;37:574–82.
- 5 Field-Fote EC. Spinal cord control of movement: implications for locomotor rehabilitation following spinal cord injury. *Phys Ther* 2000;80:477–84.

Cognitive function in the oldest old: women perform better than men

In the paper by van Exel *et al* in this issue (pp 29–32),¹ the authors examine the influence of sex and formal education on cognitive functioning in a community based sample of subjects over the age of 85. Based on the cognitive reserve theory of dementia, the authors hypothesise that women

would be expected to score more poorly than men on cognitive tests due to a lower level of formal education.

Previous studies provide support for the theory that a lower educational level is a risk factor for the development of dementia.² This relation seems to be more pronounced

in female subjects than males,³ although data regarding the influence of sex on cognitive functioning in non-demented persons were not available.

The results of the current study showed better cognitive performance in the female group, despite their lower level of formal education. One possible explanation raised by the authors was that medical risk factors (for example, atherosclerosis) may be greater in the male group. An alternative, or contributing factor, may be use of formal education as a measure of "cognitive reserve". Although years of education have traditionally been used to estimate premorbid functioning, some authors have suggested that formal education may be less important than later life experiences, such as primary occupation.⁴ A follow up to the current study might examine the role of non-educational experiences on cognitive functioning between men and women.

This paper makes an important and timely contribution to the field of aging research, with the recent emphasis on early diagnosis of dementia. Understanding variables related to

cognitive functioning in elderly people, such as the influence of sex, is essential for improving the ability to detect preclinical markers of dementia and identify "at risk" people who could benefit from clinical prevention trials.

N JOHNSON

Department of Psychiatry, Cognitive Neurology and Alzheimer's Disease Center, Northwestern University Medical School, 320 EAST Superior Street, Suite 11-499, Chicago, IL 60611-3010, USA

johnson-n@northwestern.edu

- 1 van Exel E, Gussekloo J, de Craen A J M, *et al.* Cognitive function in the oldest old: women perform better than men. *J Neurol Neurosurg Psychiatry* 2001;70:29-32.
- 2 Katzman R. Education and the prevalence of dementia and Alzheimer's disease. *Neurology* 1993;43:13-20.
- 3 Ott A, van Rossum CTM, van Harskamp F, *et al.* Education and the incidence of dementia in a large population-based study: The Rotterdam Study. *Neurology* 1999;52:663-6.
- 4 Stern Y, Alexander GE, Prohovnik I, *et al.* Relationship between lifetime occupation and parietal flow: implications for a reserve against Alzheimer's disease pathology. *Neurology* 1995;45:55-60.

1st Asia Pacific Forum on Quality Improvement in Health Care

Three day conference

Wednesday 19 to Friday 21 September 2001

Sydney, Australia

We are delighted to announce this forthcoming conference in Sydney. Authors are invited to submit papers (call for papers closes on Friday 6 April), and delegate enquiries are welcome.

The themes of the Forum are:

- Improving patient safety
- Leadership for improvement
- Consumers driving change
- Building capacity for change: measurement, education and human resources
- The context: incentives and barriers for change
- Improving health systems
- The evidence and scientific basis for quality improvement.

Presented to you by the BMJ Publishing Group (London, UK) and Institute for Healthcare Improvement (Boston, USA), with the support of the the Commonwealth Department of Health and Aged Care (Australia), Safety and Quality Council (Australia), NSW Health (Australia) and Ministry of Health (New Zealand).

For more information contact: quality@bma.org.uk or fax +44 (0) 7383 6869