Subclinical thyrotoxicosis

A C Al-Abadi

Subclinical thyrotoxicosis may be defined as a low serum thyrotrophin (TSH) concentration in an asymptomatic patient with normal serum free thyroxine (T_4) and triiodothyronine (T_3) concentrations. The secretion of TSH may be suppressed even in the presence of normal serum thyroid hormone levels. This reflects the highly sensitive response that the pituitary gland mounts to minor changes in serum free T_4 and T_3 concentrations inside the normal range of the population—exemplified by the log-linear relation between serum TSH and thyroid hormone concentrations.¹

In this review I shall address the various disorders characterised by a low serum TSH and how to differentiate them from subclinical thyrotoxicosis. I also discuss the causes of subclinical thyrotoxicosis, its potential pathophysiological and clinical consequences, and its management.

The introduction in the mid-1980s of sensitive assays for TSH allowed the measurement of serum TSH concentrations well below the normal range of 0.5 to 4.0 mIU/l. Unlike the first generation radioimmunoassays, which have a sensitivity of about 0.1 mIU/l, the new immunometric second and third generation assays have greater sensitivities of approximately 0.05 and 0.005 mIU/l, respectively, with about a 10-fold increase in sensitivity per generation.^{1,2} The thyrotrophin releasing hormone (TRH) stimulation test has since played a diminishing diagnostic role.³

Causes of low serum TSH concentrations

The main causes of low serum TSH concentrations are overt thyrotoxicosis, non-thyroidal illness, secondary (central) hypothyroidism, physiological causes, and subclinical thyrotoxicosis.

OVERT THYROTOXICOSIS

Overt thyrotoxicosis is characterised by high serum free T_4 and T_3 and very low serum TSH concentrations, which are undetectable in more than 95% of patients, even with third generation assays,⁴ and it is associated in the majority of patients with overt symptoms and signs of thyrotoxicosis. Two small subgroups of thyrotoxic patients are characterised by a normal serum T_4 in association with either a raised total T_3 and a normal thyroxine binding globulin (T_3 toxicosis)⁵ or, more rarely, a normal total T_3 but a raised free T_3 (*free T_3 toxicosis*).⁶ The diagnosis in these overtly thyrotoxic patients can therefore be readily made by the demonstration of a rise in either serum free T_4 or total or free T_3 , or both.

NON-THYROIDAL ILLNESSES

Some patients with non-thyroidal illnesses may have low serum TSH concentrations. This is invariably associated with low serum concen-

trations of T_3 and often T_4 . Although the pathogenesis of the reduction in serum TSH in these patients remains uncertain, decreased secretion of TRH, increased secretion of somatostatin, cortisol, or cytokines, and inhibition of TSH secretion by drugs such as dopamine or glucocorticoids may contribute. In addition to a low serum TSH concentration, *secondary (central) hypothyroidism* is often associated with other pituitary hormone deficiencies, and may be accompanied by neurological abnormalities related to the local effect of a hypothalamic-pituitary tumour.

PHYSIOLOGICAL CAUSES

Physiological causes of a reduction in serum TSH include pregnancy and old age. The thyroid stimulating activity of chorionic gonadotrophin, particularly in women with hyperemesis gravidarum, may increase the thyroid secretory activity enough to lower serum TSH concentration to below normal. This is a common finding near the end of the first trimester or at the beginning of the second trimester, and is significantly more common in Asian women than in those of European origin (15.7% *v* 4.8%).⁷ The pulsatile secretion of TSH may result in an interpulse nadir value that is below normal in some elderly subjects.

SUBCLINICAL THYROTOXICOSIS

The causes of subclinical thyrotoxicosis, in which TSH secretion is suppressed but the concentrations of circulating thyroid hormones remain normal—albeit in the upper normal range—can be divided into *exogenous* and *endogenous*. Drug related subclinical thyrotoxicosis may occur, as in the administration of supraphysiological doses of thyroid hormone, and in drug induced thyroiditis (amiodarone, α interferon, iodine in patients with multinodular goitre). Overtreatment with thyroid hormones may be intentional, as in patients with thyroid carcinoma in order to reduce TSH secretion to below normal, or unintentional, as in approximately one quarter of hypothyroid patients receiving treatment.⁸ L-thyroxine overtreatment is by far the commonest cause of subclinical thyrotoxicosis.⁹

When subclinical thyrotoxicosis results from an autonomous oversecretion of thyroid hormones by a diseased thyroid gland, it is termed *subclinical hyperthyroidism*. Approximately one fifth of patients with Graves' disease who have received medical (radioactive iodine)¹⁰ or surgical (subtotal thyroidectomy)¹¹ treatment and are euthyroid may be insufficiently treated and have low serum TSH concentrations. A similar proportion of patients with Graves' disease who remain euthyroid after discontinuation of antithyroid drug treatment may have low serum TSH concentrations which indicate a state of subclinical hyperthyroidism.¹¹ It is

Department of
Diabetes and
Endocrinology,
Clinical Sciences
Centre, Northern
General Hospital,
Herries Road,
Sheffield S5 7AU, UK

Correspondence to:
Dr Al-Abadi
Amira@
ammori.freemove.co.uk

Submitted 8 November 1999
Accepted 27 April 2000