Informing, communicating and sharing decisions with people who have cancer

A J Sowden, C Forbes, V Entwistle, I Watt

There is widespread agreement that people should be informed about health care options and involved in decisions about their own care. In England and Wales the National Institute for Clinical Excellence (NICE) is intent on achieving a patient focus in its work and *The NHS plan* emphasises the need for patients to have more say in their own care and more influence over the way in which the NHS works.¹ Similarly, *The NHS cancer plan* for England and Wales emphasises the need for good communication between health professionals and patients, both for delivering high quality care and for empowering people to be involved in decisions about their own care.²

This paper summarises *Effective Health Care* bulletin 6(6) which focused on communication, information giving, and sharing of decisions between health professionals and people with cancer. Evidence from systematic reviews produced by the Cochrane Consumers and Communication Group, other good quality systematic reviews, and from guidance produced by the National Cancer Guidance Steering Group is also included.

Communicating with patients

The most common complaints made by patients with cancer are about poor communication and inadequate information.^{3,4} In 1993 the General Medical Council (GMC) recognised the need to teach communication skills as part of the British undergraduate medical education. However, in a survey of all consultant non-surgical oncologists working in the UK to which 83% of the 476 consultants responded, almost half considered that they had not received sufficient training in communication skills.⁵

Communication skills programmes vary in their content and training methods. A recent systematic review evaluated communication training programmes for nurses.⁶ Of the 14 studies included, eight were based in an oncology setting (two randomised controlled trials, one controlled trial, and five pre/post tests). Only one study measured the effects on patients; this randomised controlled trial found reduced levels of anxiety but no improvement in health professionals' empathy skills (the only professional outcome measured). The other randomised controlled trial and the controlled trial found improvements in health professionals' attitudes. The five pre/post test studies

reported mixed findings. However, it is unclear whether all relevant studies were identified and the quality of the included studies was not assessed, so the findings of this review should be treated with some caution.

An overview of five studies evaluating training programmes specifically for health professionals dealing with people with cancer found positive changes in communication skills in some participants.⁷ The programmes were based around the acquisition of specific skills and/or strategies for dealing with specific situations. Those who reported more negative attitudes at the beginning of the training programme appeared to benefit most. A systematic review assessing the effects of interventions to promote patient centred care within clinical consultations will be published later this year.⁸

Informing patients

A recent survey of over 2000 people currently receiving treatment for cancer in one of 34 hospital outpatient departments in the UK found that 87% preferred to be given as much information about their illness as possible (both good and bad).⁹ Information is required for different purposes—understanding the presenting symptoms and/or disease, learning about available services, and participating in decisions about treatment options.¹⁰ Patients cannot express informed preferences about their care, or about whether and how they want to participate in care decisions unless they are given sufficient and appropriate information. Findings from recent focus groups carried out with cancer patients suggest that people still experience a dearth of information although a great deal is theoretically available.¹¹

Information materials have been found to be lacking in several areas including omission of relevant data, failure to give a balanced view of the effectiveness of different treatments, ignoring uncertainties, and rarely promoting a participative approach to decision making.¹⁰ There are instruments available to help judge the quality of written information about treatment options¹² and guides to assist in the production of good quality information materials.^{13–15} The criteria for producing “good” quality patient information are shown in table 1.

Current sources of evidence based cancer information include: the Cancer Library which

NHS Centre for Reviews & Dissemination, University of York, York YO10 5DD, UK
A J Sowden, *associate director*
C Forbes, *research fellow*

Department of Health Studies, University of York
I Watt, *professor of primary care*

Health Services Research Unit, University of Aberdeen, Aberdeen AB25 2ZD, UK
V Entwistle, *programme director*

Correspondence to:
Dr A J Sowden
ajs18@york.ac.uk

Table 1 Criteria for producing good quality information for patients

<i>Duman & Farrell</i> ¹³	<i>CHiQ</i> ¹⁶	<i>Entwistle & O'Donnell</i> ¹⁵
Informs about condition and available treatments and options Comprehensive, unbiased information about outcomes Outlines uncertainties and gaps in knowledge Caters for people from a diverse range of backgrounds Regularly reviewed and updated Integrated into a planned shared decision making programme	Accessibility: appropriate format Accuracy Appropriateness Availability Continuity: information is presented with other resources Currency: being up to date	Clear statement of aims Relevant Accurate Accessible, comprehensible, acceptable Gives further sources of information Facilitates judgement about reliability of information
Simple and easily understood language and design	Legibility: clear presentation Originality (not duplicating other sources) Patient involvement Readability Reliability	

is available via the Cochrane Library; a cancer database being developed by Macmillan Cancer Relief and The Centre for Health Information Quality (CHiQ) which will be a source of reference to information materials about cancer and its management¹⁶; and the National Electronic Library for Health which also includes information for patients and the public.

TAILORED INFORMATION

The importance of tailoring information to suit patients' educational background, cultural orientation, and general level of comprehension has been emphasised.¹⁷ A recent Scottish randomised controlled trial has shown that patients with cancer prefer information based on their own medical records to general information.¹⁸

The Teamwork Project aims to help people with cancer work in partnership with health professionals with the aid of a personal information file.¹¹ The file contains general information about cancer, tests, diagnosis, treatment and care, together with a personal health diary where a record is kept of health and health care including personal test results and details of medication prescribed. The file is currently being evaluated by several NHS trusts.

Other personalised information includes recordings or summaries of consultations between people with cancer and their doctors. A recent systematic review of eight randomised controlled trials has evaluated the effects of giving recordings or summaries of consultations to people with cancer.¹⁹ The studies varied in terms of interventions, participants, and outcomes. Interventions ranged from giving audiotapes to people at the end of the consultation to making a written summary of the key points discussed during the consultation. Across the seven studies which provided data, 83–96% of participants receiving tapes or letters reported that they had found them useful. Four of the seven studies also reported better recall in the groups which received personalised recordings or summaries than in the control groups. Of the six studies that assessed anxiety and/or depression, none reported statistically significant differences between the intervention and control groups, although one study found mixed results among participants in the intervention group—psychological morbidity increased at follow up in those with a poor prognosis compared with those with a

better prognosis. Although the quality of the trials was poor, overall the evidence suggests that providing a record of the consultation can increase the amount of information recalled and satisfaction with the information given. There is some evidence to suggest that recordings might encourage participation in subsequent consultations and there is no clear evidence to show that they affect psychological health (either positively or negatively).

THE NEEDS OF HEALTH PROFESSIONALS

Health professionals are likely to need support if patients' information needs are to be met. The learning needs of clinicians in fulfilling the information needs of patients have been identified and include: placing a higher priority on patient information; understanding the patient's needs; understanding the emotional aspects of learning; helping patients to understand; learning from the patient; and knowing about information sources.²⁰ Ensuring that such needs are addressed in both undergraduate and continuing professional education will help in achieving the goals of current government policy.

ORGANISATIONAL CHALLENGES

As well as challenges to the health professional in meeting peoples' information needs, organisational challenges have also been recognised and recommendations made.²⁰ These include: treating patients' information needs as a core activity; ensuring adequate funding, space and time devoted to patient information; producing, implementing and reviewing guidelines on patient information and auditing methods of working.²⁰

Involving patients in decision making

The shared decision making model is increasingly being advocated as a way of promoting clinical effectiveness and more appropriate and efficient use of resources.²¹ A partnership between professionals and patients in which both contribute to decisions about treatment or care is encouraged. This differs from the informed choice model where the emphasis is on the patient to make the decision and from the more traditional model where the health professional makes the decision.^{22–25}

People are likely to vary in the extent to which they want to participate in decision making as well as in which decisions. Health professionals will need the necessary skills to elicit preferences for involvement along with other communication skills.²⁶

One way to support patient involvement in decision making about treatment is through the use of decision aids. Decision aids aim to help people make specific and deliberative choices by providing information on the options and outcomes relevant to their state of health.²⁷ The effectiveness of decision aids has been assessed in three reviews.^{28–30} The search strategy used in the most recent Cochrane review has been re-run and the same inclusion criteria adopted, with the addition that participants had to have a cancer diagnosis. Four randomised controlled trials which examined the use of decision aids in cancer patients were identified, three studies of women with breast cancer^{31–33} and one of men with prostate cancer.³⁴

Little evidence was found that video disk presentations were better than usual care,³³ that interactive multimedia were better than written information,³¹ or that a tape and workbook were better than a pamphlet³² in patients with breast cancer facing major treatment decisions. A very small (non-significant) difference was found toward a lower decisional conflict (agreement/disagreement about being certain, informed, clear about values and supported in decision making) in the tape and workbook group.³² The costs of developing and evaluating the tape and workbook were estimated to be more than \$250 000 (Canadian).³²

In contrast, there was some evidence that decision aids were helpful for men with prostate cancer in reducing anxiety and influencing decision making about treatment.³⁴ Men exposed to an “empowerment” intervention (an information and question pack based on their needs and wants) assumed a significantly more active role in decision making and had lower anxiety levels than men who were given just an information pack.³⁴ Support was also found for decision aids in a systematic review that included patients with a variety of different diagnoses.³⁰ Improved knowledge, increased participation, and decreased decisional conflict were reported.³⁰

Although questions around the use of decision aids remain, including the type of aid to use, the choice of appropriate outcomes to measure, and their effect on communication between patients and professionals,²⁴ they can provide a useful way of displaying and providing information about treatment options. However, barriers to implementing decision aids in practice have been found—for example, a recent US study found time pressures to be the main barrier.³⁵ It is likely that similar barriers would exist in the UK.

Conclusions and recommendations

If the goals of current NHS policy are to be achieved, then the information needs of patients must be addressed. Initiatives such as the Cancer Information Strategy³⁶ should take into account variations in peoples’ preferences for information and involvement in decisions about their treatment and care. This means that health professionals need to know how best to elicit patients’ need and readiness for information as well as their desire for involvement in decision making. Appropriate training

in communication skills addressing such issues should be considered and any programmes offered should be appropriately evaluated, with assessments of patient outcomes as well as professional skills.

Tailored information is an option for people with cancer; in particular, recordings or summaries of key consultations may benefit adults with cancer without causing additional anxiety. The effects of such interventions will, however, need to be carefully monitored because evidence for their effectiveness is currently limited. People should also be given the opportunity for involvement in decisions about their treatment and care, but individual preferences for different levels of involvement need to be respected. One option is the use of appropriate decision aids, although again evidence about the likely effects in people with cancer is limited and further evaluation is recommended.

The Health Services Research Unit receives core funding from the Chief Scientist’s Office, Scottish Executive.

- 1 Department of Health. *The NHS plan*. London: Department of Health, 2000.
- 2 Department of Health. *The NHS cancer plan. A plan for investment. A plan for reform*. London: Department of Health, 2000.
- 3 National Cancer Alliance. *Patient centred cancer services? What patients say*. The National Cancer Alliance, 1996.
- 4 Cancer Guidance Sub-group of the Clinical Outcomes Group. *Improving outcomes in breast cancer*. London: NHS Executive, 1996.
- 5 Ramirez AJ, Graham J, Richards MA, et al. Burnout and psychiatric disorder among cancer clinicians. *Br J Cancer* 1995;71:1263–9.
- 6 Kruijver IPM, Kerkstra A, Francke AL, et al. Evaluation of communication training programs in nursing care: a review of the literature. *Patient Educ Couns* 2000;39:129–45.
- 7 National Cancer Guidance Group. *Improving outcomes in lung cancer: the research evidence*. London: NHS Executive, Department of Health, 1998.
- 8 Skea Z, Lewin S, Dick J, et al. The effects of interventions for health care providers that promote patient centred approaches in clinical consultations: results of a systematic review. In: *8th International Cochrane Colloquium*. Cape Town, South Africa, 2000.
- 9 Jenkins V, Fallowfield L, Saul J. Information needs of patients with cancer: results from a large study in UK cancer centres. *Br J Cancer* 2001;84:48–51.
- 10 Coulter A, Entwistle V, Gilbert D. Sharing decisions with patients: is the information good enough? *BMJ* 1999;318:318–22.
- 11 Smith C. The role of health professionals in informing cancer patients: findings from the Teamwork Project (phase 1). *Health Expect* 2000;3:217–9.
- 12 Charnock D, Shepperd S, Gann B, et al. DISCREN: an instrument for judging the quality of consumer health information on treatment choices. *J Epidemiol Community Health*. 1999;53:105–11.
- 13 Duman M, Farrell C. The POPPi guide: practicalities of producing patient information. London: King’s Fund, 2000.
- 14 Centre for Health Information Quality (CHIQ). *Guidelines for producing and maintaining patient information on conditions and treatments*. CHIQ, 2000.
- 15 Entwistle V, O’Donnell M. *A guide to producing health information*. Health Services Research Unit, 1999.
- 16 Centre for Health Information Quality (CHIQ). *Cancer information*. CHIQ, 2000.
- 17 Harris KA. The informational needs of patients with cancer and their families. *Cancer Practice* 1998;6:39–46.
- 18 Jones R, Pearson J, McGregor S, et al. Randomised trial of personalised computer based information for cancer patients. *BMJ* 1999;319:1241–7.
- 19 Scott JT, Entwistle VA, Sowden AJ, et al. Recordings or summaries of consultations for people with cancer (Cochrane Review). In: *The Cochrane Library*. Oxford: Update Software, 2000.
- 20 Jones R, Tweddle S, Hampshire M, et al. *Patient-led learning for the clinical professions. Fulfilling the needs of patients*. Bristol: NHS Information Authority. National Education Training and Development Programme, 2000.
- 21 Coulter A. Shared decision-making: a summary and future issues. In: Maslin AM, Powles TJ, eds. *Breast cancer. Sharing the decision*. Oxford: Oxford University Press, 1999.
- 22 Charles C, Gafni A, Whelan T. Shared decision-making in the medical encounter: what does it mean? (Or it takes at least two to tango). *Soc Sci Med* 1997;44:681–92.
- 23 Charles C, Whelan T, Gafni A. What do we mean by partnership in making decisions about treatment. *BMJ* 1999;319:780–2.

- 24 Entwistle VA, Sowden AJ, Watt IS. Evaluating interventions to promote patient involvement in decision-making: by what criteria should effectiveness be judged? *J Health Serv Res Policy* 1998;3:100-7.
- 25 Hope T. *Evidence based patient choice*. London: King's Fund, 1996.
- 26 Elwyn G, Edwards A, Gwyn R, et al. Towards a feasible model for shared decision making: focus group study with general practice registrars. *BMJ* 1999;319:753-6.
- 27 O'Connor AM, Fiset V, Rostom A, et al. Decision aids for people facing health treatment or screening decisions (protocol for a Cochrane Review). In: *The Cochrane Library*. Oxford: Update Software, 2000.
- 28 O'Connor A, Fiset V, DeGrasse C, et al. Decision aids for patients considering options affecting cancer outcomes: evidence of efficacy and policy implications. *J Natl Cancer Inst Monogr* 1999;25:67-80.
- 29 Molenaar S, Sprangers MA, Postma-Schuit FC, et al. Feasibility and effects of decision aids. *Med Decis Making* 2000;20:112-27.
- 30 O'Connor AM, Rostom A, Fiset V, et al. Decision aids for patients facing health treatment or screening decisions: systematic review. *BMJ* 1999;319:731-4.
- 31 Street RLJ, Voigt B, Geyer CJ, et al. Increasing patient involvement in choosing treatment for early breast cancer. *Cancer* 1995;76:2275-30.
- 32 Goel V, Sawka C, Thiel EC, et al. Randomized trial of a patient decision aid for choice of surgical treatment for breast cancer. *Med Decis Making* 2001;21:1.
- 33 Maslin AM, Baum M, Walker JS, et al. Using an interactive video disk in breast cancer patient support. *Nurs Times* 1998;94:52-5.
- 34 Davison BJ, Degner LF. Empowerment of men newly diagnosed with prostate cancer. *Cancer Nurs* 1997;20:187-96.
- 35 Holmes-Rovner M, Valade D, Orlowski C, et al. Implementing shared decision-making in routine practice: barriers and opportunities. *Health Expect* 2000;3:182-91.
- 36 NHS Executive. *Cancer information strategy*. London: NHS Executive, 2000.

7th European Forum on Quality Improvement in Health Care

21-23 March 2002

Edinburgh, Scotland

We are delighted to announce this forthcoming conference in Edinburgh. Authors are invited to submit papers (call for papers closes on **Friday 5 October 2001**) and delegate enquiries are welcome.

The themes of the Forum are:

- Leadership, culture change, and change management
- Achieving radical improvement by redesigning care
- Health policy for lasting improvement in health care systems
- Patient safety
- Measurement for improvement, learning, and accountability
- Partnership with patients
- Professional quality: the foundation for improvement
- Continuous improvement in education and training
- People and improvement.

Presented to you by the BMJ Publishing Group (London, UK) and Institute for Healthcare Improvement (Boston, USA). For more information contact: quality@bma.org.uk or look at the website www.quality.bmjpg.com. Tel: +44 (0)20 7383 6409; fax: +44 (0)20 7373 6869.