- 79 Greenstein D, Jeffcote N, Ilsley D, Kester RC. The menstrual cycle and Raynaud's phenomenon. *Angiology* 1996;47:427-36.
- 80 Brouwer RML, Wenting GJ, Schalekamp MADH. Acute effects and mechanism of action of ketanserin in patients with primary Raynaud's phenomenon. *J Cardiovasc Pharmacol* 1990;15:868-76.
- 81 Engelhart M, Neilsen HV, Kristensen JK. The blood supply to fingers during Raynaud's attack: a comparison of laser-Doppler flowmetry with other techniques. *Clin Physiol* 1985;5:447-53.
- 82 Pistorius MA, Planchon D, de Faucal P. Plethysmographic cold test for diagnosis and evaluation of the severity of Raynaud's phenomenon. Validation of the method by factorial analysis of correspondences in 541 patients. *Int Angiol* 1994;13:10-14.
- 83 Zamora MR, O'Brien RF, Rutherford RB, Weil JV. Serum endothelin-1 concentrations and cold provocation in primary Raynaud's phenomenon. *Lancet* 1990;336:1144-7.
- 84 Cooke ED, Bowcock SA, Smith AT. Photoplethysmography of the distal pulp in the assessment of the vasospastic hand. *Angiology* 1985;36:33-40.
- 85 Suichies HE, Aarnoudse JG, Wouda AA, Jentink HW, de Mul FFM, Greve J. Digital blood flow in cooled and contralateral finger in patients with Raynaud's phenomenon. Comparative measurements between photoelectrical plethysmography and laser Doppler flowmetry. *Angiology* 1992;43:134-41.
- 86 Kallenberg GGM, Wouda AA, Meems L, Wesseling H. Once daily felodipine in patients with primary Raynaud's phenomenon. *Eur J Clin Pharmacol* 1991;40:313-15.
- 87 Lafferty K, de Trafford JC, Potter C, Roberts VC, Cotton LT. Reflex vascular responses in the finger to contralateral thermal stimuli during the normal menstrual cycle: a hormonal basis to Raynaud's phenomenon. *Clin Sci* 1985;68:639-45.
- 88 Cooke JP, Creager MA, Osmundson PJ, Shepherd JT. Sex differences in control of cutaneous blood flow. *Circulation* 1990;82:1607-15.

Unusual and memorable

Series editor: Gary D Wright

A 46 year old Indian woman presented with a 15 year history of painless, nodular swellings over the fingers and wrist of her right hand, which were making daily activities increasingly difficult. An excision biopsy three years before had revealed a calcifying aponeurotic fibroma, but the lesion had recurred. She was otherwise well with no other rheumatological or metabolic disease. Plain radiographs showed well defined, calcified lesions that appeared to erode the adjacent bone (fig 1).

Calcifying aponeurotic fibromata are rare soft tissue tumours, usually present in childhood and typically affect the fingers and palms.¹

They are slow growing and painless, and almost never undergo malignant transformation. Histologically, they are characterised by lines of plump fibroblasts, surrounding areas of dense collagenous stroma with focal calcification. Treatment should be conservative, as local recurrence after excision is seen in 50% of cases.

¹ Enzinger FM, Weiss SW. *Soft tissue tumours*. 3rd ed. Missouri: Mosby, 1995.

Contributors: IAN MCCURDIE, SLI S JAWAD. *Rheumatology Department, The Royal Hospitals NHS Trust, Mile End Hospital, Bancroft Road, London E1 4DG.*