might reduce the economic costs to society of back pain, these effects would be limited if no account is taken of the influence of broader non-work related aspects. Moreover, for the individual, such interventions are unlikely to impact on the overall experience of back pain and consultation behaviour.

The authors thank the partners, staff, and patients of Brooklands and Bowland Road General Practices, Manchester, UK, for their support and cooperation, Lesley Jordan for help with administration, and Professor Malcolm IV Jayson for help with the design and development of the study. The study was funded by the National Back Pain Association, Teddington, and the Arthritis Research Campaign, Chesterfield.

- 1 Papageorgiou AC, Croft PR, Ferry S, Jayson MIV, Silman AJ. Estimating the prevalence of low back pain in the general population. *Spine* 1995;20:1889-94.
- 2 Leino P, Magni G. Depressive and distress symptoms as predictors of low back pain, neck-shoulder pain, and other musculoskeletal morbidity: a 10-year follow-up of metal industry employees. *Pain* 1993;53:89-94.
- 3 Bigos SJ, Battie MC, Spengler DM, Fisher LD, Fordyce WE, Hansson TH, et al. A prospective study of work perceptions and psychosocial factors affecting the report of back injury. *Spine* 1991;16:1-6.
- 4 Clinical Standards Advisory Group. *Epidemiology Review: the epidemiology and cost of back pain*. London: HMSO, 1994.
- 5 Papageorgiou AC, Macfarlane GJ, Thomas E, Croft PR, Jayson MIV, Silman AJ. Psychosocial factors in the workplace - do they predict new episodes of low back pain? *Spine* 1997;22:1137-42.
- 6 Macfarlane GJ, Thomas E, Papageorgiou AC, Croft PR, Jayson MIV, Silman AJ. Employment and physical work activities as predictors of future low back pain. *Spine* 1997;22:1143-9.
- 7 Corney RH. Development and use of a short self-rating instrument to screen for psychosocial disorder. *J R Coll Gen Pract* 1988;38:263-6.
- 8 Goldberg D, Williams P. *A user's guide to the General Health Questionnaire*. Windsor: NFER-Nelson, 1988.
- 9 Papageorgiou AC, Croft PR, Thomas E, Ferry S, Jayson MIV, Silman AJ. Influence of previous pain experience on the episode incidence of low back pain: results from the South Manchester Back Pain study. *Pain* 1996;66:181-5.
- 10 Croft PR, Papageorgiou AC, Ferry S, Thomas E, Jayson MIV, Silman AJ. Psychological distress and low back pain: evidence from a prospective study in the general population. *Spine* 1995;20:2731-7.
- 11 Croft PR, Rigby AS. Socioeconomic influences on back problems in the community in Britain. *J Epidemiol Community Health* 1994;48:166-70.
- 12 Walsh K, Cruddas M, Coggan D. Low back pain in eight area of Britain. *J Epidemiol Community Health* 1992;46:227-30.
- 13 Croft PR, Macfarlane GJ, Papageorgiou AC, Thomas E, Silman AJ. The outcome of low back pain in general practice. *BMJ* 1998;316:1356-9.
- 14 Klein-Hesselink DJ, Spruit IP. The contribution of unemployment to socioeconomic health differences. *Int J Epidemiol* 1992;21:329-37.

Unusual and memorable

Series editor: Gary D Wright

A 29 year old women presented with painful stiff fingers on her left hand. The pain was associated with intermittent swelling and limitation of motion of her digits. She was otherwise well with no history of psoriasis or other joint problems. On examination there was diffuse bony swelling of the second and third fingers of her left hand with a reduced range of movement. The overlying skin was shiny and erythematous (fig 1). Radiographs of the hands revealed endosteal hyperostosis of the second and third metacarpals (fig 2). Scintigraphy showed an increased accumulation of radionuclide in these areas (fig 3).

Melorheostosis is a rare bone disorder.¹ It usually presents with joint pain, intermittent swelling, and muscle contractures below the age of 20 years but may occur as late as the fourth or fifth decades. The disease is usually limited to a single limb, in which one or more bones may be affected. The lower limbs are more commonly affected than the upper limbs. Radiographic appearances are highly characteristic. Osseous excrescences extend along the length of the bone and may be associated with endosteal hyperostosis partially or completely obliterating the medullary cavity. Soft tissue calcification and ossification occurs frequently. The cause of melorheostosis is unknown although investigations have indicated possible associations with neurofibromatosis, tuberous sclerosis, hypophosphataemic rickets, and haemangiomas.²

1 Caudle RJ, Stern PJ. Melorheostosis of the hand. A case report with long term follow up. *J Bone Joint Surg* 1987;69A:1229-31.

Contributors: A PENDLETON, GD WRIGHT, SD ROBERTS, *Royal Victoria Hospital, Belfast.*