smokers.^{5,6} It is interesting to note that NPC is associated with the Epstein-Barr virus,⁷ one of a number of viruses which have been implicated in the pathogenesis of RA.⁸ Therefore, passive smoke might potentially predispose subjects to RA as a result of changes within the nasopharynx resulting in antigenic stimulus by a virus that triggers RA.

We agree with the authors that new information is urgently needed about any factor associated with the risk of RA. In view of emerging data highlighting smoking as an important environmental risk factor for the development of seropositive RA,^{3,9} and also this study by Heliövaara *et al*, we propose that passive smoking should be considered as a potential candidate factor for the development of RA.

D HUTCHINSON
R MOOTS
*Rheumatology Department,
University Hospital Aintree,
Liverpool, UK*

- 1 Heliövaara M, Aho K, Knekt P, Impivaara O, Reunanen A, Aromaa A. Coffee consumption, rheumatoid factor, and the risk of rheumatoid arthritis. *Ann Rheum Dis* 2000;59:631-5.
- 2 Yuan JM, Wang XL, Xiang YB, Gao YT, Ross RK, Yu MC. Non-dietary risk factors for nasopharyngeal carcinoma in Shanghai, China. *Int J Cancer* 2000;85:364-9.
- 3 Hutchinson D, Shepstone L, Moots R, Lear JT, Lynch MP. Heavy cigarette smoking is strongly associated with rheumatoid arthritis (RA), particularly in patients without a family history of RA. *Ann Rheum Dis* 2001;60:223-7.
- 4 Halldorsdottir HD, Jonsson T, Thorsteinsson J, Valdimarsson H. A prospective study on the incidence of rheumatoid arthritis among people with persistent increase of rheumatoid factor. *Ann Rheum Dis* 2000;59:149-51.
- 5 Nuorti JP, Butler JC, Farley MM, Harrison LH, McGeer A, Kolczak MS, *et al*. Cigarette smoking and invasive pneumococcal disease. *N Engl J Med* 2000;342:681-9.
- 6 Kriz P, Bobak M, Kriz B. Parental smoking, socioeconomic factors, and risk of invasive meningococcal disease in children: a population based case-control study. *Arch Dis Child* 2000;83:117-21.
- 7 Klein G. The Epstein-Barr virus and neoplasia. *N Engl J Med* 1975;293:1353-7.
- 8 Ollier W. Rheumatoid arthritis and Epstein-Barr virus: a case of living with the enemy? *Ann Rheum Dis* 2000;59:497-9.
- 9 Wilson K, Goldsmith CH. Does smoking cause rheumatoid arthritis? *J Rheumatol*. 1999;26:1-3

Authors' reply

It is not easy to measure passive smoking. Nevertheless, determinations of serum cotinine and thiocyanate recommended for the detection of indirect exposure to tobacco smoke have been shown to indicate exposures of recent occurrence.^{1,2}

To test the assumption put forward by Drs Hutchinson and Moots we studied coffee consumption for its associations with serum cotinine and thiocyanate concentrations in a sample of men who had participated in the Mini-Finland Health Survey and served as an age matched control group in a nested case-control study of lung cancer.³ A modification of the Nicotine Metabolite RIA kit method (Diagnostic Products Corporation, Los Angeles, USA) was used to determine serum cotinine concentrations.⁴ Serum thiocyanate was determined by the spectrophotometric ferric nitrate method.⁵ Of the total of 158 men in the sample, 39 reported current smoking, and 10 others had serum cotinine ≥ 200 mg/l or thiocyanate ≥ 20 nmol/l, suggesting direct exposure to tobacco. Exclusion of these men left 109 non-smokers for the final analyses.

No significant association was seen between the number of daily cups of coffee and serum cotinine concentration (age adjusted $r=0.08$, $p=0.38$). Contrary to the assumption of Drs Hutchinson and Moots, there was a negative correlation between serum thiocyanate level and coffee consumption (age adjusted $r=-0.21$, $p=0.03$). The findings are in agreement with our impression that in Finland coffee is not consumed to any greater extent in notably smoky environments. Rather, coffee consumption is confined to breakfast at home and coffee breaks both during working hours and leisure time.

On the basis of these preliminary results it seems unlikely that passive smoking would have correlated strongly enough with coffee consumption to explain the close associations between coffee consumption and the occurrence of rheumatoid factor (RF) and the risk of rheumatoid arthritis (RA) in our study. Nevertheless, our results are far from being conclusive. Whatever the links between coffee consumption, RF, and RA, we agree with Drs Hutchinson and Moots that passive smoking should be considered a potential risk factor for RA.

M HELIÖVAARA
K AHO
P KNEKT
A REUNANEN
A AROMAA
*National Public Health Institute,
Helsinki, Finland*
O IMPIVAARA
*Social Insurance Institution,
Research and Development Centre,
Turku, Finland*

Correspondence to: Dr M Heliövaara, National Public Health Institute, Mannerheimintie 166, FIN-00300 Helsinki, Finland

markku.heliovaara@ktl.fi

- 1 Tunstall-Pedoe H, Woodward M, Brown CA. Tea drinking, passive smoking, smoking deception and serum cotinine in the Scottish heart health study. *J Clin Epidemiol* 1991;44:1411-14.
- 2 Husgafvel-Pursiainen K, Sorsa M, Engström K, Einistö P. Passive smoking at work: biochemical and biological measures of exposure to environmental tobacco smoke. *Int Arch Occup Environ Health* 1987;59:337-45.
- 3 Knekt P, Raitasalo R, Heliövaara M, Lehtinen V, Pukkala E, Teppo L, *et al*. Elevated lung cancer risk among persons with depressed mood. *Am J Epidemiol* 1996;144:1096-103.
- 4 Perkins SL, Livesey JF, Escarea EA, Belcher JM, Dudley DK. High-performance liquid-chromatographic method compared with a modified radioimmunoassay of cotinine in plasma. *Clin Chem* 1991;37:1989-93.
- 5 Butts WC, Kuehneman M, Widdowson GM. Automated method for determining serum thiocyanate, to distinguish smokers from non-smokers. *Clin Chem* 1974;20:1344-8.

LETTERS TO THE EDITOR

Polyarteritis nodosa complicated by thrombotic thrombocytopenic purpura

A 56 year old woman was diagnosed with polyarteritis nodosa (PAN) in June 1998

based on the presence of fibrinoid necrosis and infiltration of polymorphonuclear cells into medium and small sized arteries on a skin biopsy specimen. She presented with erythema on her arms and legs, with fever and body weight loss. Tender masses were palpable on her right abdomen. Small erythematous lesions and livedo reticularis were seen on the arms and legs.

Laboratory investigation on admission disclosed anaemia (haemoglobin 73 g/l) and leucocytosis ($22.5 \times 10^9/l$) consisting mainly of neutrophils (85%). Creatinine clearance was 39 ml/min. Serological examination showed raised levels of C reactive protein (88.6 mg/l). Serological tests for syphilis, hepatitis B virus antigen, and antibody for hepatitis C virus were negative. A high titre of myeloperoxidase antineutrophil cytoplasmic antibody (MPO-ANCA) (201 EU) was detected in her sera. An abdominal computed tomography scan showed bilateral perirenal haemorrhages.

The patient was treated with 1000 mg of methylprednisolone for three successive days, followed by 500 mg cyclophosphamide intravenously. Plasma exchange was performed, also. Within three days, treatment had reduced the body temperature to normal and lowered the C reactive protein concentration (<10 mg/l).

The patient rapidly developed thrombocytopenia on the fifth day after admission to hospital, and the lowest platelet count was $15.0 \times 10^9/l$ on the eighth day after admission. She became unconscious, and showed features of haemolytic anaemia and renal failure. A provisional diagnosis of disseminated intravascular coagulation was made because of thrombocytopenia, a low concentration of fibrinogen (1.6 g/l), and a high fibrinogen and fibrin degradation product level (22.6 μ g/ml). We treated her with nafamostat mesilate (200 mg/day) and infusion of fresh-frozen plasma. This treatment produced a marked increase in fibrinogen concentration but failed to improve the platelet count. Further laboratory tests showed fragmented red blood cells in peripheral blood. A diagnosis of thrombotic thrombocytopenic purpura (TTP) was established based on these findings. The patient was treated with plasma exchange with 2700 ml fresh-frozen plasma.

Within one week of treatment the platelet count returned to normal and consciousness level improved dramatically. However, renal failure was irreversible and she continued to undergo dialysis. Unfortunately, the patient died from haemorrhage from a duodenal ulcer. Necropsy findings included small and medium sized polyarteritis in the kidney, uterus, pulmonary hilum, hepatic hilum,

Figure 1 Necropsy finding of ileum end. Note fibrinoid necrosis of medium sized artery. Haematoxylin and eosin stain. Magnification $\times 400$.

adrenal gland, ileum (fig 1), and ascending colon. Several stages of vasculitis existed, which was a typical finding of classic PAN. We failed to detect vasculitis affecting arterioles, venules, or capillaries. Duodenal bleeding was from peptic ulcers, which is not associated with arteritis.

A few reports have described TTP complicating certain forms of rheumatic diseases, including systemic lupus erythematosus and systemic sclerosis.^{1,2} To our knowledge there are no reports published in English of PAN complicated by TTP.

The necropsy finding was classic PAN because vasculitis affected vessels larger than arterioles and there were no pathological findings of glomerulonephritis.³ We failed to detect vasculitis affecting arterioles, venules, or capillaries. MPO-ANCA is usually a marker of microscopic polyarteritis or necrotising glomerulonephritis but does not distinguish it from classic PAN with certainty.⁴

Our case suggests the possibility of secondary TTP due to PAN. We speculate that endothelium damage by PAN may enhance the development of TTP, particularly in the presence of inflammatory processes. Our patient unfortunately died from massive bleeding from a duodenal ulcer. However, conventional treatment for TTP, including plasma infusion and plasma exchange, were effective in this case, too.⁵ Our case emphasises the need to consider TTP when thrombocytopenia occurs with vascular disease, because early and correct treatment of TTP may improve morbidity and mortality in these patients.

S YAMASAKI
M TOMINAGA
A KAWAKAMI
K MIGITA
Y KAWABE
K IZUMINO
K EGUCHI

First Department of Internal Medicine,
Nagasaki University School of Medicine,
Nagasaki, Japan

T TSUCHIYA
I SHIMOKAWA

First Department of Pathology,
Nagasaki University School of Medicine

Correspondence to: Dr K Eguchi, First Department of Internal Medicine, Nagasaki University School of Medicine, Sakamoto 1-7-1, Nagasaki 852-8501, Japan
eguchi@net.nagasaki-u.ac.jp

- 1 Fox DA, Faix JD, Coblyn J, Fraser P, Smith B, Weinblatt ME. Thrombotic thrombocytopenic purpura and systemic lupus erythematosus. *Ann Rheum Dis* 1986;45:319-22.
- 2 Miller A, Ryan PFJ, Dowling JP. Vasculitis and thrombotic thrombocytopenic purpura in a patient with limited scleroderma. *J Rheumatol* 1997;4:598-600.
- 3 Jannette JC, Falk RJ, Andrassy K, Bacon PA, Churg J, Gross WL, *et al*. Nomenclature of systemic vasculitis: proposal of an international consensus conference. *Arthritis Rheum* 1994;37:187-92.
- 4 Guillemin L, Lhote F. Polyarteritis nodosa and microscopic polyangiitis. *Clin Exp Immunol* 1995;101(suppl 1):22-3.
- 5 Rock GA, Shumak KH, Buskard NA, Blanchette VS, Kelton JG, Spasoff RA, *et al*. Comparison of plasma exchange with plasma infusion in the treatment of thrombotic thrombocytopenic purpura. *N Engl J Med* 1991;325:393-7.

Placement of intra-articular injections verified by ultrasonography and injected air as contrast medium

Intra-articular injection of long acting corticosteroid is a corner stone in rheumatological treatment. The injected intra-articular corticoid is more effective when correctly placed.^{1,2} Injection of radiographic contrast material has shown that fewer than half of the injections are correctly placed in the joint space after blind injection.¹

Generally, the clinical application of ultrasonographic examinations can be enhanced by contrast agents.³ The most commonly used technique is creation of microbubble contrast agents. Such agents, applied to the bloodstream, have been used for hepatic, nephrologic, cardiologic, and transcranial examinations.⁴ Obviously, the risk of air embolism depends on the anatomical site of the injected air contrast. Transient ischaemic attacks are described after echocardiography with air contrast⁵ and in animal models haemodynamic effects during venous air infusion can be measured.⁶ Intra-articular injection of air and subsequent lateral and posterior radiographs have shown that this technique can enhance the precision of the procedure.⁷ The disadvantage of this method is that the result can first be seen after the injection, and that a correction can only be made with a new injection. In the joint space the air is separated from the vascular system and when only small amounts of sterile air are used the risk of venous air embolism is negligible. Air is a very effective contrast medium in ultrasonography. Air sonography has been used for the diagnosis of meniscus lesions in knee joints⁸ and for rotator cuff lesions in the shoulder.⁹

We expand the applicability of this method to all joints, not only for diagnosis, but also for the correct placement of the needle before injection of medicine (steroid, osmium acid, viscosupplementation). The sterile air that is contained in the capped vial with lidocaine or steroid is used as contrast medium. The needle is guided into the joint space of the distended capsule by ultrasonography.

When the steroid and lidocaine are mixed in the syringe a small volume of air will be in the needle itself (~0.05 ml). The air in the needle is clearly seen when the injection is started and will secure the correct placement of the needle. With this technique, it is not necessary to use two separate syringes and the inclination of the syringe will not cause the air to move from the needle to the bottom of the syringe.

If the knee is injected, injection directly into the recess of the knee is recommended, which will make the small volume of air momentarily visible.

Figure 1 illustrates the ultrasonography of a metatarsophalangeal joint in a patient before and after injected air. The intra-articular air is clearly seen. We have made over 1000 ultrasonography guided intra-articular injections without any complications. This method is easy, inexpensive, without risk and radiation, and should be used routinely in rheumatology. Chemical synovectomy of the knee, especially, should always be guided by ultrasonography, and

with this technique smaller joints can also be considered for chemical synovectomy.

U FREDBERG
Medical Department (Rheumatology),
Silkeborg Centralhospital,
DK-8600 Silkeborg, Denmark

G VAN OVEREEM HANSEN
Department of Rheumatology,
Kong Chr X's Gighospital Graasten,
DK-6300 Graasten, Denmark

L BOLVIG
Department of Radiology,
University Hospital of Aarhus,
DK-8000 Aarhus C, Denmark

Correspondence to: Dr U Fredberg, Ostermarksvej 29, DK-8381 Tilst, Denmark

Fredberg@post4.tele.dk

- 1 Eustace JA, Brophy DP, Gibney RP, Bresnihan B, FitzGerald O. Comparison of the accuracy of steroid placement with clinical outcome in patients with shoulder symptoms. *Ann Rheum Dis* 1997;56:59-63.
- 2 Zingas C, Failla JM, Van Holsbeeck M. Injection accuracy and clinical relief of de Quervain's tendinitis. *J Hand Surg [Am]* 1998;23:89-96.
- 3 Campani R, Calliada F, Bottinelli O, Bozzini A, Sommaruga MG, Draghi F, *et al*. Contrast enhancing agents in ultrasonography: clinical applications. *Eur J Radiol* 1998;27(suppl 2):161-70.
- 4 Widder DJ, Simeone JF. Microbubbles as a contrast agent for neurosonography and ultrasound. *AJR Am J Roentgenol* 1986;147:347-52.
- 5 Srivastava TN, Undesser EK. Transient ischemic attack after air contrast echocardiography in patients. *Ann Intern Med* 1995;122:396.
- 6 Vik A, Jenssen BM, Brubakk AO. Comparison of haemodynamic effects during venous air infusion in pigs. *Eur J Appl Physiol* 1994;68:127-33.
- 7 Bliddal H. Placement of intra-articular injections verified by mini arthrography. *Ann Rheum Dis* 1999;58:641-3.
- 8 Hawe W, Milz P. The clinical value of air sonography in the diagnosis of meniscus lesions. A prospective study of 50 knee joints. *Sportverletz Sportschaden* 1991;5:119-26.
- 9 Hawe W. Air as contrast medium in sonography of the rotator cuff (aero-sonographic diagnosis). *Rontgenpraxis* 1991;44:75-8.

HLA class II alleles and synovial fluid cytology in RA

Rheumatoid arthritis (RA) is associated with HLA-DR4, which is encoded by the DRB1 gene. This genetic predisposition has been shown to lie within the sequence motif present in the third hypervariable region of the DRB1 gene.¹ This sequence of amino acids has been called the "disease epitope" and can be encoded by DR4 subtypes as well as non-DR4 alleles; DR1 subtypes, DR10 and DR14 subtypes.¹ Hence, it is also termed the shared epitope. In addition to imparting susceptibility to RA, HLA-DR4 has also been shown to be associated with the severity of the rheumatoid disease, including destructive erosive joint disease, rheumatoid factor positivity,² and extra-articular manifestations, such as rheumatoid nodules, vasculitis, and Felty's syndrome.^{3,4} HLA-DQ genes are in linkage disequilibrium with HLA-DR and subjects who are DR4 positive may either be DQB1*0301 or *0302 positive; certain extra-articular features of RA have been shown to be associated with HLA-DQB1*0301.^{4,5}

Synovial fluid cytology in RA is heterogeneous both with respect to total white cell