PostScript

MATTERS ARISING

Methotrexate is the leading DMARD in Finland

We read with interest the report by Aletaha *et al* on international questionnaires of the early treatment of rheumatoid arthritis (RA).¹ We would like to draw attention to the newest statistics in Finland.

In Finland the sickness insurance scheme provides reimbursement (today 75%) for prescription drugs in chronic inflammatory rheumatic diseases, principally rheumatoid arthritis (RA), psoriatic arthritis, ankylosing spondylitis, chronic reactive arthritis, juvenile idiopathic arthritis, and systemic rheumatic diseases. Eligibility entails a medical certificate written by a specialist in rheumatology and approved by the welfare authorities. All patients who are entitled to reimbursement are centrally registered by the social insurance institution. Glucocorticoids and disease modifying antirheumatic drugs (DMARDs) are medicines which can be reimbursed in this category.

As the nationwide sickness insurance scheme covers the entire Finnish population of five million, it provides a good basis for an epidemiological survey of the occurrence of all rheumatic diseases and investigation of the use of various DMARDs by these patients.

In 2001 a total of 76 552 patients had an appropriate certificate entitling them to special reimbursement for DMARDs. Of this total, only 47 967 used the drugs listed in table 1. During 2001, 4325 new certificates were granted. Table 1 shows the numbers of users of glucocorticoids and of 12 DMARDs during the past seven years. A rapid rise can be seen in the numbers using methotrexate. In 2001 the number was 16 470, a third of all users, overtaking the previously most used drug sulfasalazine. The 40 year old drug, hydroxychloroquine, has been third most used drug since 1996. The traditional sulphonylurea group drugs, sodium aurothiomalate, auranofin, and penicillamine, have lost 2860 of their earlier users (29%), whereas the newest drug, leflunomide, overtook cyclosporin in 2001. Half of the patients registered used glucocorticoids.

In Finland today rheumatic patients with specially reimbursed drugs have the code 202 on their national insurance card, but this code is the same for all rheumatic diseases. In her thesis in 1997 Kaipiainen-Seppänen counted that during four study years in a population of one million people, 3076 new patients had chronic inflammatory rheumatic disease and 491 connective tissue disease.² It can thus be calculated that around 16% of the patients in table 1 had connective tissue diseases.

Sulfasalazine was introduced in 1941, but the number of its users has continued to grow, albeit slowly, presumably owing to a sulphur allergy. Methotrexate has been for a decade the main drug used in RA globally, and it is also increasingly used in the spondyloarthropathies. Hydroxychloroquine is the preferred drug for systemic lupus erythematosus (SLE) and active Sjögren's syndrome. However, the preponderance of these three drugs is also based on their frequent use in combination therapies. Finnish doctors are well informed of the Fin-RACo study in early RA: treatment with methotrexate, sulfasalazine, hydroxychloroquine, and prednisolone has proved more successful than single therapy.³ The recent report by Aletaha *et al*, based on two matched questionnaires in 1997 and 2000, shows the same trend as table 1.¹ The most commonly used DMARDs at the onset of RA were methotrexate and sulfasalazine, with their use still increasing, and antimalarial drugs.

Table 1 shows that the use of the alkylating agents, cyclophosphamide and chlorambucil, has slightly decreased. The reason for this might be the effectiveness of methotrexate in hindering the most severe complications of arthritic diseases like AA amyloidosis.⁴ New drugs continue to be needed, as witnessed by the rapid rise in figures for leflunomide.

T Klaukka

Social Insurance Institution, Helsinki, Finland

K Kaarela

Rheumatism Foundation Hospital, Heinola, Finland

Correspondence to: Dr K Kaarela, Rheumatism Foundation Hospital, FIN 18120, Heinola, Finland; kalevi.kaarela@reuma.fi

References

- 1 Aletaha D, Eberl G, Nell VPK, Machold KP, Smolen JS. Practical progress in realisation of early diagnosis and treatment of patients with suspected rheumatoid arthritis: results from two matched questionnaires within three years. *Ann Rheum Dis* 2002;61:630-4.
- 2 Kaipiainen-Seppänen O. Incidence of chronic inflammatory rheumatic diseases in Finland during 1975-1990. Kuopio, Finland: Kuopio University Publications, 1997.
- 3 Miettinen T, Hannonen P, Leirisalo-Repo M, Nissilä M, Kautiainen H, Korpela M, *et al*. Comparison of combination therapy with single-drug therapy in early rheumatoid arthritis: a randomised trial. *Lancet* 1999;353:1568-73.
- 4 Laiho K, Tiitinen S, Kaarela K, Helin H, Isomäki H. Secondary amyloidosis has decreased in patients with inflammatory joint disease in Finland. *Clin Rheumatol* 1999;18:122-3.

Authors' reply

The Finnish data presented by Drs Klaukka and Kaarela, which confirm the trend that we saw in our patient populations¹ as well as in the survey of rheumatologists,² are highly appreciated. They clearly show how drastically the use of disease modifying antirheumatic drugs (DMARDs) has changed over the past decade. Importantly, not only has the type of DMARDs employed most frequently undergone a change, but it appears from the table presented that the frequency of combination therapies may be quite high and that more patients receive DMARD treatment than even a few years ago. Such rapid changes are well in line with the observation we made in our matched surveys. There is another noteworthy aspect, which presumably also applies to the Finnish cohort: in our patients, median methotrexate doses increased from 10 mg (quartiles 7.5; 12.5 mg) in 1995 to 15 mg (10; 20 mg) in 2001, another indication that rheumatologists may be treating patients with rheumatoid arthritis (RA) much more efficiently today than less than a decade ago. This evolution is also supported by the effects of earlier DMARD use, as is also evident from our report.²

Together with the new DMARDs, such as leflunomide, we have the tumour necrosis factor α and interleukin 1 blockers, whose appropriate dosing we also will have to learn

Table 1 Annual number of users of antirheumatic drugs (DMARDs) in Finland 1995-2001

	1995	1996	1997	1998	1999	2000	2001
Oral glucocorticoids	19670	20700	21675	22930	24100	24830	25660
Sulfasalazine	11080	12110	12940	14050	14800	15430	15980
Methotrexate	6800	8490	9870	11530	13310	15090	16470
Azathioprine	2000	2110	2145	2280	2360	2340	2310
Aurothiomalate	6540	6440	6425	6330	6120	5730	5240
Auranofin	2460	2310	2125	2010	1885	1770	1610
Hydroxychloroquine	5780	6480	7530	8740	9760	10910	11840
Cyclosporin	530	920	1200	1450	1585	1660	1600
Penicillamine	940	810	665	550	440	290	230
Chloroquine	1260	1160	1040	960	850	760	650
Cyclophosphamide	240	250	260	260	260	230	200
Chlorambucil	90	105	100	110	115	100	70
Leflunomide	0	0	0	0	0	910	2100

over time. With more agents to come in the near future, we may soon be able to treat patients with RA in a way that previously we could only dream about.

We would like to acknowledge that the reported work was performed in collaboration with the Ludwig Boltzmann Institute of Rheumatology and supported by a grant from the City of Vienna.

D Aletaha, J Smolen

Department of Rheumatology, University of Vienna, Austria

Correspondence to Professor J Smolen;
smj@2me.khl.magwien.gv.at

References

- 1 **Aletaha D**, Smolen JS. The rheumatoid arthritis patient in the clinic: comparing more than 1300 consecutive DMARD therapies. *Rheumatology (Oxford)* 2002;41:1367–74.
- 2 **Aletaha D**, Eberl G, Nell VPK, Machold KP, Smolen JS. Practical progress in realisation of early diagnosis and treatment of patients with suspected rheumatoid arthritis: results from two matched questionnaires within three years. *Ann Rheum Dis* 2002;61:630–4.

Sarcoidosis: TB or not TB?

We read with interest the lesson of the month “Sarcoidosis: TB or not TB?” by Litinsky and colleagues, published in the May 2002 issue of the *Annals of the Rheumatic Diseases*.¹

We, in India, face the reverse problem. With us tuberculosis (TB) is still a scourge. Sarcoidosis in India, though not unknown, is definitely rare in comparison with TB. Because non-caseating granulomas can occur in TB also, a diagnosis of TB is accepted even if acid fast bacilli are not demonstrated or grown in culture. For these reasons, many patients with sarcoidosis are initially treated for TB before the eventual diagnosis is made.²

The risk of a patient receiving corticosteroids developing TB depends on the dose and duration of the corticosteroid treatment. Our group has reported a high incidence of TB in patients with systemic lupus erythematosus receiving corticosteroid treatment.³

The patient in question undoubtedly had both sarcoidosis and TB, the latter probably the result of corticosteroid treatment. In our experience most patients develop TB in the first year of such corticosteroid treatment.³ We advocate routine isoniazid prophylaxis for patients receiving long term corticosteroid treatment.⁴

S D Gaitonde, C Balakrishnan, V R Joshi

P. D. Hinduja Hospital, Veer Savarkar, Marg Mahim Mumbai, Ma, India

Correspondence to: Dr V R Joshi;
dr_vjoshi@rediffmail.com

References

- 1 **Litinsky I**, Elkayam O, Flusser G, Segal R, Yaron M, Caspi D. Sarcoidosis: TB or not TB? *Ann Rheum Dis* 2002;61:385–6.
- 2 **Gaitonde S**, Mittal G, Sule A, Pathan E, Mangat G, Bhajani K, *et al*. Sarcoidosis in a rheumatology clinic J Indian Rheum Assoc 2000;8:5–8.
- 3 **Balakrishnan C**, Mangat G, Mittal G, Joshi V R. Tuberculosis in patients with systemic lupus erythematosus J Assoc Physicians India 1998;46:682–3.
- 4 **Gaitonde S**, Pathan E, Mittal G, Joshi V R. Efficacy of isoniazid prophylaxis in patients with systemic lupus erythematosus receiving long term steroid treatment *Ann Rheum Dis* 2002;61:251–3.

Authors' reply

Gaitonde *et al* raise the interesting issue of the influence of epidemiological data on clinical decisions. The differential diagnosis between tuberculosis and sarcoidosis is not easy and every case should be considered to be tuberculosis until proved otherwise, especially in countries where tuberculosis is endemic. It is less evident in countries where tuberculosis is not common. Likewise, the decision of giving routine isoniazid prophylaxis in patients treated with corticosteroids depends on the probability of exposure to *Mycobacterium tuberculosis*. Although the incidence of tuberculosis is undoubtedly increased in patients receiving corticosteroids in India,¹ justifying the use of routine prophylaxis, this issue is still a matter of controversy in non-endemic areas.² However, the ease of travel around the shrinking “global village” dissolves the boundaries between endemic and non-endemic regions and demands constant surveillance.

I Litinsky, O Elkayam

Tel Aviv Medical Centre, Israel

Correspondence to: Dr O Elkayam;
oribe14@netvision.net.il

References

- 1 **Agrawal PN**, Gupta D, Aggarwal AN, Behara D. Incidence of tuberculosis among patients receiving treatment with oral corticosteroids. *J Assoc Physicians India* 2000;48:881–4.
- 2 **Cline JC**, Davis SM. Risk of infection or reactivation of tuberculosis associated with chronic steroid therapy. *Ann Pharmacother* 1997;31:775–6.

Osteoarthritis and cardiovascular death

Haara *et al* recently published a study assessing epidemiological aspects of osteoarthritis (OA) in Finland.¹ A finding of interest was their identification of OA (in any finger joint) as a predictor of cardiovascular death among men, with the authors suggesting an undetermined metabolic factor as a mechanism.

It may be that the disability conferred by OA in the lower limbs delays presentation of patients with ischaemic heart disease owing to a lack of exertional symptoms. Thus treatment to reduce risk is delayed. Additionally, some patients who also have generalised OA and are less physically active may be at higher risk of cardiac events.²

Occupation and levels of education have been used as surrogates of social class.³ Certain jobs that require repetitive movements or heavier work intensity increase the risk of developing OA,⁴ although occupation is not always linked to the development of hand OA.⁵ Generally, manual workers, whose jobs demand higher physical input, are poorly paid compared with professionals.

Haara *et al* found no link between duration of education and OA of the fingers, but the association between OA and earlier death may, nevertheless, be mediated by an effect of social class. Death due to coronary disease is known to be associated with lower socioeconomic groups. Another Finnish study noted higher incidence of myocardial infarction in those of lower income, with higher pre-hospital, 28 day, and 12 month mortality rates.⁶ These higher rates may reflect differences in prevalence and awareness of cardiac risk factors among the

different socioeconomic groups, which may also contribute to higher cardiovascular death in men with OA.

K A Grindulis, G Bhatia, R Davis, M Sosin, D Connolly, F Khattak

Sandwell and West Birmingham NHS Trust, West Bromwich, UK

Correspondence to: Dr K A Grindulis;
Karl.Grindulis@swbh.nhs.uk

References

- 1 **Haara MM**, Manninen P, Kroger H, Arokoski JP, Karkkainen A, Knekt P, *et al*. Osteoarthritis of finger joints in Finns aged 30 or over: prevalence, determinants, and association with mortality. *Ann Rheum Dis* 2003;62:151–8.
- 2 **Wagner A**, Simon C, Evans A, Ferrieres J, Montaye M, Ducimetiere P, *et al*. Physical activity and coronary event incidence in N Ireland and France. *Circulation* 2002;105:2247–52.
- 3 **Davey Smith G**, Hart C, Hole D, MacKinnon P, Gillis C, Watt G, *et al*. Education and occupational social class: which is the more important indicator of mortality risk? *J Clin Epidemiol Community Health* 1998;52:153–60.
- 4 **National Institutes of Health Conference**. Osteoarthritis: new insights. *Ann Intern Med* 2000;133:635–46.
- 5 **Jones G**, Cooley HM, Stankovich JM. A cross sectional study of the association between sex, smoking and other lifestyle factors and osteoarthritis of the hand. *J Rheumatol* 2002;29:1719–24.
- 6 **Salomaa V**, Niemela M, Miettinen H, Ketonen M, Immonen-Raiha P, Koskinen S, *et al*. Relationship of socioeconomic status to the incidence and prehospital, 28-day, and 1-year mortality rates of acute coronary events in the FINMONICA Myocardial Infarction Register Study. *Circulation* 2000;101:1913–18.

BOOK REVIEW

Principles of glucocorticoid therapy

Eds A N Lin, S A Paget. (85.) London: Arnold, 2002. ISBN 0 340 75934-8.

Glucocorticoid therapy is one of the most widely used treatments in medicine, hence this volume is intended for “internists and the many subspecialists in internal medicine, as well as family physicians, paediatricians, and all who use corticosteroids in their practice”. This looks like an ambitious project, even before one sees the contents page. It draws together contributions written by the editors and 49 physicians, principally their colleagues in New York, USA and Edmonton, Canada. They fall into five sections—overview, basic science, effects in specific body systems (for example, the eye, the gastrointestinal tract), clinical usage (the largest section), and special management considerations. The editors aim to present “a systematic discussion of the use of corticosteroids in different organ systems, their side effects, and their use in special circumstances such as pregnancy and childhood.”

The arrangement of the chapters is disjointed. This follows from the different writing styles and the varied depth of explanation and evidence put forward in support of comments and assertions. But more than this, it is as if the project were started a few years ago but languished while some sections remained unfinished, only to be completed