SCIENTIFIC LETTERS

Increased plasma lipoprotein (a) concentrations in Behcet's disease and its relation to vascular events

Behcet's disease, which is characterised by oral and genital ulcerations and eye inflammation, was discovered by Hulusi Behcet in 1937. Other features include arthritis, thrombophlebitis, neurological abnormalities, and skin lesions. Increased incidence of vascular manifestations has been reported in patients with Behcet's disease and may constitute one of the most important causes of mortality and morbidity in those patients.¹ The disease is characterised by a relapsing inflammatory process of unknown aetiology.

Lipoprotein (a) (Lp(a)) is a cholesterol rich plasma lipoprotein that has emerged as an important risk factor for development of coronary artery disease and stroke. Lp(a) could contribute to lipid deposition on arterial walls. Moreover, the recent finding that apo(a) is analogous to plasminogen suggests the possibility that Lp(a) could be the link between thrombosis and atherosclerosis.^{2,3}

The aim of this study was to measure the plasma concentrations of Lp(a) in patients with Behcet's disease and to find relationships between Lp(a) concentrations and clinical findings including disease activity, ocular involvement, articular symptoms, skin lesions, neurologic involvement, and other lipid parameters.

The study group included 27 patients with Behcet's disease (15 men and 12 women; mean (SD) age 33.1 (10) years, range 19-65 years) and 17 healthy control subjects (10 men and 7 women; mean age 33.5 (12.1) years, range 18-59 years). The groups were matched for factors influencing lipoprotein concentrations such as diet, body mass, and exercise.

Oral aphthous lesions were found in all the patients with Behcet's disease. Ocular involvement was seen in 14 patients, genital ulcerations in 14 patients, and skin lesions (erythema nodosum, pyoderma, folliculitis, skin ulcers, etc) in 12 patients. Arthritis or arthritis-like joint involvement occurred in 12 patients, and neurologic involvement in three patients. At the time of the study, the patients were being treated with either colchicine (n = 5) or non-steroid anti-inflammatory drugs (n = 10), or both (n = 8), the remaining four patients receiving no systemic medication.

Serum total cholesterol was measured by a cholesterol oxidase enzymatic method, triglycerides by a glycerol oxidase enzymatic method, and high density lipoprotein cholesterol (HDL-C) by a cholesterol oxidase enzymatic method in supernatant after precipita-

tion with phosphotungstic acid-magnesium chloride. Apolipoproteins AI and B were determined by an immunoassay method (Sera-Pak Immuno Apo AI (code no. 6821)/Apo B (code no. 6822), Ames, Canada). Low density lipoprotein cholesterol (LDL-C) was calculated by the Friedewald formula. Lp(a) was measured using commercial anti-apo(a) polyclonal capture enzyme linked immunosorbent assay (TintElize Lp(a), catalogue no. 610210; Biopool AB, Umea, Sweden) according to the instructions of the manufacturer; an intra-assay coefficient of variation of 6.1% was obtained at the 31 mg/dl concentration of Lp(a) reference plasma (Biopool, catalogue no. 610142, n = 10).

Data were expressed as mean (SD). Lp(a) concentrations and other lipid parameters were compared by Mann Whitney U test, because of the skewed distributions of these values. Disease duration of the patients with Behcet's disease, cholesterol, triglyceride, LDL, HDL, Apo AI, and Apo B were compared with Lp(a) concentrations using regression analyses.

The results are summarised in table 1. Plasma Lp(a) concentrations in patients with Behcet's disease were significantly higher than in the healthy controls. There were no significant differences in the concentrations of total cholesterol, HDL-C, LDL-C, triglycerides, and apolipoproteins AI and B between patients and controls. As shown in table 1, plasma Lp(a) concentrations were not influenced by disease activity, but were affected by the presence of ocular, neurologic, and articular involvement, and skin lesions. Also, no correlation was observed between plasma Lp(a) concentrations and plasma concentrations of cholesterol, HDL-C, LDL-C, Apo AI, and Apo B, except for triglycerides (r = 0.66, p < 0.01).

Atherogenic, thrombogenic, and inflammatory vascular events increase in patients with Behcet's disease. High concentrations of Lp(a) (> 30 mg/dl) increase the risk of atherogenic and thrombogenic events.

Scott estimated that at plasma Lp(a) concentrations of 30 mg/dl, cellular plasminogen binding of Lp(a) is reduced 20%, thereby decreasing cell fibrinolysis and promoting a procoagulant state.⁴ Previously, Orem and colleagues from Turkey also reported that fluctuations of plasma Lp(a) concentrations with disease activity may be a contributing risk factor in the development of thrombotic complications in patients with Behcet's disease.⁵ In addition, in the present study, significant relations were found between Lp(a) concentrations and neurologic, articular, and ocular involvements and skin manifestations within the study group. These findings may be related to vascular manifestations. Also, Lp(a) concentrations were correlated with triglyceride concentrations.

This study suggests that plasma Lp(a) concentrations increase in patients with Behcet's disease. Lp(a) may therefore play a role in the

WEB TOP 10

www.heartjnl.com

These articles scored the most hits on Heart's web site during May 2000

- 1 Joint British recommendations on prevention of coronary heart disease in clinical practice**
December 1998;80(suppl 2):S1-29
- 2 Management of cardiogenic shock complicating acute myocardial infarction: towards evidence based medical practice**
S G Williams, D J Wright, L B Tan
June 2000;83:621-6 (Review)
- 3 Epidemiology, aetiology, and prognosis of heart failure**
J J McMurray, S Stewart
May 2000;83:596-602 (Education in Heart)
- 4 BNP: soon to become a routine measure in the care of patients with heart failure**
M R Cowie
June 2000;83:617-8 (Editorial)
- 5 The natural history of acute myocardial infarction**
R M Norris
June 2000;83:726-30 (Education in Heart)
- 6 Temporary cardiac pacing**
M D Gammage
June 2000;83:715-20 (Education in Heart)
- 7 Left ventricular free wall rupture: clinical presentation and management**
J Figueras, J Cortadellas, J Soler-Soler
May 2000;83:499-504 (Review)
- 8 Arrhythmogenic right ventricular cardiomyopathy: diagnosis, prognosis, and treatment**
D Corrado, C Basso, G Thiene
May 2000;83:588-95 (Education in Heart)
- 9 The cardiomyopathies: an overview**
M J Davies
Apr 2000;83:469-74 (Education in Heart)
- 10 Coronary angioplasty is like going to the dentist**
B Meier
May 2000;83:491-2 (Editorial)

thrombogenic and atherosclerotic events associated with this disease.

CIHANGIR EREM
TURAN USLU*
ORHAN DEGER†
MEHMET TOSUN*
HALIL KAVGACI

Departments of Internal Medicine,

*Department of Physical Medicine and Rehabilitation,

†Department of Biochemistry,

Karadeniz Technical University Faculty of Medicine,
K. T.Ü. Tıp Fakültesi İç Hastalıkları Anabilim Dalı,
61080, Trabzon, Turkey

- Kabbaj N, Benjelloun G, Gueddari FZ, *et al.* Vascular involvements in Behcet's disease: based on 40 patients records [abstract]. *J Radiol* 1993;74:649-56.
- Scanu AM, Lown RM, Berg K. Lipoprotein (a) and atherosclerosis. *Ann Intern Med* 1991;115:209-18.

Table 1 Mean (SD) Lp(a) concentrations in patients with Behcet's disease and controls, and their relation to the presence or absence of clinical findings

Group	Lp(a) (mg/dl)	Prevalence of Lp(a) > 30 mg/dl
Controls (n=17)	16.9 (12.1) (median 10.7)	17.6
Behcet's disease patients (n=27)	27.2 (21.1)* (median 20.3)	44.4**
	Presence (n)	Absence (n)
Active disease	28.4 (19) (n=10)	26.5 (20.1) (n=17)
Ocular involvement	30.3 (19.2) (n=14)*	24.3 (20.0) (n=13)
Skin lesions	40.7 (43.1) (n=12)*	16.4 (14.1) (n=15)
Arthritis/arthritis	24.7 (15.4) (n=10)*	34.1 (23) (n=17)
Neurologic involvement	35.2 (16.6) (n=3)*	26.2 (20.4) (n=24)

Significantly different from the other group: *p < 0.05; **p < 0.01 (Mann-Whitney U test).

- 3 Erem C, Deger O, Bostan M, *et al.* Plasma lipoprotein (a) levels in Turkish NIDDM patients with and without vascular diabetic complications. *Acta Cardiol* 1999;54:203–7.
- 4 Scott J. Thrombogenesis linked to atherogenesis at last? *Nature* 1989;341:22–3.
- 5 Orem A, Deger O, Cimsit G, *et al.* Plasma lipoprotein (a) and its relationship with disease activity in patients with Behcet's disease. *Eur J Clin Chem Clin Biochem* 1995;33:473–8.

Commentary

Increased concentrations of LP(a) have been reported previously in Behcet's disease patients. Thrombosis, in various forms, is a recognised complication. The present report confirms raised plasma LP(a) concentration in Behcet's patients and suggests that ocular, cutaneous, articular, and neurological manifestations of the disease may be particularly associated with raised plasma LP(a). Whether LP(a) is causally related to the vasculitic nature of the disease is unknown, but it is hypothesised that an atherosclerotic/thrombotic mechanism may be involved in the syndrome.

G F BAXTER
Associate Editor

Troglitazone decreases collagen accumulation in prediabetic stage of a type II diabetic rat model

Diabetic cardiomyopathy can be a major complication of diabetes. This complication is not only present in established diabetes but also in prediabetes. Increased diastolic stiffness develops early in experimental diabetic animals which is associated with interstitial accumulation of connective tissue.^{1,2}

The Otsuka Long-Evans Tokushima Fatty (DM) strain of rat was established as an animal model of congenital diabetes, manifesting stable clinical and pathological features that resemble human non-insulin dependent diabetes mellitus (NIDDM).³ In our previous study we found alteration in left ventricular diastolic function and accumulation of myocardial collagen in prediabetic rats.⁴

Troglitazone, an insulin sensitising thiazolidinedione, improves metabolic deterioration in diabetic animal models, obese subjects, and patients with NIDDM.⁵ The drug has an acute effect on cultured cardiac myocytes, on isolated rat hearts, and prevents high glucose induced insulin resistance in cultured rat fibroblasts.

Our study aimed to examine the long term effects of troglitazone on myocardial collagen content using histopathological methods in prediabetic rats.

Male DM rats and age matched male non-DM rats were randomly divided into two groups, respectively: (1) treated DM (n = 10); (2) untreated DM (n = 10); (3) treated non-DM (n = 10); (4) untreated non-DM (n = 10). Rat chow (Oriental Yeast Co, Tokyo, Japan) with or without 0.2% troglitazone (Sankyo, Japan) was given from the age of 5 weeks up to 15 weeks. Before and after treatment, blood sampling for fasting blood glucose, fasting plasma total cholesterol (cholesterol oxidase-peroxydase method), and fasting plasma triglyceride (glycerokinase-glycerol-3-phosphoxydase method) were obtained after an overnight fast. An oral glucose tolerance test (OGTT), plasma insulin concentration (enzyme linked immunosorbent assay method, Levis insulin kit, Japan), plasma alanine aminotransferase (ALT), and aspartate aminotransferase (AST) were measured at 15 weeks of age.

Figure 1 Histomorphometric analysis at 15 weeks of age in diabetic and non-diabetic rats after treatment with troglitazone. The collagen fibres appear blue in Azan-mallory stain and black in threshold image. In the threshold image, significant differences in left ventricular collagen area percentage in treated DM versus untreated DM, and untreated non-DM versus untreated DM ($p < 0.0001$) were observed.

Histomorphometrical and chemical analysis of collagen were performed as in our previous study⁴ after treatment in all four groups.

Intergroup comparisons were performed with an unpaired t test that was corrected by using a Bonferroni factor for multiple comparisons whereby significance was defined as $p < 0.05/k$, where k = number of comparisons. Group data are expressed as mean (SD).

There were no abnormal values for ALT or AST in any of the four groups. There was no significant difference in fasting blood glucose between treated and untreated groups. Troglitazone treatment decreased fasting plasma triglyceride in diabetic rats (untreated 0.64

(0.17) mmol/l *v* treated 0.26 (0.18) mmol/l, $p < 0.001$). At 15 weeks of age there were significant differences in two hour blood glucose on OGTT (mmol/l) and plasma insulin concentration (ng/ml) between untreated DM (10.4 (1.5) mmol/l, 8.1 (1.5) ng/ml) and untreated-non DM (6.1 (0.4) mmol/l, 3.8 (2.1) ng/ml, $p < 0.001$, respectively). Troglitazone treatment improved glucose tolerance and insulin sensitivity of diabetic rats (two hour blood glucose on OGTT 7.7 (1.1) mmol/l *v* untreated DM, $p < 0.005$; plasma insulin concentration 3.2 (0.4) ng/ml *v* untreated DM, $p < 0.001$).

In both the treated and untreated groups there was no dilation or hypertrophy observed on heart excisions. Troglitazone treat-

ment decreased left ventricular collagen content/dry weight in diabetic rats (1.8 (0.1) mg/g *v* untreated 2.2 (0.3) mg/g), although no significant difference was observed. No effect on non-DM rats was observed (treated 1.8 (0.4) mg/g *v* untreated 1.8 (0.5) mg/g). In histomorphometry, troglitazone treatment decreased left ventricular collagen area percentage in diabetic rats (3.3 (0.9)% *v* untreated 8.1 (1.4)%, $p < 0.0001$) but had no effect on non-DM rats (treated 4.1 (2)% *v* untreated 5.1 (1.6)%) (fig 1).

Troglitazone improves hyperinsulinaemia in diabetic humans and other animals. Hyperinsulinaemia is associated with lipid abnormalities and is an independent risk factor for ischaemic heart disease that may lead to cardiac dysfunction. Fasting plasma total cholesterol in our untreated DM rats was not significantly different from that in untreated non-DM rats, and no atherosclerotic lesion was histopathologically observed in any group. It is clear that at this stage diabetic rats are free from premature atherosclerosis (as shown by histological and serological findings) and hypertension, although these factors may be relevant in those adult rats with NIDDM and cardiac dysfunction. Administration of troglitazone decreased values of collagen content/dry weight and collagen area percentage of left ventricle in DM rats in the present study. Hepatic dysfunction may influence collagen metabolism in multiple organs, including the heart. However, in the present study, troglita-

zone treatment decreased interstitial fibrosis and collagen content without hepatic dysfunction. The improvement in cardiac fibrosis in this study is not the result of side effects of troglitazone and might have played a role in improvement of cardiac function.

Gene expression of the transforming growth factor or enhanced gene expression of collagen might participate in the onset of cardiac fibrosis by stimulating extracellular matrix synthesis.⁴ We speculate that the effect of troglitazone in the present study may have been related to improvement in the metabolic state and the regulation of gene expression of collagen in prediabetic rats. To our knowledge, this is the first study to investigate the effect of troglitazone on left ventricular collagen content in a spontaneous NIDDM rat model. Additional work is required for clarification of the mechanism of troglitazone's effect on collagen content.

L YAO
K MIZUSHIGE
T NOMA
K MURAKAMI
K OHMORI
H MATSUO

Second Department of Internal Medicine,
Kagawa Medical University,
1750-1, Ikenobe, Miki, Kita,
Kagawa 761-0793, Japan
email: katsuamz@kms.ac.jp

1 Regan TJ, Weisse AB. Diabetic cardiomyopathy.
J Am Coll Cardiol 1992;19:1165-6.

- 2 Celetano A, Vaccaro O, Tammaro P, *et al*. Early abnormalities of cardiac function in non-insulin-dependent diabetes mellitus and impaired glucose tolerance. *Am J Cardiol* 1995; 76:1173-6.
- 3 Kawano K, Hirashima T, Mori S, *et al*. Spontaneous long-term hyperglycemic rat with diabetic complications. Otsuka Long-Evans Tokushima Fatty (OLETF) strain. *Diabetes* 1992;41:1422-8.
- 4 Mizushige K, Yao L, Noma T, *et al*. Alteration in left ventricular diastolic filling and accumulation of myocardial collagen in insulin-resistant prediabetic stage of a type II diabetic rat model. *Circulation* 2000;101:899-907.
- 5 Nolan JJ, Ludvik B, Beersden P, *et al*. Improvement in glucose tolerance and insulin resistance in obese subjects treated with troglitazone. *N Engl J Med* 1994;331:1188-93.

Commentary

Thiazolidinedione agents called "glitazones" are being developed for the treatment of non-insulin dependent diabetes mellitus (NIDDM). These drugs stimulate peroxisome proliferator activated receptor- γ (PPAR- γ), an action distinct from those of existing sulfonylurea and biguanide drugs. This report shows that troglitazone, a member of this new class, exerts a myocardial antifibrotic action in a genetic animal model of NIDDM, at an early stage of disease progression. The mechanism of this novel action is unknown but may be relevant to the pathophysiology and treatment of diabetic cardiomyopathy.

G F BAXTER
Associate Editor

IMAGES IN CARDIOLOGY

Emergency DDD pacing using a single lead, balloon tip catheter

A 74 year old woman presented with cardiogenic shock and a ventricular septal defect (VSD), four days after an undiagnosed inferior myocardial infarction. An infero-apical VSD was repaired using a 4 $\times$ 3 cm Dacron patch during emergency cardiopulmonary bypass. Despite postoperative ventilation, balloon pump, and inotropic support she remained hypotensive and oliguric on the third postoperative day. In addition, inappropriate sinus bradycardia resulted in frequent ectopy and eight episodes of ventricular tachycardia/fibrillation, requiring DC cardioversion.

Ventricular demand (VVI) pacing via epicardial temporary pacing leads increased heart rate, but did not improve blood pressure or urine output. Neither VVI pacing, nor intravenous loading with amiodarone, suppressed the ventricular arrhythmia.

Dual chamber (DDD) temporary pacing was achieved using a single lead balloon flotation catheter (panel A: the two ventricular, three atrial and single indifferent superior vena cava (SVC) electrode positions are marked. Inset: 1.0 ml air filled balloon to facilitate "flow directed" catheter positioning.) This electrode catheter allows atrial pacing in DDD mode by overlapping biphasic impulses (OLBI, Biotronik) via a pair of non-contact atrial electrodes. Fluoroscopic screening was used to optimise electrode position (panel B). Reliable DDD pacing using an external generator (Eikos SLD, Biotronik) was maintained at 100 beats per minute for five days with dramatic clinical improvement. There was a 24 mm Hg rise in mean arterial pressure (panel C) enabling withdrawal of inotropic support, recovery of renal function, and a steady clinical improvement. In addition, the ventricular arrhythmia was completely suppressed apart from two further episodes of ventricular tachycardia provoked by testing ventricular thresholds in VVI mode.

J D FERGUSON
N LEVER
Y BASHIR
K M CHANNON