We select the letters for these pages from the rapid responses posted on bmj.com favouring those received within five days of publication of the article to which they refer. Letters are thus an early selection of rapid responses on a particular topic. Readers should consult the website for the full list of responses and any authors' replies, which usually arrive after our selection.

LETTERS

SIDS PAPER

Special report adds nothing new, say paper authors

We were disappointed that Gornall's article¹ was published without a response from us. It added nothing new.^{2,3} Our paper reviewed the first 57 deaths on the CONI (Care Of Next Infant), programme. We reaffirm that nine deaths were inevitable (recognised cause), and of the 48 unexpected deaths, seven were probable homicide.

Professor Emery led the CONI steering group until his death (2000). We did not materially change the cause of death for any case he knew about.

There are differences between an earlier report⁴ and subsequent full paper.² In the *Lancet*, we reported but excluded from analysis deaths over one year and clarified "non-natural" deaths. Four infants who died in bed with their parents were initially classified as "non-natural" as they were possibly accidental and avoidable. They were subsequently categorised as SIDS in line with the CESDI SUDI study.⁵ The "non-natural" group initially also included cases recorded as open verdict; their reclassification is detailed. These revisions account for the difference between 14/44 "non-natural" in our earlier report and 7/57 probable infant homicide in the *Lancet*. No case previously attributed to either NAI or suspected or proven filicide was reclassified.

The assertion that "no explanation for the decision to categorise as natural all 13 deaths for which there was insufficient evidence to reach a conclusion" is a misrepresentation of our text. This states "Eight of these CONI deaths were certified as SIDS. None were attributed to homicide or non-accidental injury." The assertion that, "In five of these

cases a police investigation was under way" is not correct. All investigations were complete before the paper was submitted for publication.

For risk factors even to suggest homicide, their prevalence must be known in all relevant groups.

R G Carpenter honorary professor, Medical Statistics Unit, Department of Epidemiology and Population Science, London School of Hygiene and Tropical Medicine, London WC1E 7HT
bob.carpenter@lshtm.ac.uk

A Waite national co-ordinator, CONI Programme, Academic Unit of Child Health, University of Sheffield, Sheffield Children's Hospital, Sheffield S10 2TH

R C Coombs consultant neonatologist
Neonatal Intensive Care Unit, Jessop Wing, Royal Hallamshire Hospital, Sheffield S10 2SF

C Daman-Willems consultant paediatrician
University Hospital, Lewisham, London SE13 6LH

A McKenzie research fellow, Medical Statistics Unit, Department of Epidemiology and Population Science, London School of Hygiene and Tropical Medicine

J Huber consultant paediatric pathologist (retired), University Hospital for Children and Youth, 3501 CA Utrecht, Netherlands

Competing interests: None declared.

- 1 Gornall J. Was message of sudden infant death study misleading? *BMJ* 2006;333:1165-8. (2 December.)
- 2 Carpenter RG, Waite WJ, Coombs RC, Daman-Willems C, McKenzie A, Huber J, et al. Repeat sudden unexpected and unexplained infant deaths: natural or unnatural. *Lancet* 2005;365:29-35.
- 3 Repeat sudden unexpected and unexplained infant deaths: Correspondence. *Lancet* 2005;365:1137-8.
- 4 Waite A, McKenzie A, Carpenter RG, Coombs R, Daman-Willems C, Emery J. *Report on 5000 babies using the CONI (care of next infant) programme*. London: Foundation for the Study of Infant Death, 1998.
- 5 Fleming P, Blair P, Bacon C, Berry J. Sudden unexpected deaths in infancy. The CESDI SUDI studies 1993-1996. London: Stationery Office, 2000.

Ellipsis marks an important omission

Gornall, in his article about the paper by Carpenter et al on recurrent infant deaths,¹ was right to argue that uncertain data should not be translated into statistics that seem clear cut. But his suggestion that a report by John Emery, the pathologist who initiated the *Lancet* study, upheld the ultra-suspiciousness of "Meadow's law" is curious.

Emery produced the report for Sally Clark's defence in 1999, six months before his death.² He wrote that in families where there were two cot deaths, a third were due to rare natural causes that had been missed at autopsy, and a third were unnatural deaths associated with abuse. The final third were "true" sudden infant death syndrome (SIDS)

in that "no suspicion of unnatural death was found and no natural cause was found."

Although Emery's words make it clear that no basis for suspicion was found in two thirds of recurrent deaths, Gornall writes that Emery reached a "stark" conclusion which he purports to quote: "The occurrence of two unexpected deaths in a family thus raises a definite suspicion of unnatural death which in my experience is confirmed ... in a third of such cases."

The ellipsis is odd. What has Gornall left out? If we check Emery's report we find it was a single word. This is what the sentence said before Gornall edited it: "The occurrence of two unexpected deaths in a family thus raises a definite suspicion of unnatural death which in my experience is confirmed only in a third of such cases."

By omitting the word "only" Gornall changes the emphasis of the entire quotation. Whereas the original implies reservations about Meadow's law, Gornall has adjusted the sense so that he can recruit Emery's conclusion in support of this law.

It is important that medical papers should categorise and report their results scrupulously. But those who seek to criticise them should show an equal degree of scrupulousness.

Richard Webster

Oxford OX2 6TX richardwebster@ntlworld.com

Competing interests: None declared.

- 1 Gornall J. Was message of sudden infant death study misleading? *BMJ* 2006;333:1165-8. (2 December.)
- 2 R v Sally Clark [2000] EWCA Crim 54 at 115-6.

Author's reply

Carpenter et al fail to address the concern of David Hall, former president of the Royal College of Paediatrics and Child Health, that the conclusions of their paper were "seriously misleading." Such defence as they do mount is equally misleading: "We did not materially change the cause of death for any case [Emery] knew about" and "No case previously attributed to either NAI or suspected or proven filicide was reclassified". All true—but beside the point. The question that remains unanswered is why, after Emery's death, all but those seven cases of "NAI or suspected or proven filicide" came to be classified as "natural" deaths.

The authors did, after all, admit that: "We cannot exclude the possibility in our study that some of the 13 cases in which enquiries were not possible were cases of covert homicide." If so, then how can they justify classifying as "natural" all 13 of these unresolved cases—including seven in which parents or their representatives flatly declined to engage with the study? This benign exercise of the benefit of the doubt may be compassionate, but it is unscientific and alone invalidates the paper's findings.

Webster's suggestion that a report by Professor Emery was doctored to support an argument is absurd. There is, as I showed in my article,¹ overwhelming evidence that Emery's view of the proportion of repeat unexpected infant deaths that could be considered to be unnatural was seriously at odds with Carpenter et al's. The key and indisputable point of the passage that Webster attempts to obscure is that Emery believed suspicion was justified in a third of cases where there had been two unexpected infant deaths in a family. Even (or "only") one third is a proportion sufficiently shocking and substantial to suggest that Professor Meadow was rather closer to the truth with his so-called "law" than his many critics are happy to admit—and to confirm that it was wrong to put Emery's name on this paper.

Jonathan Gornall freelance journalist, London E1 7LQ
jgornall@mac.com

Competing interests: JG is author of the special report.

- 1 Gornall J. Was message of sudden infant death study misleading? *BMJ* 2006;333:1165-8. (2 December.)

PRACTICE NURSES

New contract reduces quality of patient-nurse relationship

Derret and Burke raise important questions about the future for primary care nurses and health visitors and the potential for negative effects on patient care.¹ We carried out a small study on how practice nurses perceive the changes in their role since the introduction of the new General Medical Services contract. Nine practice nurses were interviewed individually, from practices in areas of high or low deprivation in Glasgow achieving high or low points on the quality outcomes framework in 2004-5. Transcripts were transcribed verbatim and analysed by using a thematic approach.

The results indicate that practice nurses generally feel that their professional roles and status are developing under the new contract. However, views on incentives

(financial reward) were mixed, with many (even from practices scoring high in the quality outcomes framework) feeling under-rewarded. All reported substantial increases in workload, with a much greater use of information technology and less time to spend with patients. All but one nurse (who had negotiated 30 minute appointments) felt that the new arrangements damaged the nurse-patient relationship, and most nurses reported a decrease in job satisfaction.

Stewart W Mercer senior clinical research fellow, University of Glasgow, Glasgow stewmercer@blueyonder.co.uk

Wendy McGregor nurse partner, Calderwood Practice, Alison Lea Medical Centre, East Kilbride G74 3HW

Competing interests: None declared.

- 1 Derrett C, Burke L. The future of primary care nurses and health visitors. *BMJ* 2006;333:1185-6. (9 December.)

OSTEOPOROSIS

What of Asian, African, and South American patients?

Poole and Compston highlight important points on osteoporosis that practising clinicians need to consider.¹ However, doctors managing patients from ethnic minorities in developed Western countries or developing countries in Asia, Africa, and South America would wish to know if and how recommendations on diagnosis, management, and prognosis apply to their individual patients. For example, in relation to waist circumference—which forms a crucial diagnostic criterion of the metabolic syndrome—the optimal screening cut-off points may be population specific and even in a given population differ according to ethnic grouping.² The relevance of findings on osteoporosis in European subjects need to be tested in other groups of patients. Such findings and recommendations are often applied directly to other ethnic groups, with little consideration given to whether they have the same benefit and, more importantly, if certain interventions may actually cause harm.³

A quick Pubmed search using the term "osteoporosis" and the names of different countries found over 1000 hits for the United Kingdom or the United States, under 10 hits each for Sri Lanka, Pakistan, Nepal, Kenya, or Peru and under 60 hits each for India, Thailand, South Africa, Brazil, or Argentina. Of the reports from the UK and the US, only a handful included patients from non-European backgrounds.⁴ Even if such information is generally sketchy it would not be wise to ignore this aspect completely. The anomaly should be

highlighted prominently in any review on the subject.

An important addition to the "Tips for the non-specialist" section of the review would be to audit the use of osteoporosis related non-drug and drug interventions in different ethnic cohorts. This may allow us to assess the degree of benefit and possible ill effects according to ethnic groupings and move us further ahead in making context specific recommendations.

Suranjith L Seneviratne consultant clinical immunologist Manchester Royal Infirmary, Manchester M13 9WL
suran200@yahoo.co.uk

Padmalal Gurugama registrar in internal medicine, Peradeniya Teaching Hospital, Sri Lanka

Devaka J Fernando consultant endocrinologist, Kings Mill Hospital, Sutton in Ashfield, Nottinghamshire NG17 4JT

Competing interests: None declared.

- 1 Poole KES, Compston JE. Osteoporosis and its management. *BMJ* 2006;333:1251-6. (16 December.)
- 2 Molarius A, Seidell JC, Sans S, Tuomilehto J, Kuulasmaa K. Varying sensitivity of waist action levels to identify subjects with overweight or obesity in 19 populations of the WHO MONICA project. *J Clin Epidemiol* 1999;52:1213-24.
- 3 Matthews HW. Racial, ethnic and gender differences in response to medicines. *Drug Metabol Drug Interact* 1995;12:77-91.
- 4 Mehta G, Taylor P, Petley G, Dennison E, Cooper C, Walker-Bone K. Bone mineral status in immigrant Indo-Asian women. *Q J Med* 2004;97:95-9.

FEEDING IN DEMENTIA

Taste in food also changes

I agree with Hoffer and would only add that often not only the quantity of food but also the taste in food changes in dementia.¹ Significant weight loss marked the beginning of the end of their lives for each of my long-lived parents and parents in law (three of whom had dementia), but once we were able to accept this and stopped trying to "feed them up," food ceased to be such a worry. They each developed idiosyncratic tastes, and my late father happily ate a doughnut a day but almost nothing else for several years. My mother, now in her 90th year, eats only toast, spicy rice dishes, and jelly babies. To her relief we have finally stopped trying to persuade her to vary her diet, and my brother and I now cook and freeze small quantities of what she likes to eat. With this bespoke meals service, my very frail mother continues to live alone in her own home and seems to be meeting her nutritional needs.

Pat Ansell senior lecturer, Department of Health Sciences, University of York, York YO10 5DD
pat.ansell@egu.york.ac.uk

Competing interests: None declared.

- 1 Hoffer LJ. Tube feeding in advanced dementia: the metabolic perspective. *BMJ* 2006;333:1214-5. (9 December.)