PRETEST AND OBJECTIVES

To receive your
credit certificate
immediately for free—

Go to
www.PSYCHIATRIST.com
and complete this
activity online.

Articles are selected for credit designation on the basis of the CME Institute's assessment of the needs of readers of *The Primary Care Companion*, with the purpose of providing readers with a curriculum of CME articles on a variety of topics throughout each volume. There are no prerequisites for participation in this CME activity.

To obtain credit, please study the designated article and complete the Posttest and Registration Form.

CME Objective

After studying the Commentary, you should be able to:

- Evaluate whether treatment algorithms should be incorporated into your clinical practice.

Accreditation Statement

The CME Institute of Physicians Postgraduate Press, Inc. is accredited by the Accreditation Council for Continuing Medical Education to provide continuing medical education for physicians.

Credit Designation

The CME Institute of Physicians Postgraduate Press, Inc. designates this educational activity for a maximum of 1 *AMA PRA Category 1 Credit(s)*TM. Physicians should only claim credit commensurate with the extent of their participation in the activity.

Date of Original Release/Review

This educational activity is eligible for AMA PRA Category 1 Credit through October 31, 2008. The latest review of this material was July 2006.

This pretest is designed to facilitate your study of the material.

1. **Evidence suggests that guideline-driven care produces better outcomes than treatment as usual.**
 - a. True
 - b. False

Pretest answer and Posttest on page 318.

Answers to Vol. 6, No. 5 Posttest 2004

1. c 2. d 3. a 4. b 5. c

Depression continues to be a treatment challenge for many physicians—psychiatrists and primary care physicians alike—in part because of the nature of the disorder, but also because of the wide variety of medications and other treatments available, each with a distinct efficacy and safety profile. One way of negotiating treatment decisions is to use treatment guidelines and algorithms. This Commentary, which appears in the September 2006 issue of The Journal of Clinical Psychiatry (2006;67:1458–1465), provides the primary care clinician with insight into the pros and cons of using treatment algorithms to guide the treatment of depression.

—Larry Culpepper, M.D.

PHYSICIANS DISCUSS THE USE OF ALGORITHMS WHEN DECIDING WHAT TREATMENTS ARE APPROPRIATE FOR PATIENTS WITH NONPSYCHOTIC DEPRESSION.

Depression continues to be challenging to manage, in part because of the myriad of treatment options available to the clinician and the patient. Treatment algorithms can help the clinician decide which treatment strategies to use, but an inherent flaw in algorithms is that they take too long to develop and therefore fall out of date quickly as new treatments are approved and new strategies are studied.

On October 12, 2005, Madhukar H. Trivedi, M.D., a Director of the CME Institute of Physicians Postgraduate Press, Inc. and a nationally recognized expert in the development of treatment algorithms, assembled a group of experts to discuss the use of algorithms in the treatment of nonpsychotic major depressive disorder. Their discussion appears here.

This special Commentary is another in a series of independent projects undertaken by the CME Institute of Physicians Postgraduate Press, Inc. as a service to its members and the broader academic and clinical community. Visit www.psychiatrist.com to listen to audio selections from this discussion.

Faculty affiliations and disclosures appear at the end of this Commentary.

The opinions expressed herein are those of the faculty and do not necessarily reflect the views of the CME provider and publisher.

This Commentary was originally published in *J Clin Psychiatry* 2006;67:1458–1465

Use of Treatment Algorithms for Depression

Madhukar H. Trivedi, M.D.; Maurizio Fava, M.D.; Lauren B. Marangell, M.D.; David N. Osser, M.D.; and Richard C. Shelton, M.D.

THE EFFECTIVENESS OF AND NEED FOR ALGORITHMS IN DEPRESSION

Dr. Trivedi: The effectiveness of treatment algorithms rests on 3 principles. The 3 broad categories requiring special attention are (1) the determination of the need for sequential treatment algorithms; (2) what should be the placement of various treatment interventions in treatment algorithms with sequenced treatments, e.g., at what point should combination treatments and somatic treatments be included; and finally, (3) what the process or procedure is to implement algorithms in the clinical setting, e.g., the use of recently developed measurement-based care.¹

Dr. Osser: It is clear to me that there are significant differences between the likely outcome of people receiving treatment as usual (TAU) as opposed to algorithm-driven treatment. In the Texas Medication Algorithm Project (TMAP),² that difference was seen primarily in the first 3 months. An important factor in that difference was that the care coordinator identified patients who were not doing well and needed to move to the next step in the algorithm. For the next 9 months of the TMAP study, there was no widening of the difference between the algorithm group and the TAU group, suggesting that, from that point on, both groups did reasonably well (Figure 1).

Dr. Trivedi: There is also evidence in primary care from Katon and colleagues,³ which showed that guideline-driven care produces better outcome than TAU. Unützer and coworkers⁴ found similar results. All of these studies have used care coordinators in the algorithm-driven treatment groups. The question becomes whether the better outcomes are related to the use of the care coordinator.

Dr. Shelton: Dr. Trivedi, according to your TMAP article from 2004,² I agree that the first quarter is where the majority of change happens, but if you look at the end of a year of treatment, there still is a substantial difference between the TAU and the algorithm treatment patients. TAU patients did not catch up by the end of a year (see Figure 1), but the differences that are achieved are achieved primarily up front and then maintained throughout the year.

Dr. Marangell: The data are fairly consistent in showing that algorithms result in not just better response rates, but improved patient satisfaction as well. Katon et al.³ found that response rates were 74% with the algorithm versus 44% with TAU and that patients who received algorithm-

Figure 1. Mean Scores on the Inventory of Depressive Symptomatology–Clinician-Rated Scale (IDS-C) During 12 Months of Algorithm-Guided Treatment or Treatment as Usual^a

^aReprinted with permission from Trivedi et al.²
* $p = .002$.

driven treatment were more likely than patients who received TAU to rate their quality of care as good or excellent (93% vs. 75%).

Dr. Fava: If you ask me whether there is a need for treatment algorithms in depression, I would say absolutely, but I question if we truly know what algorithms should look like. We often have to base these algorithms on consensus, expert opinion, or anecdotal impressions rather than scientific evidence, because evidence is lacking in many areas of depression treatment. In addition, the algorithms currently in use have never been tested against each other. One of the problems with most algorithms is their emphasis on monotherapy. Such an emphasis does not reflect clinical practice, where polypharmacy is often used from the start to enhance chances of remission. In my mind, these algorithms that emphasize monotherapy lack ecological validity.

ALGORITHM TREATMENT OPTIONS

Monotherapy Versus Combination Therapy

Dr. Trivedi: Given Dr. Fava's comments, one outstanding question in our field is when is it appropriate to start using combinations and become more aggressive with pharmacotherapy and/or psychotherapy. How far into an algorithm should that step be? The TMAP algorithm for depression,^{2,5} which was developed in the late 1990s, recommends 3 monotherapy steps before turning to combination treatment (Figure 2). Increasingly, clinicians have begun to disagree. Dr. Fava, it sounds like you were making the argument that fewer monotherapy steps should precede combination therapy in a depression algorithm. Should combination therapy be earlier, as a second step or even a first-line treatment?

Dr. Shelton: I agree with Dr. Fava that the evidence for the benefit of multiple sequential monotherapies is limited.

Dr. Marangell: That is a very important point and is separate from whether or not algorithms are useful. Sequential studies, except for projects like the Sequenced Treatment Alternatives to Relieve Depression study (STAR*D),^{6,7} and studies to compare combination treatments are rare, and so we have few data to address these questions.

Dr. Trivedi: Even with STAR*D, the design was to start with a selective serotonin reuptake inhibitor (SSRI) and then go to combination and more complicated treatments (Figure 3).⁸ The subsequent publications from STAR*D^{1,6,7} again highlight the need to be aggressive early on in terms of treatment choices and the use of adequate dose and duration of treatment exposure.

Dr. Marangell: We need to let clinicians know that the recommendation to stick with several rounds of monotherapy, particularly with a single-mechanism agent like an SSRI, is not consensus at this juncture, but the data for our suggestions for more aggressive pharmacotherapy are also limited.

Dr. Fava: It is very difficult for researchers to find support for and conduct studies that compare next-step options like combination treatment, but at the same time, these are the kind of studies that patients and consumers need so that clinicians can have more guidance when making treatment decisions.

Dr. Osser: Absolutely. We can speculate that the TMAP patients did not separate from the TAU patients any further during the last 9 months of the study because they received sequential monotherapy as opposed to combination therapy, but we have no evidence to support that speculation. There are so many options available, for both monotherapy and combination therapy. Quitkin and colleagues⁹ recently suggested that the remission rate could be as high as 95% in completers after 3 sequential monotherapies (Figure 4). That would be tough for combination therapy to beat, so I think it is difficult to have a clear sense that moving earlier to combinations would be appropriate.

Dr. Marangell: A 95% remission rate? That is very high.

Dr. Osser: It was not a controlled study; they examined data from 3 studies and extrapolated data from that pooled data set.

Dr. Fava: It is a very high rate and may be explained by considering that if there is a 50% chance of remission after every trial of monotherapy, 3 sequential monotherapy trials would have a high cumulative rate of remission. We have done a similar study¹⁰ that was truly sequential, and we found that the remission rates were about 20%. If you look at real-world populations, the chances of remission are very low after 3 sequential monotherapies.

Dr. Trivedi: The question of combination treatment versus sequential monotherapy would be difficult to investigate with a randomized trial, but the lack of evidence does not imply that combinations are not effective.

Figure 2. Texas Medication Algorithm Project (TMAP): Strategies for the Treatment of Nonpsychotic Major Depressive Disorder^a

^aThis algorithm is in the public domain and may be found on the Texas Department of State Health Services Web site at <http://www.dshs.state.tx.us/mhprograms/timamdd1algo.pdf>. For more information, see Trivedi et al.¹ and Crismon et al.⁴

Abbreviations: BUPSR = bupropion sustained release, MAOI = monoamine oxidase inhibitor, MRT = mirtazapine, NEF = nefazodone, SSRI = selective serotonin reuptake inhibitor (includes fluoxetine, sertraline, paroxetine, and citalopram), TCA = tricyclic antidepressant, VLFXR = venlafaxine extended release.

Symbols: * = Consider TCA or VLF if not tried, ** = Augment with lithium, thyroid hormone, or buspirone, *** = Skip if lithium augmentation has already failed, † = Most studied combination.

If we look at the TMAP results,² the rationale that you will continue to have additional benefit over the following 9 months with algorithm-driven treatment is valid. One might assume that the TAU group would catch up during the year of treatment if the only differences during the first 3 months were the amount of patient contact and support. However, there was no catch-up in TMAP—even at the end of 1 year, patients who had algorithm-driven treat-

ment had a more robust response than the TAU group (see Figure 1). The major benefit from using algorithm-driven treatment was accrued in the first 3 months, and that benefit was never lost compared with treatment as usual.

Dr. Shelton: You would expect that if the treatment during the first 3 months was more effective, you would have a continuing differential rate of change instead of the leveling off that we see in Figure 1.

Figure 3. Sequenced Treatment Alternatives to Relieve Depression (STAR*D) Algorithm^a

^aReprinted with permission from Rush et al.⁸

Dr. Trivedi: That is exactly where this question of modifying the algorithm so that it is more aggressive comes in—would that difference continue to grow if we use atypical antipsychotics or other combinations earlier in the algorithm? The TMAP results cannot answer that question.

Residual Symptoms and Augmentation Strategies

Dr. Trivedi: If I had to focus on what should be different in a future algorithm, I would start with the questions of whether or not to start combinations earlier, as we just discussed, and then whether or not to identify and engage in targeted treatment of specific residual symptoms—like insomnia, anhedonia, fatigue, and physical symptoms—and if so, with which augmentation strategies? Physicians are often left without clear guidance on how to treat residual symptoms. Does it help to switch antidepressants? Does it help to augment with a different type of agent? Does it help to initially prescribe a treatment that is focused on a common residual symptom? Dr. Fava, you and your group have done a fair amount of work on this topic—would you measure and target specific treatments for specific residual symptoms?

Dr. Fava: The main challenge in my mind is that when it comes to residual symptoms, we rarely really examine residual symptoms in a systematic way. For that reason, there has been a less-than-expected emphasis on the treatment of residual symptoms in algorithms, and we have often settled for incomplete response in clinical practice.

The work of Paykel and coworkers¹¹ as well as Thase and colleagues,¹² among others, has clearly shown that incomplete response is associated with a poorer

Figure 4. Proportion of Intent-to-Treat Patients and Completers in Remission^{a,b}

^aReprinted with permission from Quitkin et al.⁹ Sample 1 (N = 171) and Sample 2 (N = 420); combined (N = 591).

^bFor the percentages shown, the numerator is the number in remission for both samples, and the denominator is the number for both samples who received a new treatment during that period. For example, 165 patients (44 + 121) received a second drug and 80 patients (58 + 22) remitted. Therefore, 48% (80/165) remitted with Treatment 2.

treatment outcome and a greater chance of relapse and recurrence. Their results indicate a need for targeting residual symptoms. The 2 DSM-IV symptoms that are most commonly present as residual symptoms in depression are fatigue and sleep disturbances, which certainly suggests the importance of targeting these common residual symptoms.

On the other hand, there are many other symptoms that are not part of the DSM-IV nomenclature and yet greatly affect the well-being of patients and their quality of life. Those symptoms could also be targeted if we were to measure them systematically. An example is the presence of residual cognitive symptoms—memory, attention problems, concentration problems, problems finding words and so forth—which in my experience are common among patients who have responded to antidepressants but still have residual symptoms.

Dr. Shelton: My colleagues and I have identified at least 2 dominant domains of symptoms in depressed patients, the fatigue/anergia symptoms and the anxiety/worry/rumination types of symptoms, and our data with bupropion¹³ suggests that bupropion is robustly effective for anergia and not differentially effective from placebo for the anxiety-related symptoms. The problem that exists in pure randomized trials is that if we can identify 2 groups of people—anergic and anxious—after the initial phase of treatment and randomly assign them to one treatment or the other, we would probably get a washout of antidepressant effect, simply because some people in each group are going to respond to one treatment and some people to the other treatment.

Dr. Osser: So if you treat a patient with the “wrong” type of medication for his or her symptoms, it may be

unnecessary to add another agent for those symptoms. Instead, you could merely switch the patient to the alternative medication.

Dr. Shelton: In fact, sometimes we do that, and that is part of the quandary here—do you add on, do you switch, do you dramatically change the mechanism of action? Dr. Fava, your earlier point that the monotherapy approach is not always the most effective adds to the difficulty. In the absence of any method or mechanism for targeting symptoms, just changing to a different medication arbitrarily will probably fail to help the majority of patients.

Dr. Osser: We do have data supporting switching from one antidepressant to another, even within the same class; quite a few people do get better.^{14,15} Where is the evidence that you have a better chance of improving people if you add a medication as opposed to switching medication? There have been numerous reviews of that literature, and they conclude that the odds are about the same with those 2 interventions.

Dr. Fava: The sequential data from our group suggest that the chances of remitting with sequential monotherapy are meager.¹⁰ They certainly confirm the impression that many clinicians have that the switch data in the literature involved patients whose level of treatment resistance was not prospectively ascertained, and for that reason it is unclear whether these people were truly resistant or not.

An example is that some of the studies use a historical approach in which the average duration of failed antidepressant trials is approximately 1 year. I find it very hard to believe that a patient would continue in a failed antidepressant trial for 1 year without any augmentation. These may be patients who lost their response and cannot be equated with those to whom Dr. Shelton was alluding, and that suggests, in my mind, that many of the switch studies really involve the switch of relapsers on SSRI treatment to another SSRI, which is a completely different scenario.

A study¹⁶ that my group published recently included combination therapy right off the start with eszopiclone plus fluoxetine versus fluoxetine plus placebo. Monotherapy was associated with significantly fewer remissions and responses than the combination of fluoxetine plus eszopiclone. The Nelson et al. study¹⁷ of monotherapy with desipramine or fluoxetine versus the combination also showed significantly higher remission rates with the combination. I would disagree with the view that monotherapy switches are the best we can offer to our patients.

Dr. Trivedi: Indeed, that is absolutely correct. Another question is, can you benefit from using a dual-mechanism agent at the outset as opposed to a combination of an SSRI and a non-SSRI? I think that is an intriguing question.

Dr. Osser: Dr. Fava, you conducted 2 studies^{18,19} in which you augmented fluoxetine with desipramine or lithium or increased the dose of fluoxetine. In both of

those studies, the results were relatively unimpressive for the augmentation with lithium.

Dr. Fava: I believe that lithium is not effective as an adjunctive agent in unipolar depression. One reason that it may have worked in earlier studies is because many of the patients may have actually had bipolar depression, not unipolar depression. If you look at the study by Nierenberg and coworkers²⁰ of lithium augmentation of nortriptyline, you will see little separation from placebo, and all the most recent studies in unipolar depression have not shown a separation between lithium augmentation and placebo.

The question is, what are effective augmentation strategies? If you use them, do you get a “bigger bang for your buck,” so to speak? I think we are starting to see evidence of that.

Dr. Osser: We have a long history of believing in certain augmentations, and then over time, the data have failed to support those strategies—lithium is a good example. For years, it was considered our premier augmentation. TCA augmentation of an SSRI was held in very high esteem starting with Nelson’s studies,^{17,21} but it has now lost favor. We need more time to be convinced that the current crop of favorites for augmentation is more effective than others, or whether some sampling issue or investigator bias is the reason why they seem to be effective in the preliminary data.

Dr. Trivedi: Dr. Shelton, you have a fair amount of experience with the possibility of using atypical antipsychotics for depression. What is the evidence for that strategy?

Dr. Shelton: There are smatterings of data with the atypical antipsychotics in combination with antidepressants.²² Although the evidence strongly suggests that atypicals are effective, it is by no means definitive, yet the use of atypicals as combination therapy with antidepressants in unipolar patients is apparently common, at least as I have understood recent utilization data.²³ It may be because this particular combination is a new approach or because clinicians are seeing a robust response, but concerns about side effects and long-term adverse outcome may complicate matters. Of course, the costs of these agents can also be prohibitive. Right now, the use of atypicals in depression is probably out of proportion to the level of data that exists in the literature.

Dr. Trivedi: I think that a lot of clinicians are wondering where and how to use atypical antipsychotic agents for the treatment of depression and at what dosages. The recently published results from STAR*D^{1,6,7} also highlight the need to consider alternative options for treatment-resistant depression.

Dr. Shelton: As I understand the utilization data, the vast majority of the growth in atypical use over the last several years has been in mood disorders, including non-psychotic unipolar depression. I am often asked about the

appropriate use of these medications—should they be second-line, should they be third-line, and so on. We lack the answers to those questions.

We may also encounter barriers in trying to gather reliable data on this issue. Studies of augmentation in depression often use a core group of unresponsive patients that are quite unlikely to respond to any medication, and studies with this design may give the appearance that the treatment under investigation is not very effective. In fact, all that is true is that the medication was used as a last resort in patients who may not respond to anything—some studies with the monoamine oxidase inhibitors, for example, have this limitation.

Physical and Somatic Treatments

Dr. Trivedi: Let us turn to the role of somatic treatments, such as electroconvulsive therapy (ECT), vagal nerve stimulation (VNS), and repetitive transcranial magnetic stimulation (rTMS).

Dr. Marangell: rTMS is currently investigational in the United States. More than 60 published trials exist, but most were inadequate in terms of dose, duration, and sample size. Well-designed, adequate-duration trials are currently ongoing. The best-case scenario, if those studies are positive, would be for the rTMS data to go to the U.S. Food and Drug Administration (FDA) for approval in 2006 and for rTMS to become a viable clinical treatment soon after. I see rTMS as an acute treatment. The problem with rTMS is that patients have to go to the clinic every day to have the treatment administered there. For a lot of people, though, the idea of magnetic treatment as opposed to medication or an implanted device is appealing. There is still some work to be done to determine whether it would be a feasible long-term treatment. rTMS is similar to ECT in that ECT remains a highly effective acute treatment, but I would argue that it leaves quite a bit to be desired as a long-term treatment.

VNS is now approved by the FDA for use in adults with chronic and recurrent depression. I see VNS as fitting in very differently than rTMS or ECT. The FDA indication for VNS is 4 failed but adequate antidepressant trials during the patient's lifetime. Since VNS uses an implanted device and the data suggest that the response is seen over a longer period of time, I do not see it as competing with other treatments for an acute-treatment position on a depression treatment algorithm. It would perhaps be more appropriate to include a place in an algorithm for VNS for long-term disease modification.

In the VNS study,²⁴ which included a cohort of highly treatment-resistant patients, 46% of patients responded and 29% were in remission after 1 year of VNS treatment. However, if I had a patient who had been ill for less than 1 year with a first episode of depression and who did not respond to monotherapy, I doubt that I would use an implantable device in that person. However, if I had a patient

who clearly had chronic and recurrent disease such that they were likely to need lifetime interventions to treat and prevent episodes, I would consider VNS.

The question, then, is where VNS would fit into an algorithm. It certainly is not a first-line option, but I would argue that it should not be at the bottom of the list, either.

Dr. Trivedi: Although we have the FDA indication which states that the patient should have failed 4 antidepressant trials before becoming a candidate for VNS, in terms of this discussion, it is important to ask if it is likely that VNS deserves consideration as an earlier option for the kind of patient you just described—the patient with a recurrent, chronic, severe course for whom depression is likely a long-term illness?

Dr. Marangell: I think so. It is also important to keep in mind that VNS is usually used in combination with pharmacotherapy, not as a monotherapy. If you consider the chronic and recurrent patient who has already failed 4 antidepressant trials, you may ask whether you should try to get a response with medication while preparing the patient for VNS. Absolutely. Some people also see the question as whether you should choose ECT or VNS, but to me, those are not mutually exclusive. I see ECT as appropriate for the emergent, acute situation, and you could even implant the VNS device in patients immediately after ECT, with the idea that VNS may not work acutely, but it might help keep someone well and modify the disease course over the long run.

Dr. Shelton: To counter the problem with early relapse following ECT, one could consider the possibility of first implanting the VNS device, doing ECT, and then starting VNS treatment immediately following the ECT course. There is a delay inherent in using VNS—you have to schedule surgery for the implantation and then give the patient time to heal before you can begin the therapy. For patients who have had ECT just prior to this process, this delay may give them enough time to relapse, which would bring you back to square one. If the VNS device is in place from the beginning and then turned on immediately after the course of ECT, it might facilitate the long-term maintenance of response. All of this is speculative, of course, because we simply do not know how feasible this course of action would be, but I think it is worth consideration.

CLINICAL RECOMMENDATIONS AND FACTORS AFFECTING ALGORITHM IMPLEMENTATION

Dr. Trivedi: Until we have the kind of evidence needed to answer some of the questions we have raised today, what clinical recommendations can we make?

Dr. Shelton: The tension between the approaches we have discussed reveals a flaw in our field now, and it explains why so much variation exists in the clinical setting. We have these 2 main strategies for treating depression—

sequential monotherapy versus augmentation and combination therapies—and unfortunately, insufficient data exist for us to make a definitive conclusion regarding which strategy is more effective. We have supporting data for each.

Dr. Marangell: We always have to consider the pros and cons of each approach dependent on the clinical situation and patient preference. For a patient who started with severe symptoms and is now partially better, many clinicians may see more of an advantage in augmentation as opposed to switching. In a patient with mild depression who is a little skittish about medication and who does not want to take more than one medication, a monotherapy switch may be more attractive. Beyond the science, the clinical factors must be taken into account.

Dr. Trivedi: It is beginning to be clear that depression is very difficult to treat and that true remission rates—withstanding some of the open-label trials—in clinical practice are very low, and so we eventually have to think about more complicated treatments and their place—if not as first, second, or third steps—in a treatment algorithm. Moreover, a well-described measurement-based approach that relies on carefully characterized critical decision points that guide clinicians is absolutely crucial.

Dr. Shelton: When we talk about the treatment of depression, we are including at least 2 things—acute response and maintenance of response. When compared with placebo response, the maintenance effect is probably substantially more robust than the acute treatment effect. In a study that my colleagues and I²⁵ conducted, we took patients after 16 weeks of treatment and randomly assigned them to continuation treatment or placebo. The difference in relapse rates was noticeable, so there is a great deal of difference in what happens to patients under those circumstances.

Dr. Osser: I would like to propose a treatment sequence for augmentations. We have a wide array of possible augmentations with some positive evidence, including the atypical antipsychotics, and a huge disparity in the cost of these options. Atypical antipsychotics cost about 3 times as much as brand-name antidepressants and far more than inexpensive generic medications. One way to sequence these augmentations, if we go by the parity of evidence regarding efficacy, is to start with the least costly, which would put the atypicals toward the end of the list.

Dr. Fava: Lithium would be a fairly inexpensive augmentation, but I would still not recommend it. I would be in favor of looking at all the evidence to see where there is a signal of efficacy. That would be preferable than using cost, because basing decisions solely on cost can lead you to medications that may not be the most effective.

Dr. Osser: Cost, side effect burden, and efficacy should all be taken into consideration. I agree with you about lithium; however, if things seem apparently equal,

for example, augmentation with an SSRI or mirtazapine, which are available as generic drugs, compared with an atypical antipsychotic, then cost might be a deciding factor.

Dr. Shelton: You also have bupropion and buspirone and desipramine available as generics. Many medications that have at least some evidence supporting their effectiveness are generic. Most patients would of course prefer an inexpensive drug in the absence of any preferential effect.

Dr. Marangell: Overall, I think there really is a place for algorithms, and one of the keys from a clinician's standpoint is making sure that the algorithms are dynamic and flexible, both in terms of being able to accommodate new data as they come out, but also in terms of accommodating clinician judgment and patient preference.

Drug names: bupropion (Wellbutrin and others), buspirone (BuSpar and others), citalopram (Celexa and others), desipramine (Norpramin and others), eszopiclone (Lunesta), fluoxetine (Prozac and others), lamotrigine (Lamictal), lithium (Eskalith, Lithobid, and others), mirtazapine (Remeron and others), nortriptyline (Aventyl, Pamelor, and others), olanzapine (Zyprexa), paroxetine (Paxil, Pexeva, and others), sertraline (Zoloft and others), tranylcypromine (Parnate and others), venlafaxine (Effexor).

Disclosure of off-label usage: The chair has determined that, to the best of his knowledge, eszopiclone, lithium, olanzapine, and fluvoxamine are not approved by the U.S. Food and Drug Administration for the treatment of depression. If you have questions, contact the medical affairs department of the manufacturer for the most recent prescribing information.

Affiliations: From the Mood Disorders Program and Clinic, Department of Psychiatry, University of Texas Southwestern Medical Center, Dallas (Dr. Trivedi); the Depression Clinical and Research Program, Massachusetts General Hospital and the Department of Psychiatry, Harvard Medical School, Boston (Dr. Fava); the Mood Disorders Center, Department of Psychiatry, Baylor College of Medicine, Houston, Tex. (Dr. Marangell); the Department of Psychiatry, Harvard Medical School, Boston, Mass. (Dr. Osser); and the Department of Psychiatry, Vanderbilt School of Medicine, Nashville, Tenn. (Dr. Shelton).

Faculty disclosure: In the spirit of full disclosure and in compliance with all ACCME Essential Areas and Policies, the faculty for this CME article were asked to complete a statement regarding all relevant financial relationships between themselves or their spouse/partner and any commercial interest (i.e., a proprietary entity producing health care goods or services) occurring within the 12 months prior to joining this activity. The CME Institute has resolved any conflicts of interest that were identified. The disclosures are as follows: **Dr. Trivedi** has received grant research support from the National Institute of Mental Health, Abbott, Bayer, Bristol-Myers Squibb, Eli Lilly, Forest, GlaxoSmithKline, Janssen, Johnson & Johnson, Mead Johnson, Organon, Parke-Davis, Pfizer, Pharmacia & Upjohn, Solvay, Wyeth, and the National Alliance for Research on Schizophrenia and Depression and is a member of the speakers bureau and/or a consultant for Bristol-Myers Squibb, Cyberonics, Eli Lilly, Forest, Janssen, Organon, Pharmacia & Upjohn, Solvay, Wyeth, and Pfizer. **Dr. Fava** has received research support from Abbott, Lichtwer Pharma GmbH, and Lorex; has received honoraria from EPIX, Bayer AG, Compellis, Janssen, Knoll, Lundbeck, Dov, Biovail, BrainCells, Inc., Cypress, Fabre-Kramer, Inc., Grunenthal GmbH, MedAvante, Inc., Sepracor, and Somerset; and has received both research support and honoraria from Aspect Medical Systems, Astra-Zeneca, Bristol-Myers Squibb, Cephalon, Eli Lilly, Forest, GlaxoSmithKline, Johnson & Johnson, Novartis, Organon, Pharmavite, Pfizer, Roche, Sanofi-Synthelabo, Solvay, and Wyeth; and his spouse/partner has received research support from Parke-Davis and Pfizer Inc. **Dr. Marangell** has been a consultant for Cyberonics, Eli Lilly,

GlaxoSmithKline, Medtronic, and Pfizer; has received grant/research support from Bristol-Myers Squibb, Cyberonics, Eli Lilly, and Neuronetics; and has received honoraria from and is a member of the speakers bureau or advisory board for Cyberonics, Eli Lilly, GlaxoSmithKline, and Forest. **Dr. Shelton** has received grant/research support from Abbott, Eli Lilly, GlaxoSmithKline, Janssen, Pfizer, Roehn-Polenc-Rorer, Sanofi, SmithKline Beecham, Wyeth-Ayerst, and Zeneca; has been a consultant for Janssen and Pfizer, and is a member of the speakers bureau for Abbott, Bristol-Myers Squibb, Eli Lilly, Janssen, Pfizer, SmithKline Beecham, Solvay, and Wyeth-Ayerst. **Dr. Osser** has no significant commercial relationships to disclose relative to the presentation.

REFERENCES

- Trivedi MH, Rush AJ, Wisniewski SR, et al. Evaluation of outcomes with citalopram for depression using measurement-based care in STAR*D: implications for clinical practice. *Am J Psychiatry* 2006;163:28–40
- Trivedi MH, Rush AJ, Crismon ML, et al. Clinical results for patients with major depressive disorder in the Texas Medication Algorithm Project. *Arch Gen Psychiatry* 2004;61:669–680
- Katon W, Von Korff M, Lin E, et al. Collaborative management to achieve treatment guidelines: impact on depression in primary care. *JAMA* 1995;273:1026–1031
- Unützer J, Rubenstein L, Katon WJ, et al. Two-year effects of quality improvement programs on medication management for depression. *Arch Gen Psychiatry* 2001;58:935–942
- Crismon ML, Trivedi MH, Pigott TA, et al. The Texas Medication Algorithm Project: report of the Texas consensus conference panel on medication treatment of major depressive disorder. *J Clin Psychiatry* 1999;60:142–156
- Trivedi MH, Fava M, Wisniewski SR, et al. Medication augmentation after the failure of SSRIs for depression. *N Engl J Med* 2006;354:1243–1252
- Rush AJ, Trivedi MH, Wisniewski SR, et al. Bupropion SR, sertraline, or venlafaxine XR after failure of SSRIs for depression. *N Engl J Med* 2006;354:1231–1242
- Rush AJ, Trivedi M, Fava M. Depression, IV: STAR*D treatment trial for depression. *Am J Psychiatry* 2003;160:237
- Quitkin FM, McGrath PJ, Stewart JW, et al. Remission rates with 3 consecutive antidepressant trials: effectiveness for depressed outpatients. *J Clin Psychiatry* 2005;66:670–676
- Petersen T, Papakostas GI, Posternak MA, et al. Empirical testing of two models for staging antidepressant treatment resistance. *J Clin Psychopharmacol* 2005;25:336–341
- Paykel ES, Ramana R, Cooper Z, et al. Residual symptoms after partial remission: an important outcome in depression. *Psychol Med* 1995;25:1171–1180
- Thase ME, Simons AD, McGeary J, et al. Relapse after cognitive behavioral therapy of depression: potential implications for longer courses of treatment. *Am J Psychiatry* 1992;149:1046–1052
- Tomarken AJ, Dichter GS, Freid C, et al. Assessing the effects of bupropion SR on mood dimensions of depression. *J Affect Disord* 2004;78:235–241
- Joffe RT, Levitt AJ, Sokolov ST, et al. Response to an open trial of a second SSRI in major depression. *J Clin Psychiatry* 1996;57:114–115
- Fava M, Dunner DL, Greist JH, et al. Efficacy and safety of mirtazapine in major depressive disorder patients after SSRI treatment failure: an open-label trial. *J Clin Psychiatry* 2001;62:413–420
- Fava M, McCall WV, Krystal A, et al. Eszopiclone co-administered with fluoxetine in patients with insomnia coexisting with major depressive disorder. *Biol Psychiatry* 2006;59:1052–1060
- Nelson J, Mazure C, Jatlow P, et al. Combining norepinephrine and serotonin reuptake inhibition mechanisms for treatment of depression: a double-blind, randomized study. *Biol Psychiatry* 2004;55:296–300
- Fava M, Rosenbaum JF, McGrath PJ, et al. Lithium and tricyclic augmentation of fluoxetine treatment for resistant major depression: a double-blind, controlled study. *Am J Psychiatry* 1994;151:1372–1374
- Fava M, Alpert J, Nierenberg A, et al. Double-blind study of high-dose fluoxetine versus lithium or desipramine augmentation of fluoxetine in partial responders and nonresponders to fluoxetine. *J Clin Psychopharmacol* 2002;22:379–387
- Nierenberg AA, Papakostas GI, Petersen T, et al. Lithium augmentation of nortriptyline for subjects resistant to multiple antidepressants. *J Clin Psychopharmacol* 2003;23:92–95
- Nelson JC, Mazure CM, Bowers MB Jr, et al. A preliminary, open study of the combination of fluoxetine and desipramine for rapid treatment of major depression. *Arch Gen Psychiatry* 1991;48:303–307
- Papakostas GI. Augmentation of standard antidepressants with atypical antipsychotic agents for treatment-resistant major depressive disorder. *Essent Psychopharmacol* 2005;6:209–220
- Cooper WO, Hickson GB, Fuchs C, et al. New users of antipsychotic medications among children enrolled in TennCare. *Arch Pediatr Adolesc Med* 2004;158:753–759
- Marangell LB, Rush AJ, George MS, et al. Vagus nerve stimulation (VNS) for major depressive episodes: one year outcomes. *Biol Psychiatry* 2002;51:280–287
- Hollon SD, DeRubeis RJ, Shelton RC, et al. Prevention of relapse following cognitive therapy vs medications in moderate to severe depression. *Arch Gen Psychiatry* 2005;62:417–422

For the CME Posttest for this article, see pages 318–319.

POSTTEST

Participants may receive a maximum of 1.0 *AMA PRA Category 1 Credit(s)*[™] by reading each CME article and correctly answering at least 70% of the questions in the Posttest that follows:

Go to www.psychiatrist.com/cmehome to take this Posttest online and earn credit immediately.

Or

1. Read each question carefully and circle the answer on the Registration Form.
2. Type or print the registration information in the spaces provided and complete the evaluation.
3. Send the Registration Form to the address or fax number listed on the Registration Form.

All replies and results are confidential. Answer sheets, once graded, will not be returned. Unanswered questions will be considered incorrect and so scored. The CME Institute of Physicians Postgraduate Press, Inc. will keep only a record of participation, which indicates the completion of the activity and the designated number of *AMA PRA Category 1 Credit(s)*[™] that have been awarded. Correct answers to the Posttest will be made available to the participants of this activity upon request after the submission deadline.

Accreditation Statement

The CME Institute of Physicians Postgraduate Press, Inc. is accredited by the Accreditation Council for Continuing Medical Education to provide continuing medical education for physicians.

Answer to Pretest: 1. a

Commentary

pp. 291–298

1. Treatment algorithms are often based on which of the following?

- a. Consensus
- b. Expert opinion
- c. Anecdotal evidence
- d. All of the above

2. It is always clear when to prescribe monotherapy and when to prescribe combination treatment for depressed patients.

- a. True
- b. False

3. Which of the following best summarizes the most effective method for treating residual symptoms in depression?

- a. Switch medications
- b. Augment with a medication that targets the residual symptom in question
- c. Choose an initial medication that targets common residual symptoms like insomnia
- d. There is no single most effective method; the available evidence is unclear

4. In early studies of augmentation strategies for depression, lithium may have appeared effective because:

- a. It is truly effective in unipolar depression
- b. Patients enrolled in the studies may have had very mild unipolar depression
- c. Patients enrolled in the studies may have had bipolar depression
- d. Patients enrolled in the studies had anxiety disorders, not depression

5. In which of the following patients would an augmentation strategy probably be most effective?

- a. A first-episode patient with severe initial symptoms who is only partially better
- b. A patient with a lifetime history of multiple failed antidepressant trials, including both monotherapy and combination therapy
- c. A first-episode patient with mild initial symptoms who is skittish about taking more than 1 medication
- d. A patient with recurring mild depression who also has a history of substance abuse

REGISTRATION FORM

Circle the one correct answer for each question.

1. a b c d 4. a b c d
2. a b 5. a b c d
3. a b c d

Print or type

Name _____

Last 4 digits of Social Security Number _____
(Required to issue CME credit)

Birth Date (mm, dd, yy) _____
(Required to issue CME credit)

Degree _____ Specialty _____

Affiliation _____

Address _____

City, State, Zip _____

Phone () _____

Fax () _____

E-mail _____

☐ Hospital ☐ Private Practice ☐ Resident ☐ Intern

Deadline for submission

For a credit certificate to be issued, please complete this Registration Form no later than October 31, 2008. Online submissions will receive credit certificates immediately. Faxed or mailed submissions will receive credit certificates within 6 to 8 weeks.

Keep a copy for your files

Retain a copy of your answers and compare them with the correct answers, which will be published after the submission deadline.

Payment

If you complete the test online, no payment is necessary. A \$10 payment must accompany this form. You may pay by check, money order, or credit card (Visa or MasterCard). Make check or money order payable to Physicians Postgraduate Press, Inc. If paying by credit card, please provide the information below.

Check one: ☐ Visa ☐ MasterCard

Card number _____

Expiration date _____

Your signature _____

Please evaluate the effectiveness of this CME activity by answering the following questions.

1. Was the educational content relevant to the stated educational objectives? ☐ Yes ☐ No
2. Did this activity provide information that is useful in your clinical practice? ☐ Yes ☐ No
3. Was the format of this activity appropriate for the content being presented? ☐ Yes ☐ No
4. Did the method of presentation hold your interest and make the material easy to understand? ☐ Yes ☐ No
5. Achievement of educational objective:
 - A. Enabled me to evaluate whether treatment algorithms should be incorporated into my clinical practice. ☐ Yes ☐ No
6. Did this CME activity provide a balanced, scientifically rigorous presentation of therapeutic options related to the topic, without commercial bias? ☐ Yes ☐ No
7. Does the information you received from this CME activity confirm the way you presently manage your patients? ☐ Yes ☐ No
8. Does the information you received from this CME activity change the way you will manage your patients in the future? ☐ Yes ☐ No
9. If you answered yes, what change(s) do you intend to make in your practice?

10. Please offer comments and/or suggested topics for future CME activities.

11. How much time did you spend completing this CME activity?

12. What is your preferred format for CME activities? Circle one.
 - A. Print media (e.g., journals, supplements, and newsletters)
 - B. Internet text
 - C. Internet multimedia
 - D. Audio CD
 - E. Live meeting
13. Are you a physician? ☐ Yes ☐ No

TEAR OUT AND SEND THIS PAGE, ALONG WITH YOUR PAYMENT, TO:
CME INSTITUTE • PHYSICIANS POSTGRADUATE PRESS, INC. • P.O. BOX 752870 • MEMPHIS, TN 38175-2870
IF YOU ARE PAYING BY CREDIT CARD, YOU MAY FAX THIS PAGE TO: CME INSTITUTE AT 901-751-3444
QUESTIONS? CALL 1-800-489-1001 EXT. 8 • WWW.PSYCHIATRIST.COM