We select the letters for these pages from the rapid responses posted on bmj.com favouring those received within five days of publication of the article to which they refer. Letters are thus an early selection of rapid responses on a particular topic. Readers should consult the website for the full list of responses and any authors' replies, which usually arrive after our selection.

LETTERS

STENT THROMBOSIS

Antiplatelets alone won't do the job

Gershlick and Richardson say that premature discontinuation of clopidogrel is the greatest risk factor for thrombosis of drug eluting stents.¹ In Iakovou's series thrombosis occurred in five of 17 patients (29%) who prematurely discontinued dual antiplatelet therapy.² One of them discontinued only clopidogrel, and the others stopped both antiplatelets (aspirin and clopidogrel or ticlopidine). We do not know whether continuation of aspirin alone would have been adequate to prevent thrombosis associated with drug eluting stents. In only a minority of patients (five out of 29; 17%) did discontinuation of antiplatelets seem to have a role in the occurrence of stent thrombosis.

The tragedy of drug eluting stents is not the risk of late stent thrombosis but the fact that these devices have only a limited effect on patients' wellbeing. In several randomised trials they did not lead to a lower mortality and did not affect the incidence of acute myocardial infarction.³ Recently presented meta-analyses and data from registries show that they might increase the occurrence of acute infarctions and even late mortality.^{4,5}

Hans Van Brabandt cardiologist
B2800 Mechelen, Belgium van.brabandt@pandora.be
Competing interests: None declared.

- 1 Gershlick AH, Richardson G. Drug eluting stents. *BMJ* 2006;333:1233-4. (16 December.)
- 2 Iakovou I, Mehnan R, Dangas G. Thrombosis after implantation of drug-eluting stents. *Hellenic J Cardiol* 2006;47:31-8.
- 3 Roiron C, Sanchez P, Bouzamondo A, Lechat P, Montalescot G. Drug eluting stents: an updated meta-analysis of randomised controlled trials. *Heart* 2006;92:641-9.
- 4 Nordmann AJ, Briel M, Bucher HC. Mortality in randomized controlled trials comparing drug-eluting vs. bare metal stents in coronary artery disease: a meta-analysis. *Eur Heart J* 2006;27:2784-814.
- 5 Pfisterer M, Brunner-La Rocca HP, Buser PT, Rickenbacher P, Hunziker P, Mueller C, et al. Late clinical events after clopidogrel discontinuation may limit the benefit of drug-eluting stents: an observational study of drug-eluting versus bare-metal stents. *J Am Coll Cardiol* 2006;48:2584-91.

Patient card on discontinuing clopidogrel is available

Gershlick and Richardson mention the use of warning cards that indicate the recommended length of clopidogrel treatment.¹ The United Kingdom Clinical Pharmacy Association (UKCPA) has developed the Barts and the London clopidogrel card, designed to be given to all patients receiving combination treatment after angioplasty.

The card informs the patient of the importance of dual therapy and the reasons why clopidogrel has been initiated, daily dose, the combination with aspirin, together with the planned duration of treatment, and possible adverse effects. It also encourages the patient to show the card to any medical personnel and thereby offers important information regarding the clopidogrel and aspirin treatment to patient and medical personnel, such that doctors and other paramedical personnel do not discontinue therapy without due consideration. The card offers encouragement for further discussion and liaison regarding the patient's treatment with his or her general practitioner, or with the hospital initiating therapy.

We developed the card and have made arrangements for it to be available nationally, in a way similar to the generic warfarin booklet. This card has been endorsed by the British Cardiovascular Society and the British Cardiac Intervention Society. Cards are available free of charge from UKCPA (tel: 0116 2776999 or via email: admin@ukcpa.com).

Sotiris Antoniou principal pharmacist, cardiac services
sotiris.antoniou@bartsandthelondon.nhs.uk
Martin T Rothman professor of interventional cardiology
Barts and the London NHS Trust, London EC1A 7BE
Competing interests: None declared.

- 1 Gershlick AH, Richardson G. Drug eluting stents. *BMJ* 2006;333:1233-4. (16 December.)

Consider also low response to antiplatelets

Gershlick and Richardson suggest that the increased risk of stent thrombosis in drug eluting stents compared with bare metal stents is attributable to prolonged exposure of the stent struts.¹ We suggest that

stent thrombosis is partly attributable to clopidogrel and aspirin "low responders," and that this patient subgroup should be identified before intervention and alternative oral antiplatelet strategies considered.

In multiple trials, 5-40% of patients treated with clopidogrel displayed suboptimal inhibition of platelet aggregation when evaluated ex vivo by optical aggregometry and flow cytometry. These patients are often deemed clopidogrel low responders. A similar phenomenon—aspirin low responders—is reported in 5-20% of cases. The latter group are more likely to be women and have diabetes mellitus and a reduced response to clopidogrel.²

Bedside platelet inhibition assays have been available for some time (for example, VerifyNow P2Y12 assay, Accumetrics USA) and can identify aspirin and clopidogrel low responders beforehand.

We propose that these patients could be treated with increased oral antiplatelet loading and maintenance doses. Several other oral antiplatelet therapies (prasugrel, AZD6140) do not yet have a UK licence but have shown great promise specifically with a low incidence of pharmacodynamic low responders.³

Andrew J Wiper cardiology registrar a_wiper@yahoo.com
Lancashire Cardiac Centre, Blackpool Victoria Hospital,
Blackpool FY3 8NR

David H Roberts consultant interventional cardiologist
Competing interests: None declared.

- 1 Gershlick AH, Richardson G. Drug eluting stents. *BMJ* 2006;333:1233-4. (16 December.)
- 2 Lev EI, Patel RT, Mares KJ, Guthikonda S, Granada J, Delao T, et al. Aspirin and clopidogrel drug response in patients undergoing percutaneous coronary intervention: the role of dual drug resistance. *J Am Coll Cardiol* 2006;47:27-33.
- 3 Jernberg T, Payne CD, Winters KJ, Darstein C, Brandt JT, Jakubowski JA, et al. Prasugrel achieves greater inhibition of platelet aggregation and a lower rate of non-responders compared with clopidogrel in aspirin-treated patients with stable coronary artery disease. *Eur Heart J* 2006;27:1166-73.

MENTAL HEALTH ACT

What not to learn from devolution

Many Scottish psychiatrists do not share Simpson's enthusiasm for the Scottish Mental Health Act.¹ The 2003 act has introduced into the Scottish mental health

service a bureaucracy of previously unknown proportions. Simply allowing a patient to leave the hospital for a few hours requires a nine page form (which asks for their date of birth twice on one page). Tribunals repeatedly postpone their decisions; it is not uncommon for a patient to have three tribunal hearings in one month. Obtaining short term consultant locum cover is virtually impossible, as few agency locums are willing to go through the cumbersome process of obtaining the necessary approval for the small Scottish job market. An adversarial culture has been introduced where previously it did not exist.

The ultimate effect, rather than being one of enhanced patient care, is a system paralysed by legal requirements, while finite resources are diverted away from clinical care. Furthermore, the act itself is an incomprehensible document (even by the standards of comparable statutes). As a result, enormous importance is placed on the practice guidelines, thereby conferring undue power on the authors of those guidelines, undermining democratic accountability.

James Finlayson consultant psychiatrist
jadfinlayson@doctors.net.uk

Daniel Vincent Riordan consultant psychiatrist
New Craigs, Inverness IV3 8NP

Competing interests: None declared.

- 1 Simpson R. Mental Health Act: why not learn from devolution? *BMJ* 2006;333:1221. (9 December.)

UMBRELLA TEST

Urethral dilator is real enough

Judging by the number of new male patients who mention it, the umbrella myth still has great currency in Australia.¹

I think Bamber is correct when he says that the umbrella test may not be an urban myth.² The old urethral dilator complete with turning knob at the top, a dial and pointer, and three steel spines that expanded or retracted depending on how far the knob was turned was a fearsome enough instrument to found a myth of such

Urethral dilator—not an instrument of torture

longevity. I have one in my possession and it looks more like an instrument of torture than a surgical device (figure).

The director of the Melbourne Clinic in the 1970s (my predecessor) told me he believed in using it “as a last resort” in the treatment of recalcitrant non-specific urethritis. He said patients never came back after the treatment so “they must have been cured.” I am unsure whether he ever actually used it, or merely exhibited the instrument to the horrified gaze of the hapless patient.

David L Bradford visiting medical officer
Cairns Sexual Health Clinic, Cairns, QLD 4870, Australia
davidbradford41@hotmail.com

Competing interests: None declared.

- 1 Bradbeer B, Soni S, Ekbote A, Martin T. You're not going to give me the umbrella, are you? *BMJ* 2006;333:1287-8. (23 December.)
- 2 Bamber MG. Not an urban myth. www.bmj.com/cgi/eleletters/333/7582/1287#152021

TRAUMATISED REFUGEES

Refugee status—a yellow flag in managing back pain

The migration of refugees from central London to the outer London suburbs exposes clinicians to the problems of refugees that they may have no experience of, or training in. The recent discussion on *bmj.com* following the editorial by Basoglu presupposes that an appreciation that torture has taken place is understood in primary care and the details made available in any referral to the secondary sector.^{1,2} We have recently reported that only 50% of referral letters to a rheumatology service for refugees complaining of back or neck pain had this information provided.³ Consequently a small number were “triaged” to the physiotherapy service, where, after unsuccessful standard back treatment, they were referred on to the rheumatology service.

As a physician, I would not be so bold as to comment on the diagnosis of post-traumatic stress disorder (PTSD), but when the sleep history clearly identifies nightmares regularly disturbing sleep,⁴ the need for tricyclic antidepressant medication seems appropriate for its sedatory, as well as its pain modulating, benefits.

Usually multiple losses contribute to the clinical picture,² and the effects of torture may also affect other family members. I have seen at least two patients whose experience of spinal pain has clearly

been affected by their spouses' torture and PTSD, again elicited via the sleep history—a crucial element in managing chronic pain.⁵

Andrew O Frank consultant physician in rehabilitation medicine and rheumatology
Arthritis Centre, Northwick Park Hospital, Harrow, HA1 3UJ
andrew.frank1@btinternet.com

Competing interests: AF is medical director of Kynixa, a vocational rehabilitation company.

- 1 Basoglu M. Rehabilitation of traumatised refugees and survivors of torture. *BMJ* 2006;333:1230-1. (16 December.)
- 2 Electronic responses. Rehabilitation of traumatised refugees and survivors of torture. www.bmj.com/cgi/eleletters/333/7581/1230.
- 3 Frank AO. Retrospective study of refugees presenting to a rheumatological service with spinal pain: afraid to go to sleep? *Proceedings of the 11th world congress on pain*, Sydney, August 2005:1457-8.
- 4 McCarthy J, Frank A. Post-traumatic psychological distress may present in rheumatology clinics. *BMJ* 2002; 325:221.
- 5 De Souza LH, Frank AO. Experiences of living with chronic back pain: the physical disabilities. *Disabil Rehabil* (in press).

CAREERS ADVICE

Paper is essential reading for the tall and beautiful

My heartfelt thanks to Trilla et al for publishing their novel research.¹ I was within a whisker of dedicating myself to a specialty that I would not be suited to. I had thought that my analytical mind and desire to understand the patient holistically might have made me a physician. However being dashedly handsome and standing proudly at 1.85 m, I discover that I am wholly unsuited to my intended path.

I am queasy around blood, faint at the sight of retractors, cannot use a knife and fork, nor tie my own shoe laces. I am allergic to scrubs, terrified of taking consent, and more clumsy than any evolved primate should ever be. Should I be a surgeon? Some would say I am overqualified.

Flummoxed, I find myself with three viable options. At present I hope that late nights and early ward rounds might take their toll. Although they will certainly wizen my looks, to hope that they make me shorter are far fetched. I am not sure that surgery is ready for me yet. The career of film star, however, remains a distinct possibility.

Rhys H Thomas medical senior house officer
Gloucestershire Royal Hospital, Gloucester GL1 3NN
Rhys-Thomas@doctors.org.uk

Competing interests: None declared.

- 1 Trilla A, Aymerich M, Lacy AM, Bertran MJ. Phenotypic differences between male physicians, surgeons, and film stars: comparative study. *BMJ* 2006;333:1291-3. (23 December.)