Is a New Immune Response Mediator in the NF- κ B pathway - SUMO-4 - Related to Type 1 Diabetes?

Charles Sia

*Department of Immunology, United Biomedical Inc., 25 Davids Drive, Hauppauge, New York 11788, USA.
E-mail: csia@unitedbiomedical.com.*

Introduction

Type 1 diabetes mellitus (T1DM) is hallmarked by a complete loss of insulin secretion capacity caused by T cell-mediated destruction of pancreatic β -cells [1, 2]. The disorder has a complex pathogenesis involving genetic and environmental factors, it appears impossible hitherto to explain sufficiently how islet abnormalities arise and which mechanisms trigger immune cells to become diabetogenic. However, we already know that a series of immune-responsive mediators are involved in the occurrence of autoimmunity and the damage of β -cells; the most critical mediators are MHC class I and II molecules, cytotoxic enzymes and cytokines, such as interleukin-1 β (IL-1 β) and interferon- γ (IFN- γ) [3]. IL-1 β and IFN- γ , which are potential mediators of islet inflammation and contributors of β -cell death [4], modify the expression of a number of genes in β -cells by themselves. But indirectly these mechanisms are regulated by transcriptional modifiers through pathways involving the nuclear transcription factor NF- κ B [5], the activation of which is regarded as pro-apoptotic in pancreatic β -cells [6, 7]. Elevated activation of NF- κ B has been shown to implicate the destruction of β -cells and the development of T1DM [8]. NF- κ B was further demonstrated to be associated with T1DM [9, 10] and deficiency in this transcriptional regulator even prevents mice from streptozotocin-

induced diabetes [11, 12]. One of the modifiers of NF- κ B is a newly discovered post-transcriptional protein modifier which is, due to its functional and structural similarity with ubiquitin, called the small ubiquitin-like modifier (SUMO).

SUMO proteins control and modify a wide range of processes in eukaryotic cells including protein selection and stabilization. In particular, SUMO transfers ubiquitin-like proteins to target proteins by enzymatic cascades. This process of post-translational protein modification is termed sumoylation [13]. Sumoylation is a regulatory mechanism of protein function that involves signal transduction, glucose transport, tumor suppression and genome surveillance [14, 15]. The proteins that can be sumoylated also include I κ B α , the NF- κ B inhibitor, and the heat shock transcription factors HSF1 and HSF2. Four members of the SUMO family (SUMO1-4) have been identified to date. While SUMO1-3 have a wide tissue distribution, SUMO4 expression is restricted to immune tissues and kidney, making it a candidate mediator of autoimmune disorders such as T1DM [16, 17].

From sumoylation to NF- κ B alteration

How is sumoylation associated with T1DM etiology? The expression of cytokine and other target genes that participate in host immune responses is controlled

by transcriptional regulators of the NF- κ B family [7]. NF- κ B is activated upon phosphorylation and ubiquitylation of the NF- κ B inhibitor protein, I κ B α , which keeps the regulator inactive. Sumoylation with SUMO1 and SUMO4 is able to modify I κ B α to become resistant to proteasome-mediated ubiquitylation and thus to inhibit NF- κ B activity. In contrast, once NF- κ B is activated it triggers the transcription of several immune responsive genes including SUMO4 for the control and negative regulation of immune responses [17]. An impaired SUMO4 function induced by nucleotide polymorphism or mutation is suspected of inducing abnormal alterations in NF- κ B activity and thus contributing to the onset of T1DM [17, 18].

Two research groups have identified a SUMO4 variant, SUMO4*M55V, carrying a valine (V) substitution to replace methionine (M) at the 55th amino acid position of its native counterpart (SUM4*M) [16, 19]. The Met variant carries a change at the protein kinase phosphorylation site at position 55, which is assumed to be critical for the functional activity of SUMO4. *In vitro* studies showed that the expression of the M55 variant is associated with elevated levels of activated HSF transcription factor. No such effect was observed with the expression of the SUMO4*M55V variant. HSF1 and HSF2 are known to regulate the expression of heat shock protein (HSP) genes [20, 21], and enhanced production of HSPs has been observed in pancreatic β -cells exposed to proinflammatory cytokines [6]. It is therefore possible that SUMO4*M conjugated HSF1 and HSF2 are ineffective in mediating down-regulation of HSP generation during insulinitis to protect pancreatic islet cells from undergoing apoptosis. In contrast, no clear effects have been reported for the Val55 allele. While Bohren *et al.* detected no differences in NF- κ B reporter gene expression between both SUMO4 alleles, Gou *et al.* observed a 5.5-fold increase in NF- κ B transcriptional activity associated with the M55V substitution [16, 19]. The SUMO4*M55V gene product binds to one of the NF- κ B inhibitors, I κ B α and renders the sumoylated I κ B α to lose its positive role in shutting down NF- κ B activity following its activation. This condition is thought to drive prolonged transcription of all the genes regulated via NF- κ B activation. Indeed, significant enhancement of IL-12 production by peripheral monocytes/dendritic cells is found in diabetic patients carrying the

SUMO4*M55V variant, and increased IL-12 gene transcription is implicated to contribute to pancreatic β -cell destruction in the NOD mice [16].

Association of SUMO4 with T1DM development

The linkage between NF- κ B and T1DM development has prompted two independent groups, one led by Bohren and the other by Gou and Wang, to seek for polymorphisms of genes within the IDDM5 locus on chromosome 6q25 linked to NF- κ B regulation [16, 19]. Gou *et al.* have found the new polymorphic M55V allele to be associated with T1DM in an ethnically heterogeneous population, including European American and Asian patients and controls, with higher frequency in the G (Val) allele among children of European ancestry in the US as judged by observations made from multiplex and simplex family studies [16]. Interestingly, the same group found a positive association of the A (Met55) allele with T1DM in a British cohort, as did the group of Bohren with the patients and controls they have studied [16, 19]. Two other studies failed to prove the association [22, 23], but another study has confirmed the positive association in a Korean population [24]. The discrepancies in the results of these association studies may be due to several factors, such as sample size, genotypic errors, selection biases, genetic heterogeneity and population differences [16]. In order to rule out genotypic errors Wang *et al.* have extended their study to include 195 more patients and additionally 1,060 control subjects of European ancestry in Florida to confirm the positive association of SUMO4*M55V with T1DM. A further test of independence with the combined Florida data set has consolidated their findings on a significant level ($p = 0.0001$) [25]. Therefore, the authors attribute the discrepancies of the results to genetic heterogeneity due to the different populations analyzed [16].

In conclusion, the identification of the conserved form SUMO4*M and its variant SUMO4*M55V contributes to our understanding of the cause(s) of T1DM. In this regard, further studies are suggestive to unravel the complex mechanism of unbalanced NF- κ B activation that may trigger the destruction of β -cells. Association studies with enlarged sample sizes and heterogeneous populations could reduce the biases and may lead to more consistent results.

References

1. Kay TW, Thomas HE, Harrison LC, Allison J. The beta cell in autoimmune diabetes: many mechanisms and pathways of loss. *Trends Endocrinol Metab* 2000. 11(1):11-15.
2. Kurrer MO, Pakala SV, Hanson HL, Katz JD. Beta cell apoptosis in T cell-mediated autoimmune diabetes. *Proc Natl Acad Sci U S A* 1997. 94(1):213-218.
3. Daniel D, Gill RG, Schloot N, Wegmann D. Epitope specificity, cytokine production profile and diabetogenic activity of

- insulin-specific T cell clones isolated from NOD mice. *Eur J Immunol* 1995. 25(4):1056-1062.
4. **Rabinovitch A.** An update on cytokines in the pathogenesis of insulin-dependent diabetes mellitus. *Diabetes Metab Rev* 1998. 14:129-151.
 5. **Ghosh S, May MJ, Kopp EB.** NF-kappa B and Rel proteins: evolutionarily conserved mediators of immune responses. *Annu Rev Immunol* 1998. 16:225-260.
 6. **Cardozo AK, Heimberg H, Heremans Y, Leeman R, Kutlu B, Kruhoffer M, Orntoft T, Eizirik DL.** A comprehensive analysis of cytokine-induced and nuclear factor-kappa B-dependent genes in primary rat pancreatic beta-cells. *J Biol Chem* 2001. 276(52):48879-48886.
 7. **Hayashi T, Faustman D.** NOD mice are defective in proteasome production and activation of NF-kappaB. *Mol Cell Biol* 1999. 19(12):8646-8659.
 8. **Eizirik DL, Mandrup-Poulsen T.** A choice of death - the signal-transduction of immune-mediated beta-cell apoptosis. *Diabetologia* 2001. 44(12):2115-2133.
 9. **Gylvin T, Bergholdt R, Nerup J, Pociot F.** Characterization of a nuclear-factor-kappa B (NFkappaB) genetic marker in type 1 diabetes (T1DM) families. *Genes Immun* 2002. 3(7):430-432.
 10. **Hegazy DM, O'Reilly DA, Yang BM, Hodgkinson AD, Millward BA, Demaine AG.** NFkappaB polymorphisms and susceptibility to type 1 diabetes. *Genes Immun* 2001. 2(6):304-308.
 11. **Lgssiar A, Hassan M, Schott-Ohly P, Friesen N, Nicoletti F, Trepicchio WL, Gleichmann H.** Interleukin-11 inhibits NF-kappaB and AP-1 activation in islets and prevents diabetes induced with streptozotocin in mice. *Exp Biol Med (Maywood)* 2004. 229(5):425-436.
 12. **Mabley JG, Hasko G, Liaudet L, Soriano F, Southan GJ, Salzman AL, Szabo C.** NFkappaB1 (p50)-deficient mice are not susceptible to multiple low-dose streptozotocin-induced diabetes. *J Endocrinol* 2002. 173(3):457-464.
 13. **Matunis MJ, Pickart CM.** Beginning at the end with SUMO. *Nat Struct Mol Biol* 2005. 12(7):565-566.
 14. **Seeler JS, Dejean A.** Nuclear and unclear functions of SUMO. *Nat Rev Mol Cell Biol* 2003. 4(9):690-699.
 15. **Muller S, et al.** SUMO: a regulator of gene expression and genome integrity. *Oncogene* 2004. 23(11):1998-2008.
 16. **Guo D, Li M, Zhang Y, Yang P, Eckenrode S, Hopkins D, Zheng W, Purohit S, Podolsky RH, Muir A, et al.** A functional variant of SUMO4, a new I kappa B alpha modifier, is associated with type 1 diabetes. *Nat Genet* 2004. 36(8):837-841.
 17. **Li M, Guo D, Isaacs CM, Eizirik DL, Atkinson M, She JX, Wang CY.** SUMO wrestling with type 1 diabetes. *J Mol Med* 2005. 83(7):504-513.
 18. **Kosoy R, Concannon P.** Functional variants in SUMO4, TAB2, and NFkappaB and the risk of type 1 diabetes. *Genes Immun* 2005. 6(3):231-235.
 19. **Bohren KM, Nadkarni V, Song JH, Gabbay KH, Owerbach D.** A M55V polymorphism in a novel SUMO gene (SUMO-4) differentially activates heat shock transcription factors and is associated with susceptibility to type I diabetes mellitus. *J Biol Chem* 2004. 279(26):27233-27238.
 20. **He H, Soncin F, Grammatikakis N, Li Y, Siganou A, Gong J, Brown SA, Kingston RE, Calderwood SK.** Elevated expression of heat shock factor (HSF) 2A stimulates HSF1-induced transcription during stress. *J Biol Chem* 2003. 278(37):35465-35475.
 21. **Alastalo TP, Hellesuo M, Sandqvist A, Hietakangas V, Kallio M, Sistonen L.** Formation of nuclear stress granules involves HSF2 and coincides with the nucleolar localization of Hsp70. *J Cell Sci* 2003. 116:3557-3570.
 22. **Smyth DJ, Howson JM, Lowe CE, Walker NM, Lam AC, Nutland S, Hutchings J, Tuomilehto-Wolf E, Tuomilehto J, Guja C, et al.** Assessing the validity of the association between the SUMO4 M55V variant and risk of type 1 diabetes. *Nat Genet* 2005. 37(2):110-111.
 23. **Qu H, Bharaj B, Liu XQ, Curtis JA, Newhook LA, Paterson AD, Hudson TJ, Polychronakos C.** Assessing the validity of the association between the SUMO4 M55V variant and risk of type 1 diabetes. *Nat Genet* 2005. 37(2):111-112.
 24. **Park Y, Park S, Kang J, et al.** Additional support for a genetic association between SUMO4 and type 1 diabetes in the Korean population. *Nat Genet* (in press).
 25. **Wang CY, Yang P, She JX.** Genetic heterogeneity of the IDDM5 (SUMO4) locus. *Nat Genet* (in press).