Published in final edited form as:

Neurosci Lett. 2007 January 3; 411(1): 1–5.

Decreased blood-brain barrier permeability to fluorescein in streptozotocin-treated rats

Brian T. Hawkins, Scott M. Ocheltree, Kristi M. Norwood, and Richard D. Egleton*

Department of Medical Pharmacology, The University of Arizona College of Medicine, Tucson, AZ 85724

Abstract

Investigations of the blood-brain barrier (BBB) in diabetes have yielded contradictory results. It is possible that diabetes differentially affects paracellular and transcellular permeabilities via modulation of tight junction and transport proteins, respectively. Fluorescein (FL), a marker for paracellular permeability, is a substrate for the transport proteins organic anion transporter (OAT)-3 and multidrug resistance protein (MRP)-2 at the BBB. Furthermore, MRP-2-mediated efflux of FL can be upregulated by glucose. In this study, streptozotocin-induced diabetes led to decreased brain distribution of FL measured by in situ brain perfusion, consistent with activation of an efflux transport system for FL at the BBB. This change was paralleled by increased protein expression of MRP-2, but not OAT-3, in cerebral microvessels. These data indicate that diabetes may lead to changes in efflux transporters at the BBB and have implications for delivery of therapeutics to the central nervous system.

Keywords

blood-brain barrier; diabetes; multidrug resistance protein; organic anion transporter; fluorescein; in situ brain perfusion

Introduction

Diabetes is characterized by multifocal microvascular pathologies. However, the specific effects of diabetes on the blood-brain barrier (BBB) remain controversial. Studies utilizing the streptozotocin (STZ)-induced model of diabetes to assess its effect on the BBB have indicated that the BBB remains intact [4], whereas recent work [29] indicates that BBB permeability is increased in STZ diabetes. Clinical studies of BBB function in diabetes have been similarly disparate [7,38].

One explanation for these conflicting results is that paracellular and transcellular permeabilities of the BBB are differentially regulated in diabetes. Whereas permeability to macromolecules and most small, water-soluble molecules is limited through the paracellular pathway by epithelial-like tight junctions (TJ) [12], transcellular movement of lipid-soluble molecules and organic ions is highly regulated by numerous transport and efflux proteins [27].

Fluorescein (FL) has been used as a marker of paracellular BBB disruption by osmotic shock [13]. However, FL is also a substrate for at least two groups of transporters expressed at the BBB: multidrug resistance proteins (MRP) and organic ion transporters (OAT) [39]. Therefore, the distribution of FL into the brain can be affected by both the integrity of TJ and the

*Correspondence: R.D. Egleton, Dept. of Medical Pharmacology, 1501 N. Campbell Ave, PO Box 245050, Tucson, AZ 85724-5050, USA. Tel: +1 520-626-4997, Fax: +1 520-626-2204, email: rde@email.arizona.edu.

expression/activity of these transporters. Interestingly, MRP-2-mediated efflux of FL in the gut can be stimulated by hyperglycemia [25], indicating that efflux transport may be upregulated in diabetes.

We previously demonstrated that TJ are compromised in STZ diabetes, and that BBB permeability to sucrose is concurrently increased [Hawkins et al., submitted]. In the present study, we have examined the effects of STZ treatment on distribution of FL into the brain using in situ brain perfusion and on the expression of the FL-transporting proteins MRP-2 and OAT-3 in cerebral microvessels.

Materials and Methods

Animals and reagents

Animal protocols used in this study were approved by the University of Arizona Institutional Animal Care and Use Committee and conform to NIH guidelines. Male Sprague-Dawley rats (328 ± 4 g, 2–4 months old) were purchased from Harlan (Indianapolis, IN) and fasted for 6 h prior to i.p. injection of either 60 mg/kg streptozotocin (STZ) in 0.9% saline or saline alone (SAL). Following injection, animals were returned to their cages, maintained under standard 12:12 light dark conditions, and given food and water *ad libitum* for the remainder of the study (14 days for all endpoints). Development of diabetes was confirmed by blood glucose analysis; STZ-treated animals with non-fasting blood glucose < 300 mg/dl at sacrifice (2 out of 56, 4%) were excluded from the study.

Rabbit polyclonal affinity-purified antibody to rat OAT-3 was obtained from Alpha Diagnostic (San Antonio, TX). Mouse monoclonal antibody to MRP-2 was obtained from Alexis (San Diego, CA). All other reagents were purchased from Sigma (St. Louis, MO), unless otherwise stated.

In situ brain perfusion

Animals were anesthetized with 1 ml/kg ketamine cocktail (acepromazine 0.6 mg/ml, ketamine 78.3 mg/ml, and xylazine 3.1 mg/ml) and given heparin (10,000 units/kg, i.p.), then perfused with an oxygenated Ringer's solution containing FL (sodium fluorescein salt, Sigma) via bilateral common carotid artery cannulation as previously described [13]. Cerebral hemispheres were mechanically homogenized in 7.5 % (w/v) trichloroacetic acid (TCA) and neutralized with 5 N NaOH for fluorimetric determination of FL (excitation 485 nm, emission 535 nm). Samples of perfusate were collected from the perfusion circuit, diluted in 7.5% TCA, and measured by the same methods. The ratio of FL in brain to that in the perfusate, R_{br} , was calculated by the following:

$$R_{br}(\mu l / g) = [FL]_{susp} \cdot V_{susp} / [FL]_p \cdot m_{sam} \quad (1)$$

where $[FL]_{susp}$ is the measured concentration of FL in the tissue suspension, V_{susp} is the total volume of the tissue suspension, $[FL]_p$ is the measured concentration of FL in the perfusate, and m_{sam} is the mass of the brain tissue sampled.

Unidirectional transfer coefficients (K_{in}) were calculated from the slope of the least-squares fit to linear regressions of R_{br} versus time (T):

$$R_{br} = K_{in}(T) + A \quad (2)$$

Alternatively, non-linear uptake data was fit to the equation:

$$R_{br} = K_{in} / k_{out} \cdot (1 - e^{-k_{out}T}) \quad (3)$$

where k_{out} is an efflux constant [26] using Sigma Plot 2001 v. 7.0.

Cerebral microvessel isolation and western blots

Cerebral microvessel protein was isolated from freshly harvested brains as previously described [14] and quantified using the bicinchoninic acid method (Pierce, Indianapolis, IL) with BSA as a standard. Samples were frozen at -20°C for use within a week or at -80°C for use at a later time. 40 μg protein was loaded onto Tris-HCl gels (Criterion, Biorad, Hercules, CA), separated (200 V, 60 min) and transferred to PVDF membranes (240 mA, 45 min). Membranes were sealed in plastic and frozen at -20°C until use, blocked 4 h in TBS with 0.5% Tween-20 and 5% nonfat dry milk, and incubated in primary antibody 1 h at RT. Membranes were washed several times with TBS/Tween-20/milk followed by TBS/Tween-20 without milk prior to incubation with HRP-conjugated secondary antibody (Amersham) for 30 min at RT, washed again, developed using enzyme chemiluminescence (ECLplus, Amersham), and visualized on X-ray film. Semiquantitation of scanned films was performed using a Kodak Image Station, with gel staining (Gelcode; Pierce, Rockford, IL) as a loading control. Results are reported as % expression of control. Primary and secondary antibodies were diluted in 0.5% BSA in PBS (anti-OAT-3, 1:500, anti-MRP-2, 1:50, anti-Rb and anti-Ms secondary, 1:4000).

Immunofluorescence microscopy

Animals were anesthetized as described above. Following transcardiac perfusion with 0.9% saline, brains were removed and snap-frozen in 2-methyl butane with dry ice. Brains were stored at -80°C until cutting into 20- μm coronal sections and mounting onto gelatin-coated slides (minimum of 48 hours). All slides were stored at -80°C until use. Slides were brought to room temperature, air dried, fixed in 100% ethanol (acetone was used for MRP-2 per manufacturer's indications) for 10 min, then washed in PBS followed by wash buffer (1% BSA/0.2% Tween-20 in PBS). After blocking 90 min in normal goat serum (Vector Labs, Burlingame, CA) diluted 1:50 in wash buffer, slides were incubated with primary antibody diluted in wash buffer (anti-MRP-2 diluted 1:50, anti OAT-3 diluted 1:500) in humidified chambers overnight at 4°C . After rinsing with wash buffer, slides were incubated with appropriate fluorescent-tagged secondary antibody (Alexafluor488 conjugated anti-Rb or anti-Ms IgG, Molecular Probes, Eugene, OR) diluted 1:500 in wash buffer for 1 h at room temperature, then rinsed with wash buffer and PBS and coverslips were mounted with Vectashield (Vector Labs). Microvessels were visualized on a Zeiss 510 Metaseries laser scanning confocal microscope.

Statistical Analysis

Student's *t* test was used for the comparison of two means (SAL vs. STZ). Linear regression coefficients were compared by methods described in [14].

Results

STZ treatment led to significant changes in blood chemistry by day 14 (Table 1) and in weight (weight changes were $+41 \pm 5$ g, and -16 ± 5 g for SAL and STZ, respectively).

Permeability of the BBB to FL was determined by in situ brain perfusion (Figure 1). In the SAL group, brain entry of FL was limited ($K_{in} = 0.81 \pm 0.12 \mu\text{l g}^{-1} \text{min}^{-1}$) and linear with respect to perfusion time ($R^2 = 0.98$) as previously reported [13]. STZ treatment significantly ($p < 0.01$) decreased brain entry of FL ($K_{in} = 0.35 \pm 0.07 \mu\text{l g}^{-1} \text{min}^{-1}$, $R^2 = 0.90$), determined by comparison of linear regression coefficients [14]. Least-squares fitting of the STZ data to

equation (3) improved the fit ($R^2 = 0.96$) and indicated an efflux component of FL transport not apparent in the SAL group ($k_{out} = 0.43 \pm 0.13 \text{ min}^{-1}$), while K_{in} ($0.93 \pm 0.33 \mu\text{l g}^{-1} \text{ min}^{-1}$) was not significantly different than SAL. Data from the SAL group could not be fit to the nonlinear model, suggesting that efflux of FL at the BBB is negligible in the control condition.

Expression of the FL-transporting proteins MRP-2 and OAT-3 was examined by western blotting of cerebral microvessel extracts [11,18]. The MRP-2 antibody M₂III-6 recognized a single band at approximately 170 kDa in liver extract (LE) [19], and faint band at approximately 150 kDa in cerebral microvessel extracts (Figure 2A). STZ treatment was associated with a 49% increase in MRP-2 expression in cerebral microvessels (Figure 2A). The OAT-3 antibody recognized a single band at 70 kDa in protein extracts from choroid plexus (CP) [40] and cerebral microvessels (Figure 2B). No significant change was observed in protein expression of OAT-3 (Figure 2B). Expression of both MRP-2 and OAT-3 was confirmed in cerebral microvessels by immunofluorescence microscopy of brain slices (Figure 3).

Discussion

FL has been used as a marker for studying BBB permeability for decades [1,15,41]. Though FL is a substrate for efflux transporters at the BBB [17,39], osmotic BBB disruption increases permeability to FL [13], indicating brain distribution of FL can be increased via opening of the paracellular route across the BBB. STZ-induced diabetes is associated with increased paracellular permeability of the BBB and decreased expression and/or degradation of BBB tight junction proteins [6,29]. In this study, brain uptake of FL was decreased in STZ-treated rats compared to controls (Figure 1). Taking previously reported increases paracellular permeability into account, it is unlikely that this observation reflects a tightening of the diffusion barrier to FL.

In STZ-treated animals, FL uptake departs from linear with a flattening of the curve between 3 and 4 min (Figure 1). Normally, due to the high concentration gradient between the perfusate and brain for FL (i.e., $[\text{FL}]_p \gg [\text{FL}]_{\text{brain}}$) we would expect unidirectional transfer (blood to brain) and the plot of uptake versus time to be linear; therefore, this departure from linearity indicates an efflux mechanism [3]. Thus, linear regression analysis would underestimate K_{in} . Previous studies have used fits to equation (3) to calculate K_{in} and k_{out} [26,35]. Fitting the STZ data to equation (3) improved the fit of the STZ data ($R^2 = 0.96$ vs. 0.90 in the linear regression) and indicated no significant change in K_{in} . Rather, an efflux component to FL transport is indicated that is not observable in the control condition, as SAL data could not be fitted to the non-linear model. Furthermore, extrapolation of the linear regression to $t = 0$ suggests an initial volume of distribution > 0 in the STZ group (Figure 1). As the vascular volume has been corrected for by washout of the tracer at the end of the perfusion [13], this curve should intersect very close to the origin, further supporting fitting this data to the non-linear model.

Expression of MRP-2 at the BBB is controversial, due to differential expression patterns among species, strains, and individuals [8]. Studies in postmortem human brain sections have variously indicated no brain immunoreactivity for MRP-2 [31] and heterogeneous expression in brain endothelium and parenchymal cells among individuals [23]. Mice have strain-specific expression patterns of MRP-2, with FVB mice in particular lacking MRP-2 in brain but not liver [37]. MRP-2 has been immunolocalized in rat brain microvessels [28], a finding supported by increased brain distribution of MRP-2 substrates in TR- (MRP-2 deficient) rats [33]. Based on transport studies in isolated cerebral microvessels [28], MRP-2 is thought to be expressed at the luminal membrane of the capillary endothelium, where it is ideally situated to mediate efflux of substrates in a manner analogous to its role at the apical membranes of polarized epithelia in the kidney, liver, and gut [8].

It has been suggested that MRP-2 expression at the rat BBB is induced by status epilepticus but is below detection limits under normal conditions [16]. In this study, immunoreactivity for MRP-2 was observed in our enriched microvessel preparations from control animals but was very faint compared with the positive control (Fig 2A). 14-d STZ treatment increased expression of MRP-2, indicating that diabetes may lead to an upregulation of MRP-2 at the BBB.

Upregulation MRP-2-mediated efflux of FL in response to glucose was observed in preparations of rat jejunum [25]. Though that study examined activity of MRP-2 rather than expression, reactive oxygen species produced by hyperglycemia [5] can upregulate expression of p-glycoprotein and MRP isoforms [2]. In the liver, expression of MRP-2 can be regulated by several nuclear receptors [20,21], which can in turn be regulated by glucose [9] and hyperlipidemia [10]. Both hyperglycemia and hyperlipidemia are hallmarks of the STZ model, and were seen in this cohort (Table 1). Further study is warranted to see these transporters are regulated at the BBB by similar mechanisms.

OAT-3, a member of the solute carrier (SLC) family of transporters, acts in concert with other transporters to mediate vectorial transport of xenobiotics and metabolites from blood to brain [24]. OAT-3 immunoreactivity has been shown at the abluminal membrane of cerebral microvessels [30,32], though there is evidence of luminal expression as well [22]. Western blots for OAT-3 have indicated molecular weights ranging from 40 to 130 kD, depending on tissue, species, and assay conditions [34]. In rat brain capillaries, OAT-3 has been reported at approximately 75 kD [22], similar to the band at approximately 70 kDa we observed (Figure 2B). Though no changes in OAT-3 expression were observed in this study (Figure 2B), this does not eliminate the possibility that OAT-3 activity is upregulated in STZ-treated animals [36]. Alternatively, the increase in MRP-2 activity alone may be sufficient to account for the changes in FL transport observed.

In summary, STZ diabetes leads to upregulation of efflux transport of FL at the BBB paralleled by increased expression of MRP-2. Substrates for MRP-2 include antiepileptic drugs [33] and HIV protease inhibitors [8], suggesting that the brain distribution of these therapeutics to the CNS may be altered in diabetic patients. As transcriptional mechanisms regulating the expression of multiple drug transporters may overlap and interact [2], it is possible that other efflux transporters such as p-glycoprotein or MRP-1 may be altered in diabetes as well. Additional study may prove critical for informed management of CNS therapeutic regimens in diabetic patients.

Acknowledgements

This work was supported by NIDDK grant DK065003 to R.D.E.

References

1. Abraham CS, Deli MA, Joo F, Megyeri P, Torpier G. Intracarotid tumor necrosis factor- α administration increases the blood-brain barrier permeability in cerebral cortex of the newborn pig: quantitative aspects of double-labelling studies and confocal laser scanning analysis. *Neurosci Lett* 1996;208:85–8. [PubMed: 8859896]
2. Bauer B, Hartz AM, Fricker G, Miller DS. Modulation of p-glycoprotein transport function at the blood-brain barrier. *Exp Biol Med* (Maywood) 2005;230:118–27. [PubMed: 15673560]
3. Blasberg RG, Fenstermacher JD, Patlak CS. Transport of alpha-aminoisobutyric acid across brain capillary and cellular membranes. *J Cereb Blood Flow Metab* 1983;3:8–32. [PubMed: 6822623]
4. Bradbury MW, Lightman SL, Yuen L, Pinter GG. Permeability of blood-brain and blood-nerve barriers in experimental diabetes mellitus in the anaesthetized rat. *Exp Physiol* 1991;76:887–98. [PubMed: 1768415]

5. Ceriello A. Oxidative stress and diabetes-associated complications. *Endocr Pract* 2006;12(Suppl 1): 60–2. [PubMed: 16627383]
6. Chehade JM, Haas MJ, Mooradian AD. Diabetes-related changes in rat cerebral occludin and zonula occludens-1 (ZO-1) expression. *Neurochem Res* 2002;27:249–52. [PubMed: 11958524]
7. Dai J, Vrensen GF, Schlingemann RO. Blood-brain barrier integrity is unaltered in human brain cortex with diabetes mellitus. *Brain Res* 2002;954:311–6. [PubMed: 12414115]
8. Dallas S, Miller DS, Bendayan R. Multidrug resistance-associated proteins: expression and function in the central nervous system. *Pharmacol Rev* 2006;58:140–61. [PubMed: 16714484]
9. Duran-Sandoval D, Mautino G, Martin G, Percevault F, Barbier O, Fruchart JC, Kuipers F, Staels B. Glucose regulates the expression of the farnesoid X receptor in liver. *Diabetes* 2004;53:890–8. [PubMed: 15047603]
10. Handschin C, Meyer UA. Regulatory network of lipid-sensing nuclear receptors: roles for CAR, PXR, LXR, and FXR. *Arch Biochem Biophys* 2005;433:387–96. [PubMed: 15581595]
11. Hartz AM, Bauer B, Fricker G, Miller DS. Rapid regulation of P-glycoprotein at the blood-brain barrier by endothelin-1. *Mol Pharmacol* 2004;66:387–94. [PubMed: 15322229]
12. Hawkins BT, Davis TP. The blood-brain barrier/neurovascular unit in health and disease. *Pharm Rev* 2005;57:173–185. [PubMed: 15914466]
13. Hawkins BT, Egleton RD. Fluorescence imaging of blood-brain barrier disruption. *J Neurosci Methods* 2006;151:262–7. [PubMed: 16181683]
14. Hawkins BT, Egleton RD, Davis TP. Modulation of cerebral microvascular permeability by endothelial nicotinic acetylcholine receptors. *Am J Physiol Heart Circ Physiol* 2005;289:H212–9. [PubMed: 15708958]
15. Hoffman HJ, Olszewski J. Spread of sodium fluorescein in normal brain tissue. A study of the mechanism of the blood-brain barrier. *Neurology* 1961;11:1081–5. [PubMed: 13908102]
16. Hoffmann K, Gastens AM, Volk HA, Loscher W. Expression of the multidrug transporter MRP2 in the blood-brain barrier after pilocarpine-induced seizures in rats. *Epilepsy Res* 2006;69:1–14. [PubMed: 16504477]
17. Huai-Yun H, Secrest DT, Mark KS, Carney D, Brandquist C, Elmquist WF, Miller DW. Expression of multidrug resistance-associated protein (MRP) in brain microvessel endothelial cells. *Biochem Biophys Res Commun* 1998;243:816–20. [PubMed: 9500978]
18. Huber JD, Hau VS, Borg L, Campos CR, Egleton RD, Davis TP. Blood-brain barrier tight junctions are altered during a 72-h exposure to lambda-carrageenan-induced inflammatory pain. *Am J Physiol Heart Circ Physiol* 2002;283:H1531–7. [PubMed: 12234806]
19. Johnson BM, Zhang P, Schuetz JD, Brouwer KL. Characterization of transport protein expression in multidrug resistance-associated protein (mrp) 2-deficient rats. *Drug Metab Dispos* 2006;34:556–62. [PubMed: 16204465]
20. Johnson DR, Klaassen CD. Regulation of rat multidrug resistance protein 2 by classes of prototypical microsomal enzyme inducers that activate distinct transcription pathways. *Toxicol Sci* 2002;67:182–9. [PubMed: 12011477]
21. Kast HR, Goodwin B, Tarr PT, Jones SA, Anisfeld AM, Stoltz CM, Tontonoz P, Kliewer S, Willson TM, Edwards PA. Regulation of multidrug resistance-associated protein 2 (ABCC2) by the nuclear receptors pregnane X receptor, farnesoid X-activated receptor, and constitutive androstane receptor. *J Biol Chem* 2002;277:2908–15. [PubMed: 11706036]
22. Kikuchi R, Kusuhara H, Sugiyama D, Sugiyama Y. Contribution of organic anion transporter 3 (Slc22a8) to the elimination of p-aminohippuric acid and benzylpenicillin across the blood-brain barrier. *J Pharmacol Exp Ther* 2003;306:51–8. [PubMed: 12684544]
23. Kubota H, Ishihara H, Langmann T, Schmitz G, Stieger B, Wieser HG, Yonekawa Y, Frei K. Distribution and functional activity of P-glycoprotein and multidrug resistance-associated proteins in human brain microvascular endothelial cells in hippocampal sclerosis. *Epilepsy Res* 2006;68:213–28. [PubMed: 16361082]
24. Kusuhara H, Sugiyama Y. Active efflux across the blood-brain barrier: role of the solute carrier family. *NeuroRx* 2005;2:73–85. [PubMed: 15717059]

25. Legen I, Kristl A. D-glucose triggers multidrug resistance-associated protein (MRP)-mediated secretion of fluorescein across rat jejunum in vitro. *Pharm Res* 2004;21:635–40. [PubMed: 15139520]
26. Lockman PR, Mumper RJ, Allen DD. Evaluation of blood-brain barrier thiamine efflux using the in situ rat brain perfusion method. *J Neurochem* 2003;86:627–34. [PubMed: 12859676]
27. Loscher W, Potschka H. Role of drug efflux transporters in the brain for drug disposition and treatment of brain diseases. *Prog Neurobiol* 2005;76:22–76. [PubMed: 16011870]
28. Miller DS, Nobmann SN, Gutmann H, Toeroek M, Drewe J, Fricker G. Xenobiotic transport across isolated brain microvessels studied by confocal microscopy. *Mol Pharmacol* 2000;58:1357–67. [PubMed: 11093774]
29. Mooradian AD, Haas MJ, Batejko O, Hovsepyan M, Feman SS. Statins ameliorate endothelial barrier permeability changes in the cerebral tissue of streptozotocin-induced diabetic rats. *Diabetes* 2005;54:2977–82. [PubMed: 16186401]
30. Mori S, Takanaga H, Ohtsuki S, Deguchi T, Kang YS, Hosoya K, Terasaki T. Rat organic anion transporter 3 (rOAT3) is responsible for brain-to-blood efflux of homovanillic acid at the abluminal membrane of brain capillary endothelial cells. *J Cereb Blood Flow Metab* 2003;23:432–40. [PubMed: 12679720]
31. Nies AT, Jedlitschky G, Konig J, Herold-Mende C, Steiner HH, Schmitt HP, Keppler D. Expression and immunolocalization of the multidrug resistance proteins, MRP1-MRP6 (ABCC1-ABCC6), in human brain. *Neuroscience* 2004;129:349–60. [PubMed: 15501592]
32. Ohtsuki S, Kikkawa T, Mori S, Hori S, Takanaga H, Otagiri M, Terasaki T. Mouse reduced in osteosclerosis transporter functions as an organic anion transporter 3 and is localized at abluminal membrane of blood-brain barrier. *J Pharmacol Exp Ther* 2004;309:1273–81. [PubMed: 14762099]
33. Potschka H, Fedrowitz M, Loscher W. Multidrug resistance protein MRP2 contributes to blood-brain barrier function and restricts antiepileptic drug activity. *J Pharmacol Exp Ther* 2003;306:124–31. [PubMed: 12663688]
34. Robertson EE, Rankin GO. Human renal organic anion transporters: characteristics and contributions to drug and drug metabolite excretion. *Pharmacol Ther* 2006;109:399–412. [PubMed: 16169085]
35. Smith QR. Brain perfusion systems for studies of drug uptake and metabolism in the central nervous system. *Pharm Biotechnol* 1996;8:285–307. [PubMed: 8791815]
36. Soodvilai S, Chatsudthipong V, Evans KK, Wright SH, Dantzer WH. Acute regulation of OAT3-mediated estrone sulfate transport in isolated rabbit renal proximal tubules. *Am J Physiol Renal Physiol* 2004;287:F1021–9. [PubMed: 15238352]
37. Soontornmalai A, Vlaming ML, Fritschy JM. Differential, strain-specific cellular and subcellular distribution of multidrug transporters in murine choroid plexus and blood-brain barrier. *Neuroscience* 2006;138:159–69. [PubMed: 16361063]
38. Starr JM, Wardlaw J, Ferguson K, MacLullich A, Deary IJ, Marshall I. Increased blood-brain barrier permeability in type II diabetes demonstrated by gadolinium magnetic resonance imaging. *J Neurol Neurosurg Psychiatry* 2003;74:70–6. [PubMed: 12486269]
39. Sun H, Miller DW, Elmquist WF. Effect of probenecid on fluorescein transport in the central nervous system using in vitro and in vivo models. *Pharm Res* 2001;18:1542–9. [PubMed: 11758761]
40. Sweet DH, Miller DS, Pritchard JB, Fujiwara Y, Beier DR, Nigam SK. Impaired organic anion transport in kidney and choroid plexus of organic anion transporter 3 (Oat3 (Slc22a8)) knockout mice. *J Biol Chem* 2002;277:26934–43. [PubMed: 12011098]
41. Tervo T, Joo F, Palkama A, Salminen L. Penetration barrier to sodium fluorescein and fluorescein-labelled dextrans of various molecular sizes in brain capillaries. *Experientia* 1979;35:252–4. [PubMed: 421851]

Figure 1. *In situ* brain perfusion with sodium fluorescein

Data are mean $R_{br} \pm \text{S.E.M.}$, $n = 6-11$ per time point. Closed circles = SAL; open circles = STZ; solid lines = linear regressions fit to both data sets; dotted line = nonlinear regression of STZ data fit to Equation 3.

Figure 2. Protein expression of MRP-2 and OAT-3

Shown are summary data and representative western blots for MRP-2 (A) and OAT-3 (B) in cerebral microvessels isolated from SAL (solid bar) and STZ (open bar) animals. Data are mean % relative expression normalized to SAL $\pm$ S.E.M., $n = 6$ per treatment group. Consistency of protein loading was confirmed by gel staining. LE: liver extract (positive control for MRP-2); CP: choroid plexus (positive control for OAT-3). ** $p < 0.01$.

Figure 3. Immunofluorescence of MRP-2 and OAT-3 in brain slices

Shown are representative images at 40x of cerebral sections stained for MRP-2 (A) and OAT-3 (B). Images were taken at 30% laser power, gain 500–700, pinhole = 66 μm . Localization to the vascular/perivascular region was confirmed by costaining with vascular markers ZO-1 and occludin (data not shown).

Table 1

Blood chemistry data.

	SAL	STZ
glucose	242 ± 9	546 ± 12 ***
ketones	6.1 ± 0.5	10.1 ± 0.8 ***
triglycerides	54 ± 6	180 ± 24 ***
cholesterol	85 ± 1	113 ± 3 ***
HDL	32 ± 2	52 ± 2 *
LDL	38 ± 2	31 ± 3

Data are mean blood concentration (mg/dl) ± S.E.M.

*
= p < 0.05 vs. SAL.***
= p < 0.001 vs. SAL.