these were aged over 60 years and two men were younger. Three women and three men died of circulatory failure at the intensive care ward. Two of the women were unconscious on admission, and three men died postoperatively. Of those one was aged 33 and the other 38 years, and they died of pancreatic abscess and haemorrhage respectively. The third man died of circulatory failure 20 days after recovering from his pancreatitis.

The mortality in this series was high in the first five years, but no patient died during 1969-70, probably owing to the fact that peritoneal dialysis was started earlier and the patients were not so ill on admission. During 1971 and 1972 we have so far treated 15 cases of acute haemorrhagic pancreatitis, of whom three have died. Since this regime was introduced there have been some cases of pancreatic cyst formation which have required surgical interference. A remarkable improvement in the conditions of the patients has always occurred and the pain has usually disappeared or decreased shortly after the start of the dialysis.

The mode of action of peritoneal dialysis in acute pancreatitis is not quite clear. Several factors are probably involved—a removal of oedema and pancreatic exudate with toxic substances; correction of electrolytes and blood gases; relieving the load on the kidneys by removing toxic substances from the blood; and supply of calories by absorption of glucose from the dialysing fluid.—I am, etc.

JON GJESSING

Department of Anaesthetics,
Sundsvall Hospital,
Sweden

- 1 Gjessing, J., and Nilsson, L., *Läkartidningen*, 1965, **62**, 3706.
- 2 Gjessing, J., and Dencker, H., *Acta Chirurgica Scandinavica*, 1968, **134**, 351.
- 3 Gjessing, J., and Jacobs, S., *Nordisk Medicin*, 1971, **86**, 927.
- 4 Gjessing, J., Oskarsson, B. M., Tomlin, P. J., and Brock-Utne, J., *British Medical Journal*, 1972, **1**, 617.

Foreign Exchange Visits

SIR,—While it is common for medical specialists to have travelling clubs we believe that visits to other countries under the auspices of postgraduate medical education centres are something new, and readers may be interested to know of two successful exchange visits we have carried out.

In 1969 we asked the Danish Medical Association whether doctors from a town or towns similar in population to ours would be interested in visiting us for four or five days. As a result a group of 26 doctors and their wives came from Esbjerg and Slagelse. These two Danish towns have neither medical schools nor universities but have well developed medical services. Each Danish couple stayed with an English family matched so far as possible for age, interest, or specialty.

The programme consisted of lectures on various subjects, an explanation of the evolution and functioning of the National Health Service, discussions on current world problems such as abortion and population control, on the development of coronary and intensive care units. There was, of course, an energetic social programme.

In 1970 we were invited to Denmark. So successful was this that this year we invited doctors from Holland, and 21 Dutch doctors

with their wives came from Hengelo and stayed five days. Once again this was highly successful.

We are left with the impression that these visits are not only extremely pleasurable but promote international goodwill and understanding. Our visitors were impressed with the development of postgraduate medical education in Britain. For our part our overall impression is that our outlook on medical care is very similar to that of the Danes and the Dutch. If other area clinical tutors are thinking of doing something on these lines we will be happy to give further information on how we managed these exchange visits.—We are, etc.,

I. S. HODGSON-JONES

H. J. VOSS

Kettering and District Hospital,
Kettering, Northants

Respiratory Distress Syndrome

SIR,—Your leading article on the respiratory distress syndrome (7 October, p. 2) raises interesting possibilities. The close correlation of the results of prenatal examination of amniotic fluid with the probability of developing respiratory distress syndrome¹⁻³ would suggest that its onset in an infant should not come as a surprise to the medical attendants. Though it may often be possible to postpone the timing of elective interruption of pregnancy there will still be cases where the child is delivered before term, and even in infants of 35 weeks gestation or more the mortality from respiratory distress syndrome is 10%.⁴

Though surfactant deficiency is an important factor in the cause of respiratory distress syndrome⁵ there is confusion about the reasons for the deficiency. As you point out, immaturity is one factor but failure of synthesis is unlikely to be the only cause, since most infants who subsequently die of respiratory distress syndrome are in good condition at birth^{4,5} and infants who die of respiratory distress syndrome associated with maternal diabetes have no gross surfactant deficiency at the time of death.⁷ Since deficiency of surfactant is also a feature of oxygen toxicity, and since the lung pathology of oxygen toxicity and respiratory distress syndrome are similar,⁶ it has been suggested^{8,9} that the development of respiratory distress syndrome is due to a pathological response to the oxygen in the normal atmosphere.

This problem could be approached from two routes—either by altering the inspired gases or by supporting the mechanisms which protect the lungs against the effects of oxygen. It has been shown that in established respiratory distress syndrome by applying continuous positive pressure to the airway the oxygen tension in the inspired gases can be reduced towards normal while maintaining the arterial oxygen level.¹⁰ If this method was applied early it might allow the oxygen tension in the inspired gases to be reduced below the concentration in the atmosphere. Recent work⁹ on the role of the second gas effect on the onset of pulmonary oxygen toxicity found that helium, the lightest gas, accentuated the oxygen toxicity while the heaviest gas, sulphur hexafluoride, showed the greatest amelioration. This work suggests an alternative possibility of modifying or preventing the disease by varying the non-oxygen component to a composition less likely to damage the abnormal lungs.

In exposure to hyperbaric oxygen ascorbic acid has a protective action against the toxic pulmonary effects of oxygen,¹¹ and indeed the ascorbic acid appears to be mobilized to the lungs and destroyed by oxygen.¹² Pentobarbital and other anaesthetic agents have a protective effect against lung lesions which develop in association with seizures induced by hyperbaric oxygen.¹³ Sympatholytic agents also have a protective action against pulmonary lesions, but in contrast to the anaesthetic agents they have no effect on the seizures.¹³ Since these agents protect a lung with normal protective mechanisms against the effects of a grossly raised inspired oxygen tension they may be applicable to the lung predisposed to the onset of the respiratory distress syndrome because of immature antioxidant mechanisms.—I am, etc.,

E. L. LLOYD

Royal Infirmary,
Edinburgh

- 1 Bhagwanani, S. G., Fahmy, D., and Turnbull, A. C., *Lancet*, 1972, **1**, 159.
- 2 Clements, J. A., et al., *New England Journal of Medicine*, 1972, **286**, 1077.
- 3 Whitfield, C. R., Chan, W. H., Sproule, W. B., and Stewart, A. D., *British Medical Journal*, 1972, **2**, 85.
- 4 Scopes, J., *British Journal of Hospital Medicine*, 1970, **3**, 579.
- 5 Reynolds, E. O. R., Robertson, N. R. C., and Wigglesworth, J. S., *Pediatrics*, 1968, **42**, 758.
- 6 Reynolds, E. O. R., *American Journal of Obstetrics and Gynecology*, 1970, **106**, 780.
- 7 Boughton, K., Gandy, G., and Gairdner, D., *Archives of Disease in Childhood*, 1970, **45**, 311.
- 8 Prathap, K., *Lancet*, 1972, **1**, 745.
- 9 Shanklin, D. R., and Lester, E. P., *Biology of the Neonate*, 1972, **20**, 140.
- 10 Gregory, G. A., et al., *New England Journal of Medicine*, 1971, **284**, 1333.
- 11 Jamieson, D., and Van der Brenk, H. A. S., *Biochemical Pharmacology*, 1964, **13**, 159.
- 12 Shanklin, D. R., Cunningham, J. J., Sotelo-Avila, C., and Crussi, F., *Archives of Pathology*, 1967, **84**, 451.
- 13 Wood, J. D., in *Physiology and Medicine of Diving and Compressed Air Work*. Ed. P. B. Bennett and D. H. Elliott. London, Bailliere, 1969.

Duodenogastric Reflux and Pyloric Surgery

SIR,—Much discussion goes on about whether pyloric reflux or gastric retention causes benign lesser curve gastric ulcer. I was interested to read that Dr. C. D. T. MacLean (14 October, p. 113) found pyloric reflux in 40% of patients with a normal barium meal but a higher incidence in those with gastric ulcer. Du Plessis,¹ who supported the reflux hypothesis, concluded that gastric resection must be done for gastric ulcer. So also did Mr. W. M. Capper (personal communication).

Those who believe that gastric ulceration is caused by retention secondary to duodenal ulceration or pyloric channel disease are led immediately to vagotomy—and indeed to highly selective vagotomy without drainage—for all benign gastric ulcers. I have treated 150 patients with benign lesser curve gastric ulcer by vagotomy since 1962. In 31 of these highly selective vagotomy without drainage has been used. Our results are the same as those of Johnston and his colleagues² in Leeds. The operation seems to be as curative for gastric ulcer as it is for duodenal.

Many of the lesser curve gastric ulcer specimens in the pathology museums in London show obvious evidence of pyloric channel disease. The important work of Liebermann-Meffert and Alloway³ in Basle confirms this association. They describe all the known pathological features of pyloric channel disease in their specimens. No doubt

basal duodenal disease and pyloric channel disease may lead to pyloric reflux. The deformed fibrotic pyloric opening in advanced pyloric channel disease is shown in the Fig.—I am, etc.,

HAROLD BURGE

West London Hospital,
London W.6

- ¹ Du Plessis, D. J., *South African Medical Journal*, 1960, 34, 101.
- ² Johnston, D., Humphrey, C. S., Smith, R. B., and Wilkinson, A. R., *British Journal of Surgery*, 1972, 59, 787.
- ³ Liebermann-Meffert, D., and Allgower, M., *Archives Françaises des Maladies de L'Appareil Digestif*, 1972, 61, Suppl. p. 284C.

Chlamydia in Chronic Prostatitis

SIR,—As you pointed out (1 July, p. 1), it is rarely possible to trace the causative agent of chronic prostatitis. Lately TRIC agent has been suspected as a fairly common cause of non-specific genital infections.^{1,2} We studied sera from 79 untreated patients with chronic or subchronic prostatitis and 72 age-matched registered male blood donors for complement-fixing antibodies to chlamydia. It is known that an antibody response to infections with chlamydia may be detected by complement fixation tests to shared group antigens. We used a psittacosis antigen (Wellcome). Complement-fixing antibodies in a titre of at least 1:5 were found in 33% of the patients with prostatitis but in only two of the blood donors (see Table).

Cultures of urethral swabs and of expressed prostatic and seminal fluids from the patients with prostatitis almost always gave growth only of such bacteria as normally constitute the flora in the distal urethra. None grew gonococci, nor did cultures of rectal swabs. But immuno-fluorescent studies of seminal smears suggested infection with gonococci in five of the cases. In a few cases it seemed that *Candida albicans*, *Corynebacterium vaginalis*, mycoplasmas, *Trichomonas vaginalis*, or a virus might be the causal agent.

Treatment with metacycline proved more successful ($P < 0.1$) in the patients with complement-fixing antibodies to chlamydia than in those without, as judged from the relief of symptoms. Chlamydia infections may be treated with antibiotics such as

tetracyclines, but recent work suggests that such treatment may not be so effective as supposed.³—We are, etc.,

PER-ANDERS MÅRDH

Institute of Medical Microbiology,
University Hospital,
Lund, Sweden

STIG COLLEEN
BO HOLMQUIST

Department of Urology,
University Hospital,
Lund, Sweden

- ¹ Jones, B. R., *British Journal of Venereal Diseases*, 1972, 48, 13.
- ² Trehanne, J. D., Davey, S. J., Gray, S. J., and Jones, B. R., *British Journal of Venereal Diseases*, 1972, 48, 18.
- ³ Garrod, L. P., and O'Grady, F., in *Antibiotic and Chemotherapy*, p. 448. Edinburgh and London, Churchill-Livingstone, 1972.

Rosette Inhibition Test and Renal Transplantation

SIR,—Despite the detailed report of the method employed by Mr. A. Munro and others (31 July, 1971, p. 271) to assay rosette inhibition, many difficulties may be encountered before reproducible results are obtained. This is to be expected in a test whose mechanism is ill-understood, and a recent report on the influence of physical factors on rosette inhibition¹ serves to emphasize these difficulties. Once reproducible results are obtained, however, in a test that utilizes 10 serial dilutions a shift of two or more dilutions from examination to examination should be regarded as significant.

Our group has found that unimmunosuppressed subjects exhibit a mean inhibitory concentration (M.I.C.) of antilymphocyte globulin between 1:16,000 and 1:32,000. Well suppressed transplant patients have a M.I.C. of about 1:128,000. We note that 24-48 hours before rejection may be diagnosed clinically the patient's M.I.C. sinks to 1:4,000-1:8,000. That is four to five dilution steps difference from "suppressed" to "unsuppressed." We believe that once a patient has been given the immunosuppressive drugs referred to a "normal" level is meaningless. The change in M.I.C. from a level indicating "immunosuppression" to one, two or more dilution steps lower, however, is important.

Our results with the rosette inhibition test seem to confirm the findings of Bewick's group. We are impressed by the early warning of impending rejection that the test appears to give.—We are, etc.,

RICHARD FORREST

ROGER HÄLLGREN

KARL-ERIK FJELLSTRÖM

University Hospital Medical Clinic,
Uppsala, Sweden

- ¹ Morton, H., and Clunie, G. J. A., *Transplantation*, 1972, 14, 211.

Diagnosis of Renal Agenesis

SIR,—Drs. N. C. De and J. R. Harper (16 September, p. 696) draw attention to the problem of resuscitation in infants subse-

quently found to have severe renal anomalies. With improvements in care the perinatal mortality and morbidity directly attributable to obstetric causes has declined and the recognition of infants with external manifestations of severe renal malformations assumes greater significance. We also have encountered instances where intensive resuscitative measures have been applied for considerable periods to such infants who had no hope of survival. Diagnosis is not easy, especially when pressures from labour confuse the characteristic facial appearances in the early neonatal period. Also somewhat similar features can be seen associated with the leaking liquor syndrome.¹ These infants, like renal agenetics, have pulmonary hypoplasia and consequently present problems in resuscitation, but it is important to distinguish them as they may be saved.

Many other factors besides the facial appearance should be considered. Help may be obtained from a careful obstetric history. In infants with severe renal malformations the presence of bilateral talipes and spade-like hands may be of assistance in diagnosis.² Birth weight tends to be low in renal agenesis but not in cases of severe renal cystic dysplasia or of congenital urinary tract obstruction such as urethral atresia or stenosis. Another factor which can prove of great value in diagnosis is macroscopic examination of the placenta, which in most cases of oligohydramnios associated with failure of fetal micturition will reveal the presence of amnion nodosum.^{3,4} Amnion nodosum may be present in association with the leaking liquor syndrome but is much less common in this condition.

Over-emphasis tends to be placed on the facial appearance of infants with severe renal malformations without due consideration of all the other features. Nevertheless, diagnosis will in a few instances present difficulties and resuscitative measures should be undertaken in doubtful cases lest a normal infant affected only by prolonged leakage of liquor be allowed to die.—We are, etc.,

A. D. BAIN

Department of Pathology,
Royal Hospital for Sick Children and
University of Edinburgh

J. S. SCOTT

Department of Obstetrics and Gynaecology,
University of Leeds

- ¹ Bain, A. D., Smith, I. I., and Gauld, I. K., *British Medical Journal*, 1964, 2, 598.
- ² Bain, A. D., and Scott, J. S., *British Medical Journal*, 1960, 1, 841.
- ³ Landing, B. H., *American Journal of Obstetrics and Gynecology*, 1950, 60, 1339.
- ⁴ Scott, J. S., and Bain, A. D., *Proceedings of the Royal Society of Medicine*, 1958, 51, 512.

Immunological Responses in Pregnancy

SIR,—In their paper demonstrating depression of cell-mediated immunity during pregnancy, Dr. Ronald Finn and others (15 July, p. 150) predict that the function of the lymphocytes in the bursa-derived system would not be reduced and might even be increased during pregnancy. They refer to the work of Woodrow and others,³ who found no reduction of humoral antibody production in pregnant rabbits.

We have examined the effect of pregnancy on insulin-binding antibodies present in the serum of insulin-treated patients. In all patients studied insulin antibody levels fell as pregnancy progressed so that by the third

Group	No. of Subjects	Complement-fixing Antibody Titres					
		1/ <5	1/5	1/10	1/20	1/40	1/80
Prostatitis	79	53	3	12	6	4	1
Controls	72	70	—	1	1	—	—