Role of combination antiviral therapy in pandemic influenza

It is impossible to predict which drugs will be effective against a new pandemic strain of influenza. **Sotirios Tsiodras** and colleagues argue that failure to stockpile both major classes of antiviral drugs could prove costly

We currently have two classes of drugs that are effective against influenza viruses: the M2 ion channel inhibitors (amantadine and rimantadine) and the neuraminidase inhibitors (oseltamivir, zanamivir).¹ Although ion channel inhibitors are effective against several subtypes of influenza A viruses,² they are not being widely stockpiled for a future influenza pandemic.³ This is because they cause unacceptable side effects¹ and their use is associated with a rapid emergence of resistance^{1 4 5} without any demonstrable reduction in transmissibility or pathogenicity.⁶ Resistance of influenza A viruses to amantadine is increasing worldwide,⁷ and the US Centers for Disease Control and Prevention recommended against the use of ion channel inhibitors for treatment or prophylaxis during the 2005-6 influenza season.⁸ Policy makers and some medical experts thus consider ion channel inhibitors inappropriate as first line treatment or prophylaxis for pandemic influenza. We believe their role should be reconsidered for three reasons: firstly, the unpredictability of antiviral susceptibility in novel strains of influenza; secondly, on economic grounds; and lastly, for reasons of long term chemical stability.

Why stockpile ion channel inhibitors?

The strongest argument for having ion channel inhibitors available for a pandemic is that we cannot predict the antiviral susceptibility of a novel influenza strain. Ion channel inhibitors show a broad antiviral spectrum against influenza A strains and naturally occurring resistance in new strains is very low.^{9 10} In addition, a recombinant virus possessing the M segment of the 1918 strain was inhibited effectively both in tissue culture and in vivo by amantadine and rimantadine.¹¹ Similarly, influenza strains with pandemic potential occurring later in the 20th century were susceptible to ion channel inhibitors.¹²

Moreover, avian H5N1 strains continue to evolve.¹³ Isolates from human cases in Vietnam, Thailand, and Cambodia differ genotypically from isolates from Indonesia, China, and Eastern European cases¹³ and these differences affect their antiviral susceptibility. The strains from Vietnam, Thailand, and Cambodia have characteristic polymorphisms that confer amantadine resistance¹⁴ whereas isolates from the other countries are mostly susceptible to ion channel inhibitors.¹⁵

Emergence of a pandemic strain resistant to neurami-

Editorial, p268

Sotirios Tsiodras

lecturer in infectious diseases, Fourth Academic Department of Internal Medicine, University of Athens Medical School, Athens 12462, Greece

John D Mooney

public health specialist trainee, South West Public Health Training Programme, Dorset

Angelos Hatzakis

professor, Department of Hygiene and Epidemiology, University of Athens Medical School

Correspondence to: S Tsiodras

nidase inhibitors cannot be excluded, especially as the use of these drugs is expected to increase. Only one mutation is required to lead to full resistance,^{16 17} as is the case for resistance to M2 ion channel inhibitors, and resistance has already occurred in H5N1 strains that have infected humans.¹⁸ Moreover, wide scale prophylaxis with a single drug may lead to resistance becoming commonplace.

Combination therapy

Importantly, the combination of ion channel and neuraminidase inhibitors in vitro reduced the emergence of resistance and may even act synergistically against influenza A viruses.^{19 20} Even low concentrations of oseltamivir prevented the emergence of amantadine resistant variants of the highly pathogenic avian influenza H5N1 virus isolated from Hong Kong in 1997.¹⁹ Combination therapy may also allow the use of a lower dose of ion channel inhibitors, which is known to reduce side effects.²¹

Thus combined use of these drugs may be useful in a future pandemic either for prophylaxis (this needs to be further assessed) or, more importantly, for treatment. The World Health Organization recently recommended combined use of ion channel and neuraminidase inhibitors against the H5N1 subtype.²² However, this recommendation was deemed weak because it was based on low quality evidence (largely because of a scarcity of good trial data), and neuraminidase inhibitors continue to be the preferred treatment when available.²² We urgently need adequately sized trials with combinations of anti-influenza drugs to improve the evidence base.

Costs

Ion channel inhibitors are an older and considerably cheaper class of drug than neuraminidase inhibitors (a sevenfold difference in the September 2006 *British National Formulary*). The additional cost implications of maintaining stocks of both types of antiviral drug are therefore modest. The low cost of ion channel inhibitors, however, could also be a disincentive for drug companies to research their potential role in a flu pandemic. This may partly explain the current lack of evidence and suggests that we may need publicly funded trials into combined regimens. The potential benefits of other therapies alone or in combination with existing antiviral

Ion channel inhibitors could yet have an important role in our armoury against a future flu pandemic

Molecular model of amantadine

drugs also needs exploring. Interventions against the inflammatory cascade that is believed to explain the lethality of H5N1 have been recently highlighted²³ and statins may have a role here.²⁴

In conclusion, ion channel inhibitors could yet have an important role in our armoury against a future flu pandemic. To preserve their activity we recommend that they are not used as monotherapy or for prophylaxis against seasonal or avian influenza. Several countries including the US, the UK, and Greece are already stockpiling ion channel inhibitors. Other countries should consider following suit. As well as being cheap, these drugs are chemically stable, giving them a long shelf life.²⁵ Combined antiviral therapy with neuraminidase inhibitors has the potential to reduce both their side effect profile and the likelihood of resistance. More clinical data are urgently needed to verify such an effect.

Contributors and sources: ST, JDM, and AH have all worked extensively in public health issues relevant to pandemic influenza in their respective countries. This article arose from discussions and planning for pandemic preparedness. ST and AH conceived the idea of the manuscript and ST, JDM, and AH drafted and critically revised the manuscript. ST and AH can serve as guarantors for this article.

Competing interests: None declared.

SUMMARY POINTS

Ion channel inhibitors are not recommended for monotherapy or prophylaxis against seasonal or avian influenza

Combined use with neuraminidase inhibitors may reduce side effects and the risk of resistance

Adequately sized trials of such combinations are urgently needed

Stockpiling of ion channel inhibitors should be considered for potential use in a flu pandemic

Several countries including the US, the UK, and Greece are already stockpiling ion channel inhibitors. Other countries should consider following suit

- 1 Monto AS. The role of antivirals in the control of influenza. *Vaccine* 2003;21:1796-800.
- 2 Dolin R, Reichman RC, Madore HP, Maynard R, Linton PN, Webber-Jones J. A controlled trial of amantadine and rimantadine in the prophylaxis of influenza A infection. *N Engl J Med* 1982;307:580-4.
- 3 World Health Organization. *WHO pandemic influenza draft protocol for rapid response and containment*. 17 March 2006. www.who.int/csr/disease/avian_influenza/guidelines/pandemicfluprotocol_17.03a.pdf.
- 4 Hayden FG, Hay AJ. Emergence and transmission of influenza A viruses resistant to amantadine and rimantadine. *Curr Top Microbiol Immunol* 1992;176:119-30.
- 5 Shiraishi K, Mitamura K, Sakai-Tagawa Y, Goto H, Sugaya N, Kawaoka Y. High frequency of resistant viruses harboring different mutations in amantadine-treated children with influenza. *J Infect Dis* 2003;188:57-61.
- 6 Fleming DM. Managing influenza: amantadine, rimantadine and beyond. *Int J Clin Pract* 2001;55:189-95.
- 7 Bright RA, Medina M, Xu X, Perez-Oroz G, Wallis TR, Davis XM, et al. Incidence of adamantane resistance among influenza A (H3N2) viruses isolated worldwide from 1994 to 2005: a cause for concern. *Lancet* 2005;366:1175-81.
- 8 Centers for Disease Control and Prevention. *CDC recommends against the use of amantadine and rimantadine for the treatment or prophylaxis of influenza in the United States during the 2005-06 influenza season*. www.cdc.gov/flu/han011406.htm.
- 9 Belshe RB, Burk B, Newman F, Cerruti RL, Sim IS. Resistance of influenza A virus to amantadine and rimantadine: results of one decade of surveillance. *J Infect Dis* 1989;159:430-5.
- 10 Ziegler T, Hemphill M, Ziegler ML, Perez-Oroz G, Klimov AI, Hampson AW, et al. Low incidence of rimantadine resistance in field isolates of influenza A viruses. *J Infect Dis* 1999;180:935-9.
- 11 Tumpey TM, Garcia-Sastre A, Mikulasova A, Taubenberger JK, Swayne DE, Palese P, et al. Existing antivirals are effective against influenza viruses with genes from the 1918 pandemic virus. *Proc Natl Acad Sci* 2002;99:13849-54.
- 12 Monto AS, Gunn RA, Bandyk MG, King CL. Prevention of Russian influenza by amantadine. *JAMA* 1979;241:1003-7.
- 13 Garten RJ, Smith C, Xu X, Balish A, Klimov A, Lien P, et al. *Analysis of H5N1 influenza viruses from humans and birds between 2003 and 2005 reveals an increase in the spectrum of viruses with pandemic potential*. Abstract 64. Fifth International Conference on Emerging Infectious Diseases, 2006. www.iceid.org/documents/AbstractsFinal.pdf.
- 14 The World Health Organization Global Influenza Program Surveillance Network. Evolution of H5N1 avian influenza viruses in Asia. *Emerg Infect Dis* 2005;11:1515-21.
- 15 Cheung CL, Rayner JM, Smith GJ, Wang P, Naipospos TS, Zhang J, et al. Distribution of amantadine-resistant H5N1 avian influenza variants in Asia. *J Infect Dis* 2006;193:1626-9.
- 16 Gubareva LV, Bethell R, Hart GJ, Murti KG, Penn CR, Webster RG. Characterization of mutants of influenza A virus selected with the neuraminidase inhibitor 4-guanidino-Neu5Ac2en. *J Virol* 1996;70:1818-27.
- 17 Kiso M, Mitamura K, Sakai-Tagawa Y, Shiraishi K, Kawakami C, Kimura K, et al. Resistant influenza A viruses in children treated with oseltamivir: descriptive study. *Lancet* 2004;364:759-65.
- 18 De Jong MD, Thanh TT, Khanh TH, Vo MH, Smith GJ, Nguyen VC, et al. Oseltamivir resistance during treatment of influenza A (H5N1) infection. *N Engl J Med* 2005;353:2667-72.
- 19 Ilyushina NA, Bovin NV, Webster RG, Govorkova EA. Combination chemotherapy, a potential strategy for reducing the emergence of drug-resistant influenza A variants. *Antiviral Res* 2006;70:121-31.
- 20 Govorkova EA, Fang HB, Tan M, Webster RG. Neuraminidase inhibitor-rimantadine combinations exert additive and synergistic anti-influenza virus effects in MDCK cells. *Antimicrob Agents Chemother* 2004;48:4855-63.
- 21 Reuman PD, Bernstein DI, Keefer MC, Young EC, Sherwood JR, Schiff GM. Efficacy and safety of low dosage amantadine hydrochloride as prophylaxis for influenza A. *Antiviral Res* 1989;11:27-40.
- 22 WHO. *WHO rapid advice guidelines on pharmacological management of humans infected with avian influenza A (H5N1) virus*, 2006. www.who.int/medicines/publications/WHO_PSM_PAR_2006.6.pdf.
- 23 De Jong MD, Simmons CP, Thanh TT, Hien VM, Smith GJD, Chau TNB, et al. Fatal outcome of human influenza A (H5N1) is associated with high viral load and hypercytokinemia. *Nat Med* 2006;12:1203-7.
- 24 Fedson DS. Pandemic influenza: a potential role for statins in treatment and prophylaxis. *Clin Infect Dis* 2006;43:199-205.
- 25 Scholtissek C, Webster RG. Long-term stability of the anti-influenza A compounds—amantadine and rimantadine. *Antiviral Res* 1998;38:213-5.

Accepted: 16 January 2007