Published in final edited form as:
Front Biosci. ; 12: 1450–1462.

Roles of gap junctions and hemichannels in bone cell functions and in signal transmission of mechanical stress

Jean Xin Jiang, Arlene Janel Siller-Jackson, and Sirisha Burra

Department of Biochemistry, University of Texas Health Science Center, 7703 Floyd Curl Drive, San Antonio, TX 78229-3900

Abstract

Gap junctions formed by connexins (Cx) play an important role in transmitting signals between bone cells such as osteoblasts and osteoclasts, cells responsible for bone formation and bone remodeling, respectively. Gap junction intercellular communication (GJIC) has been demonstrated to mediate the process of osteoblast differentiation and bone formation. Furthermore, GJIC propagates Ca^{2+} signaling, conveys anabolic effects of hormones and growth factors, and regulates gene transcription of osteoblast differentiation markers. GJIC is also implicated to regulate osteoclast formation, survival and apoptosis. Compared with other bone cells, the most abundant type are osteocytes, which express large amounts of connexins. Mechanosensing osteocytes connect and form gap junctions with themselves and other cells only through the tips of their dendritic processes, a relatively small percent of the total cell surface area compared to other cells. Recent studies show that in addition to gap junctions, osteoblasts and osteocytes express functional hemichannels, the un-opposed halves of gap junction channels. Hemichannels are localized at the cell surface and function independently of gap junctions. Hemichannels in osteocytes mediate the immediate release of prostaglandins in response to mechanical stress. The major challenges remaining in the field are how the functions of these two types of channels are coordinated in bone cells and what the asserted, distinct effects of these channels are on bone formation and remodeling processes, and on conveying signals elicited by mechanical loading.

Keywords

Bone; Gap Junctions; Hemichannels; Cx43; Connexin; Osteoblast; Osteocyte; Mechanical Stress; Review

2. INTRODUCTION

2.1. Gap Junctions in Bone Cells

Gap junctions are transmembrane channels, which connect the cytoplasm of adjacent cells. These channels permit molecules with molecular weights approximately less than 1 kDa such as small metabolites, ions, and intracellular signaling molecules (i.e. calcium, cAMP, inositol triphosphate) to pass through. Gap junction channels have been demonstrated to be important in modulating cell signaling and tissue function in many organs, such as heart, liver, peripheral nerve, ovary, ear and lens of the eye (1–8). Gap junctions are formed by members of a family of sequentially and structurally related proteins known as connexins. Approximately twenty connexins have been identified and cloned from various tissues and cells (9–11). Six monomers of connexins are joined head-to-head across the extracellular “gap” between two adjacent cells

to form intercellular channels. Connexins are membrane proteins, which consists of four conserved membrane spanning domains and two extracellular loop domains, sharing more than 95% homology. Sequences in the intracellular loop and especially, those in the carboxyl-terminus are divergent between connexins (12).

Three types of connexins, Cx43, Cx45 and Cx46, are expressed in bone tissues. Cx43, a ubiquitously expressed connexin, is identified in virtually all types of bone cells, including cultured osteoblasts from newborn rat calvaria (13), human bone marrow stromal cells, trabecular bone osteoblasts (14), murine osteoblasts (15), primary osteocytes in vivo (16), mandibular bone and periodontal ligament cells of rat teeth (17), osteoblast-like MC3T3-E1 cells (18,19), osteocyte-like MLO-Y4 cells (20,21) and chondrocytes from articular and growth plate cartilage (22–24). Cx45 and Cx46 are expressed in several established osteoblast cell lines. Both Cx43 and Cx46 are expressed in osteoblast-like ROS17/2.8 cells, however, the latter connexin fails to assemble into multimeric complexes and to be expressed on the cell surface to form gap junction channels (25). Cx45 is expressed in UMR-102 and hFOB 1.19 osteoblastic, and MLO-Y4 osteocytic cell lines, but to a much lesser degree than Cx43 (21, 26,27). However, we found that Cx45 protein is expressed in bone marrow, but not in osteoblasts, osteocytes and osteoclasts in the alveolar bone tissues of the tooth (28).

Functional gap junctions in osteoblasts were first demonstrated with electrical conductance and dye injection (29). Voltage-sensitive gap junction currents were detected in osteoblastic cells derived from calvarias of new-born rats with a single gap junction channel conductance of approximate 100 pS (30). Fluorescent dye injected into rat calvarial subperiosteal osteoblasts spreads to neighboring osteoblastic cells via GJIC. Using dye-transfer assays, it has also been observed that dye spreads rapidly between a numbers of odontoblasts (31) and osteoblasts (32). Microinjection of anti-Cx43 antibody in MC3T3-E1 cells blocks cell coupling (33). By using electron microscopy and histocytochemistry, the morphological proof of the existence of gap junction structures has also been obtained for periosteal fibroblasts, osteoblasts, and osteocytes in vivo (32,34–36). Interestingly, in addition to linear gap junctions, stacked, oval, and annular junction structures are found, especially within the osteocyte cytoplasm and in osteocyte cell processes within the canaliculi (36). However, the functional significance of these unusual gap junction structures inside the cells is still obscure. Together, connexins are richly expressed in bone cells and form functional gap junction channels.

2.2. Gap Junctions in Osteoblast Differentiation and Bone Formation

Gap junctions and Cx43 play essential roles in osteoblast differentiation and bone development in human and animal models in vivo (reviewed by Stains and Civitelli (37)). Cx43 mutations have been reported to cause the pleiotropic phenotypes of oculodentodigital dysplasia (ODDD) (38–41) and one of the mutations also leads to the skin disorder palmoplantar keratoderma (41). A mouse model of ODDD was identified from mutagenesis screening, which possesses a Cx43 mutation and displays similar bone defect phenotypes as human ODDD. A recent report shows that ODDD mutants of Cx43 form nonfunctional gap junctions and hemichannels, unapposed halves of gap junction channels (see section 2.6) in C6 glioma cells (42). A mouse knockout model with a genetic deletion of a Cx43 gene has delayed ossification, craniofacial abnormalities, and misshapen ribs (43). Osteoblasts isolated from calvaria of Cx43-null mice display reduced differentiation and mineralization (43). The overall skeletal phenotypes of these knockout mice, however, are not as severe as human ODDD. A possible explanation could be that these analyses had to be conducted at the early developmental stages since these animals died prenatally caused by the blockade of ventricle outflow (4,44). More severe bone phenotypes might be expected if the assay were performed on adult animals. Knockdown of Cx43 using antisense oligonucleotides in chick leads to craniofacial abnormalities (45–47), a phenotype remarkably analogous to ODDD. Moreover, in organ cultures of embryonic chick

mandibular mesenchyme, bone formation is markedly reduced in the presence of the combined antisense oligonucleotides to Cx43 and Cx45, whereas bone formation is still evident when treated with antisense to either connexin (48), implying the compensating effect by these connexins. Two models of osteoblast-specific Cx43 gene deletion driven by either the osteocalcin OG2 promoter or collagen type I promoter were developed (49). These models will provide invaluable tools to investigate the specific role of Cx43 in osteoblasts and in the formation of bone in the adult skeleton. In another model system, recent studies show that mutations in the Cx43 gene cause *sof*, a mutant with a short fin, in zebrafish (50). This study suggests the crucial role of gap junctions in the regulation of bone size and growth.

Gap junctions have been demonstrated to play an important role in osteoblast differentiation and maintenance of a differentiated osteoblast phenotype *in vitro*. Gap junction inhibitors, antisense, and even Cx45 attenuate the stimulatory effect of Cx43 on differentiation of osteoprogenitors or osteoblasts, which is associated with a reduction in the expression of osteoblast differentiation markers, such as alkaline phosphatase, osteocalcin, and collagen type I and decreased mineralization (51–54). Expression of a dominant negative Cx43 construct in MC3T3-E1 cells leads to a decrease in alkaline phosphatase activity and calcium deposition (55). Interestingly, inhibition of GJIC induces the trans-differentiation of osteoblasts to an adipocytic phenotype *in vitro* (56). Conversely, overexpression of Cx43 promotes proliferation and differentiation of osteoblasts (52,57). Gap junction function and Cx43 expression are shown to parallel osteoblast differentiation (27). Promotion of osteoblast differentiation by gap junctions could be interpreted by its role in mediating intercellular passage of signaling factors that are elicited by differentiation-stimulating growth factors or hormones. Indeed, inhibition of gap junctions attenuates the effects of parathyroid hormone on osteoblast differentiation and mineralization (58,58,59). In addition to osteoblasts, GJIC is also involved in the expression of alkaline phosphatase and type I collagen in bone marrow stromal cells when co-cultured with endothelial cells (51). A subsequent study from this group found that alkaline phosphatase activity is only increased by the direct contact of human osteoprogenitor cells with human vascular endothelial cell types, while a transformed human bone marrow endothelial cell line has no such effect (60). Disruption of GJIC attenuates this osteogenic effect. Both studies imply that heterotypic GJIC between two different types of cells is necessary to transmit the signals pertinent to osteoblast differentiation.

The role of gap junctions in osteoblast differentiation could be directly correlated to certain molecules passing through these channels. Cx43 and Cx45 have different gap junctional permeability in osteoblasts (26). Cx43 in osteoblast ROS17/2.8 cells mediated cell-cell coupling for both small ions and larger molecules, but Cx45 in UMR106 cells allowed passage only of small ions. This data implies the distinct roles of gap junction channels formed by these two connexins in passage of ions, signaling molecules and small metabolites in bone cells. Expression of Cx45 alters gap junctional permeability in osteoblastic cells expressing endogenous Cx43 (61). Overexpression of Cx45, which appears to function as a dominant negative, disrupts gap junctions formed by endogenous Cx43. This leads to a decrease in basal activities of the osteocalcin and type I collagen promoters (52). Both Cx43 and Cx45 interact with zonular occludens-1 (ZO-1) in osteoblast cells (62); however, the functional significance of this interaction in osteoblast differentiation is still unclear. Gap junctions have been reported to regulate gene transcription of osteoblastic differentiation markers. Stains and Civitelli (63) report that gap junctions regulate ERK/PI3 kinase (PI3K) signaling, which affects gene transcription through the gap junction-dependent osteocalcin connexin response element. Tbx, a developmental transcription regulatory factor, represses gene transcription of Cx43 both in osteoblasts and possibly in developing embryos (64). Mice deficient in Cx40, a Tbx-5-regulated connexin, display axial and appendicular skeletal malformations with similar phenotypes to Tbx5-deficient mice. Tbx5 exerts its role in bone growth and maturation by controlling gene expression of Cx40 (65).

2.3. Gap Junctions in Osteoclast Formation

In addition to its expression in osteoblast and osteocytes, Cx43 has also been identified in osteoclasts. An osteoclast is a multinucleated cell that degrades and reabsorbs bone. They are involved in the natural turnover of bone tissue along with osteoblasts. Osteoclasts arise from haemopoietic cells of the monocyte/neutrophil lineage. Gap junctions are suggested to play important functional roles in osteoclast precursor fusion to form multinucleated mature osteoclasts, communication in the bone multicellular unit in bone remodeling and osteoclast formation (66–68). Inhibitors of gap junctions, 18 α -glycyrrhetic acid and oleamide, inhibited parathyroid hormone (PTH) and 1,25-(OH) $_2$ D $_3$ (an active form of vitamin D $_3$)-stimulated osteoclastic pit formation, an assay for detection of osteoclast formation (68). The connexin-mimetic peptide GAP 27, that inhibits GJIC, decreases the numbers of TRAP-positive osteoclasts and increases the numbers of apoptotic osteoclasts (69). These experimental evidences suggest that GJIC is crucial for osteoclast formation and survival. Recently, the gap junction inhibitor carbenoxolone was shown to significantly inhibit osteoclastogenesis stimulated by PTH, prostaglandin E $_2$ (PGE $_2$) and 1,25(OH) $_2$ D $_3$ in mouse bone marrow cultures (70), further implicating the role of GJIC in conveying the stimulating signals of these hormones on osteoclast formation and bone remodeling.

2.4. Gap Junction Regulation by Hormones, Growth Factors and Other Components

The functions and expression of gap junctions and Cx43 are regulated by hormones, and other signaling and regulatory molecules. PTH stimulates the expression of Cx43 in osteoblasts and osteocytes (71–74). The stimulatory effect of PTH appears to depend upon the developmental state of the osteoblasts along the osteoblastic differentiation pathway (74). Stable transfection antisense cDNA of Cx43 blocks the increase of cAMP elicited by PTH in ROS17/2.8 cells (59). Additionally, blocking of GJIC attenuates the stimulatory effect of PTH on osteoblast mineralization (58). Treatment with PGE $_2$, in most cases, increases GJIC and connexin expression in osteoblastic cell lines (59,71,72,74–78). Recently, an antiarrhythmic peptide analog rotigaptide (ZP123) has been shown to increase GJIC in human osteoblasts (79). Interestingly, this peptide also completely prevents the loss of femoral trabecular bone strength in ovariectomized rats. 17 β -estradiol and 1,25(OH) $_2$ D $_3$ do not affect Cx43 expression in UMR106 cells (74), but somehow cause cell uncoupling in primary osteocytes (80,81). In contrast to PTH and PGE $_2$, transforming growth factor β , osteogenin and bone morphogenetic protein-2 inhibit GJIC in MC3T3-E1 cells (82). pH plays a critical role in bone formation, by which alkaline pH supports mineral deposition while an acidic pH promotes mineral dissolution (83). GJIC is sensitive to pH changes in MC3T3-E1 cells and lower pH causes a reduction in cell coupling and the transcription rate of Cx43 (84,85). Elevated intracellular cAMP that increases GJIC induces an increase in the expression of osteocalcin, a marker for osteoblast differentiation and a decrease in alkaline phosphatase activity (73,86). ERK/PI3K signaling is suggested to functionally influence GJIC (87).

Components of the extracellular matrix (ECM) have direct impact on osteoblast differentiation, GJIC and connexins. ECM components, such as, type 1 collagen, fibronectin, vitronectin, are present during specific stages of bone development and have been shown to have different roles in supporting adhesion of mechanically strained osteoblasts (88). Modified extracellular matrix molecule, sulfated hyaluronan, enhances the expression of Cx43 and N-cadherin, further resulting in remarkable induction of the alkaline phosphatase activity in rat osteoblast cells (89). Hydroxy apatite (HA) microspheres, which enhance the differentiation of osteoblasts, increase GJIC in osteoblasts (90).

2.5. Gap Junctions in Ca $^{2+}$ Signaling and Propagation in Bone Cells

Ca $^{2+}$ signaling evoked by hormones, growth factors and mechanical stimulation in bone cells is thought to be crucial for bone formation and remodeling. Acting through the ERK signaling

pathway in primary calvarial osteoblasts, extracellular Ca^{2+} induces COX-2 transcription and PGE_2 production (91). Gap junction-dependent and independent mechanisms have been proposed to modulate the Ca^{2+} wave propagation in bone cells. In Cx43-expressing ROS17/2.8 cells, calcium waves are propagated by GJIC and influx of extracellular calcium. However, in Cx45-predominantly expressed UMR106 cells, the propagation of Ca^{2+} is mediated through gap junction-independent pathway, by which calcium waves are transmitted by activation of purinergic receptors, causing a release of intracellular calcium stores (92). In cultured primary osteoblasts and in bone marrow stromal cells, elevated intracellular Ca^{2+} reduces intercellular coupling (72,81). Extracellular Ca^{2+} influx through L-type Ca^{2+} channels triggers the surge of intracellular Ca^{2+} , which is transmitted via Cx43 gap junction channels (93). GJIC is also demonstrated to mediate intercellular signaling in isolated chondrocytes (22,23,94) and paired chondrocytes in situ (24). Gap junctions mediate the propagation of intracellular Ca^{2+} waves in mechanically stimulated articular chondrocytes (95) and between chondrocytes and synovial cells (96,97). Recent evidence from long-term culture of human osteoblast-like cells shows that in less differentiated cells, purinergic receptor P2Y-mediated intracellular calcium waves are primarily involved, but as cells differentiate in culture, gap junction-mediated intracellular Ca^{2+} become more prominent (98). Gap junctions provide a direct transmitting pathway for intracellular Ca^{2+} propagation, which results in organizing the cells as a group of cells as a syncytium for the coordinated action of bone formation and remodeling.

2.6. Gap Junction Hemichannels

In addition to being the major components of gap junction channels, connexins have recently been shown to exist and function in the form of un-apposed halves of gap junction channels called hemichannels (99–101). These channels are localized at the cell surface, independent of physical contact with adjacent cells. Hemichannels, like gap junction channels, display relatively low substrate selectivity and permit molecules with molecular weights less than 1 kDa to pass through. However, the function of hemichannels is very different from gap junctions; the former mediate communication between cells and the extracellular matrix, while the latter is involved in the communication between adjacent cells. An atomic force microscopic study reveals the structures of Cx43 hemichannels as randomly distributed individual particles and clusters, showing a lack of preferential orientation in a lipid membrane (102). Extracellular domains of these undocked hemichannels are structurally different from connexons in docked gap junctional plaques. Hemichannels appear to provide a mechanism for ATP and NAD^+ release, which raises intracellular Ca^{2+} levels and promotes Ca^{2+} wave propagation in astrocytes, bone cells, epithelial cells, and retinal cells (103–111). Hemichannels in astrocytes are shown to be involved in the release of the neurotransmitter glutamate (112). Existence of functional hemichannels formed by Cx43 has been reported in neural progenitors, neurons (113,114), astrocytes (115,116), and heart (117). These hemichannels are regulated by voltage, protein kinase C, extracellular Ca^{2+} , and retinoic acid (106,118,119). Electrophysiological studies demonstrate the opening of hemichannels formed by Cx45 when the external Ca^{2+} concentration is reduced (120). The direct interaction of Ca^{2+} with Asp residues is responsible for preventing voltage-gated opening of Cx32 hemichannels. Disruption of the binding site, which is linked to a hereditary peripheral neuropathy, causes complete Ca^{2+} deregulation of hemichannels (118). Hemichannels are reported to regulate cell volume in response to the change in extracellular physiological calcium (121). Metabolic inhibition also induces opening of Cx43 hemichannels in cortical astrocytes and isolated ventricular myocytes (122), and intriguingly, GJIC is reduced (123). Hemichannels are demonstrated to exist in osteoblasts and osteocytes (109,124,125). These hemichannels expressed in bone cells appear to function as essential transducers of anti-apoptotic effects of bisphosphonates (124,125). We reported that hemichannels formed by Cx43 serve as the pathway for the exit of elevated intracellular PGE_2 in osteocytes induced by

fluid flow shear stress (126,127). However, the physiological roles of hemichannels *in vivo* remain largely unknown.

2.7. Gap Junctions and Hemichannels in Mechanosensing Bone Osteocytes

The skeleton adapts to mechanical usage (128) and mechanical loading promotes bone formation and remodeling. In general, most bone cells may be involved in mechanosensing since several types of cells have been shown to be sensitive to mechanical stress (129). However, several arguments have been raised in favor of osteocytes as the mechanosensory cells (130–132). If osteocytes are the cells responsible for bone modeling and remodeling, key issues are how mechanical loading is sensed, how these signals are conveyed to other non-sensing cells and how these signals are translated into chemical signals. The application of force to bone results in several potential stimuli for osteocyte function including hydrostatic pressure and fluid flow shear stress. Over four decades, various theoretical and experimental studies argue that flow of interstitial fluid driven by extravascular pressure as well as by the applied cyclic mechanical loading is likely the stress-related factor that informs bone cells about mechanical loading and stimulates bone formation. *In vivo* experiments have demonstrated the circulation of interstitial fluid in the lacuno-canalicular network by mechanical loading (133–135). Furthermore, bone cells including osteoblasts and osteocytes are more sensitive to fluid flow shear stress than mechanical strain, stretching and compression (136–140). Theoretical models and other experimental evidence also suggest that osteocytes are the most likely mechanosensor cells in the bone and shear stress induced by fluid flow shear stress is the major mechanism of mechanical loading (130–132,141–146). A cable model predicts that the diffusion time for the spread of current along the membrane of the osteocytic processes is about 0.03 seconds, close to the same predicted pore pressure relaxation time for the draining of the bone fluid into the osteonal canal (147). It has been hypothesized that the bending of primary cilia of an osteocyte by extracellular fluid sends signals into cells through gap junctions (148). Osteocytes have been shown to be more sensitive with respect to the release of PGE₂ following both hydrostatic compression and fluid flow treatment than osteoblasts. However, fluid flow shear stress is even more effective than hydrostatic compression in eliciting this response in osteocytes (138). It has been found that mechanical forces applied to bone cause fluid flow through the canaliculi surrounding the osteocyte that is probably responsible for the deformation of the cell membrane (141,149,150). Fluid flow shear stress is likely to stimulate chemical responses, and the signaling molecules generated are likely to be transmitted between cells by gap junction channels connected through the extensive networks of dendritic processes, and through hemichannels between osteocytes and the extracellular matrix. In turn, the signaling cascade activated in this fashion leads to the expression of important regulatory molecules crucial for modulating bone formation and remodeling.

Osteocytes are dispersed throughout the mineralized matrix. The mature osteocyte is described as a stellate or star-shaped cell with a large number of slender, cytoplasmic processes radiating in all directions but perpendicular to the bone surface. These cell processes pass through the bone in thin canals called canaliculi. It has been hypothesized that matrix-embedded osteocytes emit signals that stimulate osteoblast differentiation into osteocytes (32,131,151). The cell processes of osteocytes are connected with each other and the lining cells via gap junctions (34,35,152), thereby allowing direct cell-to-cell coupling. The rapid intracellular passage of ions and signaling molecules mediated by gap junction channels and the extracellular signaling by hemichannels renders them suitable for sensing mechanical strain and later transmitting these mechanical signals into biochemical events. Therefore, gap junctions and hemichannels in osteocytes appear to have essential, distinctive roles in transmitting the signals elicited by mechanical stimulation to other bone cells and the extracellular matrix, and further promote bone formation and remodeling. In osteocyte-like MLO-Y4 cells and primary osteocytes, we

and others have shown that Cx43 is the major connexin expressed (153,154) and expression patterns are similar in these two types of cells (20,154). In addition to Cx43, Cx45 was also detected in MLO-Y4 cells (21). However, only Cx43, but not Cx45, was detected in osteocytes of alveolar bone in the tooth (28). Yellowley and coworkers (155) have shown that MLO-Y4 cells can couple through gap junctions to osteoblast-like MC3T3-E1 cells detected by the passage of calcein between cells and the mechanically induced calcium response.

2.8. Gap Junctions and Hemichannels in Mechanical Signal Transduction of Bone

There are multiple molecular sensory molecules that sense and somehow concert mechanical stimulation into biological signals in bone cells (for recent review, see Rubin et al. (156)). It has been postulated that mechanical usage of bone results in strain-induced signals that are sensed by the mechanostat and GJIC defines the mechanostat set point (157). Gap junctions through the propagation of intracellular signals contribute to mechanotransduction in bone and contribute to the regulation of bone cell differentiation (158). Few studies have been conducted examining the expression and regulation of Cx43 in response to mechanical stimulation using in vivo animal models. By using a rat experimental tooth movement model, Su et al. (17) reported that Cx43 is increased in osteoclasts and periodontal ligament cells in compression zones, and in osteoblasts and osteocytes in tension zones of the periodontal ligament. They also observed high expression of Cx43 mRNA in osteocytes. Our studies using a mouse tooth movement model show the similar stimulation of Cx43 mRNA expression by mechanical loading in osteoblasts and lining cells in bone formation sites and osteoclasts in resorption sites (28). However, there is no discernible increase of Cx43 mRNA in osteocytes, but in contrast, level of Cx43 protein in alveolar osteocytes is highly up-regulated in response to mechanical loading. This discrepancy between their studies and our work could be caused by the different mechanical models used, variation of tooth movement applied, and duration of loading regimes. These in vivo studies suggest that expression of Cx43 in bone is highly responsive to mechanical loading.

Several in vitro studies have been conducted to determine the relationship between mechanical stress and function and expression of gap junctions and connexins in bone cells. Mechanical loading through cyclical stretching enhances the phosphorylation of Cx43 and GJIC in osteoblastic cells, implying the formation of functional gap junction channels (159). Mechanical stimulation by magnetic field exposure inhibits osteoblast growth through a mechanism independent of gap junction coupling, while the alteration in alkaline phosphatase activity appears to be stimulated by the induced electric field in a gap junction-dependent manner (160). Mechanical stress causes the release of PGE₂, which is shown to be dependent upon gap junctions (161,162). A dominant negative mutant of Cx43 diminishes fluid flow-induced release of PGE₂, but not Ca²⁺ responses (161). Consistently, the fluid flow-induced PGE₂ response of ROS17/2.8 cells is gap junction-mediated and independent of intracellular Ca²⁺ (162). Gap junction activities and accumulation of Cx43 protein in osteocytes are altered by mechanical loading (20,21,87). Oscillating fluid flow has been shown to up-regulate GJIC in MLO-Y4 cells by an ERK1/2 MAP kinase-dependent mechanism (87). We have shown that fluid flow stimulates GJIC and increases Cx43 expression in osteocyte-like MLO-Y4 cells (20). Recently, we found that in addition to gap junctions, Cx43 forming hemichannels mediate the biological responses elicited by fluid flow (127). Fluid flow increased surface expression of Cx43 and induced the rapid opening of hemichannels, which in turn mediated the release of PGE₂ in MLO-Y4 cells. All this evidence suggests that signals generated by mechanical stimulation are likely to be transmitted between bone cells through gap junction channels, and between cells and the extracellular matrix through hemichannels. The role of hemichannel opening in response to mechanical stress is postulated in a model diagram (Figure 1).

Prostaglandin increases the skeletal response to mechanical loading (163). Prostaglandin released by bone cells is generally thought to be a skeletal anabolic agent as these agents can increase bone mass in animals (164–166). A recent report shows that fluid flow induced PGE₂ release by MLO-Y4 cells is reduced by the degradation of the glycocalyx on the cell surface, a hypothesized mechanosensor in osteocytes (167). We show that PGE₂ released via Cx43-hemichannels in response to fluid flow functions in an autocrine fashion to activate EP₂ receptor signaling, including increased intracellular cAMP and activated protein kinase A, which in turn, stimulates gap junction function and Cx43 expression (76,77,127). These data suggest that hemichannels are likely to provide an important novel means, distinct from gap junctions, for modulating the immediate biological responses of osteocytes to mechanical stress.

2.9. Gap Junction Hemichannel Regulation

The experimental evidences revealing the regulatory mechanisms of connexin-forming hemichannels have started to emerge. Dephosphorylation of Cx43 due to ATP depletion and activation of Ca²⁺-dependent protein phosphatases are proposed to induce the opening of hemichannels (168). The opening of hemichannels by ATP-depletion supports a role for hemichannels in causing injury in epithelial cells in general and in renal-tubule cells (169, 170). Liposomes in which nonphosphorylated Cx43 is constituted show greater hemichannel permeability than those phosphorylated by mitogen-activated protein kinase (171). Moreover, hemichannels formed by Cx43(S368A), a PKC site mutant, remains preferentially open (172). Oxidation of Cx43 by enhanced generation of reactive oxygen-derived species also promotes the opening of hemichannels (168). Conversely, a free-radical scavenger (Trolox) inhibits opening of Cx43 hemichannels in metabolically inhibited astrocytes (123,168), implying the involvement of redox potential in the opening of hemichannels. A recent study by Retamal et al. (173) shows that metabolic inhibition increases the levels of Cx43 on the cell surface and induces dephosphorylation and nitrosylation of Cx43.

We recently found that the opening of hemichannels is likely to be modulated by the association of Cx43 with $\alpha 5$ integrin (our unpublished result, (174)). Previous evidence suggests the involvement of integrins in regulation of GJIC and Cx43 expression (175–177). Interaction of $\alpha 3\beta 1$ with laminin 5 promotes gap junction coupling (175). Cx43 and Cx26 expression and distribution, formation of gap junction plaques and GJIC are regulated by matrix proteins (177). Previous immunohistochemical studies have shown the co-localization of Cx43 with cell and substrate adhesion molecules during intramembranous bone formation (178). However, the role of integrin and adhesion molecules in regulation of hemichannels has not been elucidated. It is likely that integrins and cell or substrate adhesion molecules serve as molecular tethers that sense mechanical stress and transducer the effects on gap junctions and hemichannels. Integrins are reported to be mechanical sensors on the cell surface (179,180) and have been proposed to be the candidate mechanosensors in bone cells (181). Fluid flow stimulates pathways that are regulated by integrin binding to the extracellular matrix (182). Among various isotypes of integrins, $\alpha 5$ and $\beta 1$ integrins are expressed in virtually all cell types in bone (183–185) and cartilage (186), and are found to induce responses to mechanical stimuli (181). Integrins, $\alpha 2$, $\alpha 5$ and $\beta 1$ induce ERK activation in osteocytic cells (187). The potential association between integrin and connexins in bone cells, particularly, in osteocytes, may provide a novel, effective mechanism in the process of mechanotransduction mediated by gap junctions and hemichannels.

3. CONCLUSION AND FUTURE PERSPECTIVES

In summary, ample experimental evidences support the notion that gap junction and hemichannels provide important signaling pathways in all major bone cells and play crucial

regulatory roles in the various phases of bone modeling and remodeling processes. These channels pass the signals generated by growth factors, hormones, mechanical stress and other physiological factors between cells and their extracellular environments, and convey the anabolic or catabolic effects of these factors on bone cells.

One of important future directions will involve in the elucidation of the specific role of hemichannels, distinct from gap junctions, and their regulatory components in vivo. At present time, this research has been hampered by a lack of specific dominant negative connexin mutants that can specifically inhibit function of hemichannels, but not gap junctions or vice versa. With the identification of these mutants, the distinct roles of hemichannels as well as gap junctions in bone cells are likely to be revealed. Another line of possible future research will be on the investigation of the roles of gap junctions and hemichannels in osteocytes, the most abundant cell type in normal human bone. Little is known about the function of this cell type compared to any other bone cells. As a mechanosensor cell, it is not clear how osteocytes transduce signals generated by mechanical stress via gap junctions and hemichannels, subsequently convert mechanical stress to chemical signals, and convey these signals between osteocytes and the extracellular matrix, which in turn regulate the bone remodeling process. This avenue of research will increase our understanding of the mechanistic roles of osteocytes in sensing mechanical stress, and in bone biology.

Acknowledgements

The work is supported by NIH grant (AR46798) and Welch Foundation Grant (AQ-1507).

References

1. Bennett MVL, Goodenough DA. Gap junctions, electronic coupling, and intercellular communication. *Neurosciences Res Prog Bull* 1978;16:373–486.
2. Bergoffen J, Scherer SS, Wang S, Scott MO, Bone LJ, Paul DL, Chen K, Lensch MW, Chance PF, Fischbeck KH. Connexin mutations in X-linked Charcot-Marie-Tooth disease. *Science* 1993;262:2039–2042. [PubMed: 8266101]
3. Britz-Cunningham SH, Shah MM, Zuppan CW, Fletcher WH. Mutations of the *connexin43* gap-junction gene in patients with heart malformations and defects of laterality. *New Engl J Med* 1995;332:1323–1329. [PubMed: 7715640]
4. Reaume AG, De Sousa PA, Kulkarni S, Langille BL, Zhu D, Davies TC, Juneja SC, Kidder GM, Rossant J. Cardiac malformation in neonatal mice lacking connexin43. *Science* 1995;267:1831–1834. [PubMed: 7892609]
5. Simon AM, Goodenough DA, Li E, Paul DL. Female infertility in mice lacking connexin37. *Nature (Lond)* 1997;385:525–529. [PubMed: 9020357]
6. Kelsell DP, Dunlop J, Stevens HP, Lench NJ, Liang JN, Parry G, Mueller RF, Leigh IM. Connexin 26 mutations in hereditary non-syndromic sensorineural deafness. *Nature* 1997;387:80–83. [PubMed: 9139825]
7. Gong X, Li E, Klier G, Huang Q, Wu Y, Lei H, Kumar NM, Horwitz J, Gilua NB. Disruption of alpha3 connexin gene leads to proteolysis and cataractogenesis in mice. *Cell* 1997;91:833–843. [PubMed: 9413992]
8. Jiang JX, White TW, Goodenough DA. Changes in connexin expression and distribution during chick lens development. *Dev Biol* 1995;168:649–661. [PubMed: 7729595]
9. Eiberger J, Degen J, romualdi A, Deutsch U, Willecke K, Sohl G. Connexin genes in the mouse and human genome. *Cell Commun Adhes* 2001;8:163–165. [PubMed: 12064582]
10. Willecke K, Eiberger J, Degen J, Eckardt D, romualdi A, Guldenagel M, Deutsch U, Sohl G. Structural and functional diversity of connexin genes in the mouse and human genome. *Biol Chem* 2002;383:725–737. [PubMed: 12108537]
11. Saez JC, Berthoud VM, Branes MC, Martinez AD, Beyer EC. Plasma membrane channels formed by connexins: their regulation and functions. *Physiol Rev* 2003;83:1359–1400. [PubMed: 14506308]

12. Goodenough DA, Goliger JA, Paul DL. Connexins, connexons, and intercellular communication. *Annu Rev Biochem* 1996;65:475–502. [PubMed: 8811187]
13. Schirmacher K, Schmitz I, Winterhager E, Traub O, Bremmer F, Jones D, Binbmann D. Characterization of gap junctions between osteoblast-like cells in culture. *Calcif Tissue Int* 1992;51:285–290. [PubMed: 1330238]
14. Civitelli R, Beyer EC, Warlow PM, Roberson AJ, Geist ST, Steinberg TH. Connexin43 mediates direct intercellular communication in human osteoblastic cell networks. *J Clin Invest* 1993;1888–1896. [PubMed: 8387535]
15. Edelson E. Conduits for cell/cell communication. *MOSAIC* 1990;21:48–56.
16. Mason DJ, Hillam R, Skerry TM. Constitutive in vivo mRNA expression by osteocytes of β -actin, osteocalcin, connexin-43, IGF-1, c-fos and c-jun, but not TNF-alpha nor tartrate-resistant acid phosphatase. *J Bone Miner Res* 1996;11:350–357. [PubMed: 8852945]
17. Su M, Borke JL, Donahue HJ, Li Z, Warshawsky NM, Russell CM, Lewis JE. Expression of connexin 43 in rat mandibular bone and periodontal ligament (PDL) cells during experimental tooth movement. *J Dent Res* 1997;76:1357–1366. [PubMed: 9207768]
18. Chiba H, Sawada N, Oyamada M, Kojima T, Nomura S, Ishii S, Mori M. Relationship between the expression of the gap junction protein and osteoblast phenotype in a human osteoblastic cell line during cell proliferation. *Cell Structure & Function* 1993;18:419–426. [PubMed: 8033223]
19. Yamaguchi DT, Ma D, Lee A, Huang J, Gruber HE. Isolation and characterization of gap junctions in the osteoblastic MC3T3-E1 cell line. *J Bone Miner Res* 1994;9:791–803. [PubMed: 8079655]
20. Cheng B, Zhao S, Luo J, Sprague E, Bonewald LF, Jiang JX. Expression of functional gap junctions and regulation by fluid flow shear stress in osteocyte-like MLO-Y4 cells. *J Bone Miner Res* 2001;16:249–259. [PubMed: 11204425]
21. Thi MM, Kojima T, Cowin SC, Weinbaum S, Spray DC. Fluid flow stress remodels expression and function of junctional proteins in cultured bone cells. *Am J Physiol (Cell Physiol)* 2003;284:C389–C403. [PubMed: 12388096]
22. Donahue HJ, Guilak F, Vander Molen MA, McLeod KJ, Rubin CT, Grande DA, Brink PR. Chondrocytes isolated from mature articular cartilage retain the capacity to form functional gap junctions. *J Bone Miner Res* 1995;10:1359–1364. [PubMed: 7502708]
23. Schwab W, Hofer A, Kasper M. Immunohistochemical distribution of connexin 43 in the cartilage of rats and mice. *Histochem J* 1998;30:413–419. [PubMed: 10192540]
24. Chi SS, Rattner JB, Matyas JR. Communication between paired chondrocytes in the superficial zone of articular cartilage. *J Anat* 2004;205:363–370. [PubMed: 15575885]
25. Koval M, Harley JE, Hinck E, Steinberg TH. Connexin46 is retained as monomer in a trans-golgi compartment of osteoblastic cells. *J Cell Biol* 1997;137:847–857. [PubMed: 9151687]
26. Steinberg TH, Civitelli R, Geist ST, Robertson AJ, Hick E, Veenstra RD, Wang HZ, Warlow PM, Westphale EM, Laing JG, et al. Connexin43 and connexin45 form gap junctions with different molecular permeabilities in osteoblastic cells. *EMBO J* 1994;13:744–750. [PubMed: 8112289]
27. Donahue HJ, Li Z, Zhou Z, Yellowley CE. Differentiation of human fetal osteoblastic cells and gap junctional intercellular communication. *Am J Physiol (Cell Physiol)* 2000;278:C315–C322. [PubMed: 10666026]
28. Gluhak-Heinrich J, Gu S, Pavlin D, Jiang JX. Mechanical loading stimulates expression of connexin 43 in alveolar bone cells in the tooth movement model. *Cell Commun Adhes* 2006;13:115–125. [PubMed: 16613785]
29. Jeanson BG, Feagin FF, Mcminn RW, Shoemaker RL, Rehm WS. Cell-to cell communication of osteoblasts. *J Dent Res* 1979;58:1415–1423. [PubMed: 285112]
30. Schirmacher K, Ramanan SV, Cronin K, Peterson E, Brink PR. Voltage sensitivity of gap junction currents in rat osteoblast-like cells. *Biochim Biophys Acta* 1997;1327:89–96. [PubMed: 9247170]
31. Ushiyama J. Gap junctions between odontoblasts revealed by transjunctional flux of fluorescent tracers. *Cell Tissue Res* 1989;258:611–616. [PubMed: 2482135]
32. Palumbo C, Palazzini S, Marotti G. Morphological study of intercellular junctions during osteocyte differentiation. *Bone* 1990;11:401–406. [PubMed: 2078434]
33. Yamguchi DT, Ma D, Lee A, Huang J, Gruber HE. Isolation and characterization of gap junctions in the osteoblastic MC3T3-E1 cell line. *J Bone Miner Res* 1994;9:791–803. [PubMed: 8079655]

34. Doty SB. Morphological evidence of gap junctions between bone cells. *Calcif Tissue Int* 1981;33:509–512. [PubMed: 6797704]
35. Jones SJ, Gray C, Sakamaki H, Arora M, Boyde A, Gourdie R, Green C. The incidence and size of gap junctions between the bone cells in rat calvaria. *Anat Embryol* 1993;187:L343–352.
36. Shapiro F. Variable conformation of gap junctions linking bone cells: a transmission electron microscopic study of linear, stacked linear, curvilinear, oval and annular junctions. *Calcif Tissue Int* 1997;61:285–293. [PubMed: 9312198]
37. Stains JP, Civitelli R. Gap junctions in skeletal development and function. *Biochim Biophys Acta* 2005;1719:69–81.
38. Paznekas WA, Boyadjiev SA, Shapiro RE, Daniels O, Wollnik B, Keegan CE, Innis JW, Dinulos MB, Christian C, Hannibal MC, Jabs EW. Connexin 43 (GJA1) mutations cause the pleiotropic phenotypes of oculodentodigital dysplasia. *Am J Hum Genet* 2003;72:408–418. [PubMed: 12457340]
39. Kjaer KW, Hansen L, Eiberg H, Leicht P, Opitz JM, Tommerup N. Novel connexin 43 (GJA1) mutation causes oculo-dento-digital dysplasia with curly hair. *Am J Med Genet* 2004;127A:152–157.
40. Richardson R, Donnai D, Meire F, Dixon MJ. Expression of Gja1 correlates with the phenotype observed in oculodentodigital syndrome/type III syndactyly. *J Med Genet* 2004;41:60–67. [PubMed: 14729836]
41. van Steensel M, Spruijt L, van der Burgt I, Bladergroen RS, Vermeer M, Steijlen PM, van Geel M. A 2-bp deletion in the GJA1 gene is associated with oculo-dento-digital dysplasia with palmoplantar keratoderma. *Am J Med Genet* 2005;132:171–174.
42. Lai A, Le DN, Paznekas WA, Gifford WD, Jabs EW, Charles AC. Oculodentodigital dysplasia connexin43 mutations results in non-functional connexin hemichannels and gap junctions in C6 glioma cells. *J Cell Sci* 2006;119:532–541. [PubMed: 16418219]
43. Lecanda F, Warlow PM, Sheikh S, Furlan F, Steinberg TH, Civitelli R. Connexin43 deficiency causes delayed ossification, craniofacial abnormalities, and osteoblast dysfunction. *J Cell Biol* 2000;151:931–943. [PubMed: 11076975]
44. Ya J, Erdtsieck-Ernste EB, de Boer PA, van Kempen MJ, Jongsma H, Gros D, Moorman AF, Lamers WH. Heart defects in connexin43-deficient mice. *Circ Res* 1998;82:360–366. [PubMed: 9486664]
45. Becker DL, McGonnell I, Makarenkova HP, Patel K, Tickle C, Lorimer J, Green CR. Roles for a connexin in morphogenesis of chick embryos revealed using a novel antisense approach. *Dev Gene* 1999;24:33–42.
46. McGonnell IM, Green CR, Tickle C, Becker DL. Connexin43 gap junction protein plays an essential role in morphogenesis of the embryonic chick face. *Dev Dyn* 2001;222:420–438. [PubMed: 11747077]
47. Law LY, Lin JS, Becker CR, Green CR. Knockdown of connexin43-mediated regulation of the zone of polarizing activity in the developing chick limb leads to digit truncation. *Dev Growth Differ* 2002;44:537–547. [PubMed: 12492512]
48. Minkoff R, Bales ES, Kerr CA, Struss WE. Antisense oligonucleotide blockade of connexin expression during embryonic bone formation: evidence of functional compensation within a multigene family. *Dev Gene* 1999;24:43–56.
49. Castro CH, Stains JP, Sheikh S, Szejnfeld VL, Willecke K, Theis M, Civitelli R. Development of mice with osteoblast-specific connexin43 gene deletion. *Cell Commun Adhes* 2003;10:445–450. [PubMed: 14681055]
50. Iovine MK, Higgins EP, Hindes A, Coblitz B, Johnson SL. Mutations in connexin43 (GJA1) perturb bone growth in zebrafish fins. *Dev Biol* 2005;278:208–219. [PubMed: 15649473]
51. Villars F, Guillotin B, Amédée T, Dutoya S, Bordenave L, Bareille R, Amédée J. Effect of HUVEC on human osteoprogenitor cell differentiation needs heterotypic gap junction communication. *Am J Physiol* 2002;282:C775–C785.
52. Lecanda F, Towler DA, Ziambaras K, Cheng SL, Koval M, Steinberg TH, Civitelli R. Gap junctional communication modulates gene expression in osteoblastic cells. *Mol Biol Cell* 1998;9:2249–2258.
53. Li Z, Zhou Z, Yellowley CE, Donahue HJ. Inhibiting gap junctional intercellular communication alters expression of differentiation markers in osteoblastic cells. *Bone* 1999;25:661–666. [PubMed: 10593410]

54. Schiller PC, D'Ippolito G, Balkan W, Roos BA, Howard GA. Gap-junctional communication is required for the maturation process of osteoblastic cells in culture. *Bone* 2001;28:362–369. [PubMed: 11336916]
55. Upham BL, Suzuki J, Chen G, Wang Y, McCabe LR, Chang CC, Krutovskikh VA, Yamasaki H, Trosko JE. Reduced gap junctional intercellular communication and altered biological effects in mouse osteoblast and rat liver oval cell lines transfected with dominant-negative connexin 43. *Mol Carcinog* 2003;37:192–201. [PubMed: 12891628]
56. Schiller PC, D'Ippolito G, Brambilla R, Roos BA, Howard GA. Inhibition of gap-junctional communication induces the trans-differentiation of osteoblasts to an adipocytic phenotype *in vitro*. *J Biol Chem* 2001;276:14133–14138. [PubMed: 11278824]
57. Gramsch B, Gabriel HD, Wiemann M, Grummer R, Winterhager E, Bingmann D, Schirmacher K. Enhancement of connexin 43 expression increases proliferation and differentiation of an osteoblast-like cell line. *Exp Cell Res* 2001;264:397–407. [PubMed: 11262196]
58. Schiller PC, D'Ippolito G, Balkan W, Roos BA, Howard GA. Gap-junctional communication mediates parathyroid hormone stimulation of mineralization in osteoblastic cultures. *Bone* 2001;28:38–44. [PubMed: 11165941]
59. Van der Molen MA, Rubin CT, Mcleod KJ, Mccauley LK, Donahue HJ. Gap junctional intercellular communication contributes to hormonal responsiveness in osteoblastic networks. *J Biol Chem* 1996;271:12165–12171. [PubMed: 8647809]
60. Guillotin B, Bourget C, Remy-Zolgadri M, Bareille R, Fernandez P, Conrad V, Amedee-Vilamitjana J. Human primary endothelial cells stimulate human osteoprogenitor cell differentiation. *Cell Physiol Biochem* 2004;14:325–332. [PubMed: 15319536]
61. Koval M, Geist ST, Westphale EM, Kemendy AE, Civitelli R, Beyer EC, Steinberg TH. Transfected connexin45 alters gap junction permeability in cells expressed endogenous connexin43. *J Cell Biol* 1995;130:987–1005. [PubMed: 7642714]
62. Laing JG, Manley-Markowski RN, Koval M, Civitelli R, Steinberg TH. Connexin45 interacts with zonula occludens-1 and connexin43 in osteoblastic cells. *J Biol Chem* 2001;276:23051–23055. [PubMed: 11313345]
63. Stains JP, Civitelli R. Gap junctions regulate extracellular signal-regulated kinase (ERK) signaling to affect gene transcription. *Mol Biol Cell* 2005;16:64–72. [PubMed: 15525679]
64. Chen JR, Chatterjee B, Meyer R, Yu JC, Borke JL, Isales CM, Kirby ML, Lo CW, Bollag RJ. Tbx2 represses expression of connexin43 in osteoblastic-like cells. *Calcif Tissue Int.* 2004
65. Pizard A, Burgon PG, Paul DL, Bruneau BG, Seidman CE, Seidman JG. Connexin 40, a target of transcription factor Tbx5, patterns wrist, digits, and sternum. *Mol Cell Biol* 2005;25:5073–5083. [PubMed: 15923624]
66. Ilvesaro J, Väänänen K, Tuukkanen J. Bone-resorbing osteoclasts contain gap junctional connexin-43. *J Bone Miner Res* 2000;15:919–926. [PubMed: 10804022]
67. Ilvesaro J, Tuukkanen J. Gap-junctional regulation of osteoclast function. *Crit Rev Eukaryot Gene Expr* 2003;13:133–146. [PubMed: 14696962]
68. Ransjo M, Sahli J, Lie A. Expression of connexin43 mRNA in microisolated murine osteoclasts and regulation of bone resorption *in vitro* by gap junction inhibitors. *Biochem Biophys Res Commun* 2003;303:1179–1185. [PubMed: 12684060]
69. Ilvesaro J, Tavi P, Tuukkanen J. Connexin-mimetic peptide Gap 27 decreases osteoclastic activity. *BMC Musculoskelet Disord* 2001;2:10. [PubMed: 11747476]
70. Matemba SF, Lie A, Ransjo M. Regulation of osteoclastogenesis by gap junction communication. *J Cell Biochem.* 2006in press
71. Civitelli R, Ziambaras K, Warlow PM, Lecanda F, Nelson T, Harley J, Atal N, Beyer EC, Steinberg TH. Regulation of connexin43 expression and function by prostaglandin E2 (PGE2) and parathyroid hormone (PTH) in osteoblastic cells. *J Cell Biochem* 1998;68:8–21. [PubMed: 9407310]
72. Massas R, Benayahu D. Parathyroid hormone effect on cell-to-cell communication in stromal and osteoblastic cells. *J Cell Biochem* 1998;69:81–86. [PubMed: 9513049]
73. Schiller PC, Mehta PP, Roos BA, Howard GA. Hormonal regulation of intercellular communication: parathyroid hormone increases connexin 43 gene expression and gap-junctional communication in osteoblastic cells. *Molecular Endocrinology* 1992;6:1433–1440. [PubMed: 1331776]

74. Schiller PC, Roos BA, Howard GA. Parathyroid hormone up-regulation of connexin 43 gene expression in osteoblasts depends on cell phenotype. *J Bone Miner Res* 1997;12:2005–2013. [PubMed: 9421233]
75. Shen V, Rifas L, Kohler G, Peck WA. Prostaglandins change cell shape and increase intercellular gap junctions in osteoblasts cultured from rat fetal calvaria. *J Bone Miner Res* 1986;1:243–249. [PubMed: 3503541]
76. Cheng B, Kato Y, Zhao S, Luo J, Sprague E, Bonewald LF, Jiang JX. Prostaglandin E₂ is essential for gap junction-mediated intercellular communication between osteocyte-like MLO-Y4 cells in response to mechanical strain. *Endocrinology* 2001;142:3464–3473. [PubMed: 11459792]
77. Cherian PP, Cheng B, Gu S, Sprague E, Bonewald LF, Jiang JX. Effects of mechanical strain on the function of gap junctions in osteocytes are mediated through the prostaglandin EP₂ receptor. *J Biol Chem* 2003;278:43146–43156. [PubMed: 12939279]
78. Donahue HJ, McLeod KJ, Rubin CT, Andersen J, Grine EA, Hertzberg EL, Brink PR. Cell-to-cell communication in osteoblastic networks: cell-line-dependent hormonal regulation of gap junction function. *J Bone Miner Res* 1995;10:881–889. [PubMed: 7572312]
79. Jorgensen NR, Teilmann SC, Henriksen Z, Meier E, Hansen SS, Jensen JE, Sorensen OH, Petersen JS. The antiarrhythmic peptide analog rotigaptide (ZP123) stimulates gap junction intercellular communication in human osteoblasts and prevents decrease in femoral trabecular bone strength in ovariectomized rats. *Endocrinology* 2005;146:4745–4754. [PubMed: 16109789]
80. Massas R, Korenstein R, Benayahu D. Estrogen modulation of osteoblastic cell-to-cell communication. *J Cell Biochem* 1998;69:282–290. [PubMed: 9581867]
81. Schirmacher K, Nonhoff D, Wiemann M, Peterson-Grine E, Brink PR, Bingmann D. Effects of calcium on gap junctions between osteoblast-like cells in culture. *Calcif Tissue Int* 1996;59:259–264. [PubMed: 8781049]
82. Rudkin GH, Yamaguchi DT, Ishida K, Peterson WJ, Bahadosingh F, Thye D, Miller TA. Transforming growth factor-beta, osteogenin, and bone morphogenetic protein-2 inhibit intercellular communication and alter cell proliferation in MC3T3-E1 cells. *J Cell Physiol* 1996;168:433–441. [PubMed: 8707879]
83. Kreiger NS, Sessler NE, Bushinsky DA. Acidosis inhibits osteoblastic and stimulates osteoblastic activity. *Am J Physiol* 1992;262:F442–F448. [PubMed: 1558161]
84. Yamaguchi DT, Huang JT, Ma DF. Regulation of gap junction intercellular communication by pH in MC3T3-E1 osteoblastic cells. *J Bone Min Res* 1995;10:1891–1899.
85. Yamaguchi DT, Mia D. Mechanism of pH regulation of connexin 43 expression in MC3T3-E1 cells. *Biochem Biophys Res Comm* 2003;304:736–739.
86. Romanello M, Moro L, Pirulli D, Crovella S, D'Andrea P. Effects of cAMP on intercellular coupling and osteoblast differentiation. *Biochem Biophys Res Commun* 2001;282:1138–1144.
87. Alford AI, Jacobs CR, Donahue HJ. Oscillating fluid flow regulates gap junction communication in osteocytic MLO-Y4 cells by an ERK1/2 MAP kinase-dependent mechanism. *Bone* 2003;33:64–70. [PubMed: 12919700]
88. Lacouture ME, Schaffer JL, Klickstein LB. A comparison of type I collagen, fibronectin, and vitronectin in supporting adhesion of mechanically strained osteoblasts. *J Bone Miner Res* 2002;17:481–492. [PubMed: 11874239]
89. Nagahata M, Tsuchiya T, Ishiguro T, Matsuda N, Nakatsuchi Y, Teramoto A, Hachimori A, Abe K. A novel function of N-cadherin and connexin43: marked enhancement of alkaline phosphatase activity in rat calvarial osteoblast exposed to sulfated hyaluronan. *Biochem Biophys Res Comm* 2004;315:603–611.
90. Nakaoka R, Ahmed S, Tsuchiya T. Hydroxy apatite microspheres enhance gap junctional intercellular communication of human osteoblasts composed of connexin 43 and 45. *J Biomed Mater Res* 2005;74:181–186.
91. Choudhary S, Wadhwa S, Raisz LG, Alander C, Pilbeam CC. Extracellular calcium is a potent inducer of cyclo-oxygenase-2 in murine osteoblasts through an ERK signaling pathway. *J Bone Miner Res* 2003;18:1813–1824. [PubMed: 14584892]
92. McIntire SL, Reimer RJ, Schuske K, Edwards RH, Jorgensen EM. Identification and characterization of the vesicular GABA transporter. *Nature* 1997;389:870–876. [PubMed: 9349821]

93. Jorgensen NR, Teilmann SC, Henriksen Z, Civitelli R, Sorensen OH, Steinberg TH. Activation of L-type calcium channels is required for gap junction-mediated intercellular calcium signaling in osteoblastic cells. *J Biol Chem* 2003;278:4082–4086. [PubMed: 12446698]
94. D'Andrea P, Vittur F. Gap junctions mediate intercellular calcium signalling in cultured articular chondrocytes. *Cell Calcium* 1996;389–397. [PubMed: 8955553]
95. D'Andrea P, Calabrese A, Capozzi I, Grandolfo M, Tonon R, Vittur F. Intercellular Ca²⁺ wave in mechanically stimulated articular chondrocytes. *Biorheology* 2000;37:75–83. [PubMed: 10912180]
96. D'Andrea P, Calabrese A, Grandolfo M. Intercellular calcium signaling between chondrocytes and snovial cells in co-culture. *Biochem J* 1998;329:681–687. [PubMed: 9445399]
97. Grandolfo M, Calabrese A, D'Andrea P. Mechanism of mechanically induced intercellular calcium waves in rabbit articular chondrocytes and in HIG-82 synovial cells. *J Bone Miner Res* 1998;13:443–453. [PubMed: 9525345]
98. Henriksen Z, Hiken JF, Steinberg TH, Jorgensen NR. The predominant mechanism of intercellular calcium wave propagation changes during long-term culture of human osteoblast-like cells. *Cell Calcium* 2006;39:435–444. [PubMed: 16545868]
99. Goodenough DA, Paul DL. Beyond the gap: Functions of unpaired connexon channels. *Nature Rev* 2003;4:1–10.
100. Bennett MVL, Contreras JE, Bukauskas FF, Saez JC. New roles for astrocytes: Gap junction hemichannels have something to communicate. *Trends Neurosci* 2003;26:610–617. [PubMed: 14585601]
101. Ebihara L. New roles for connexons. *News Physiol Sci* 2003;18:100–103. [PubMed: 12750444]
102. Thimm J, Mechler A, Lin H, Rhee S, Lal R. Calcium-dependent open/closed conformations and interfacial energy maps of reconstituted hemichannels. *J Biol Chem* 2005;280:10646–10654. [PubMed: 15615707]
103. Stout CE, Costantin JL, Naus CCG, Charles AC. Intercellular calcium signaling in astrocytes via ATP release through connexin hemichannels. *J Biol Chem* 2002;277:10482–10488. [PubMed: 11790776]
104. Vanoye CG, Vergara LA, Reuss L. Isolated epithelial cells from amphibian urinary bladder express functional gap junction hemichannels. *Am J Physiol (Cell Physiol)* 1999;276:C279–C284.
105. Gomes P, Srinivas SP, Van Driessche W, Vereecke J, Himpens B. ATP release through connexin hemichannels in corneal endothelial cells. *Invest Ophthalmol Vis Sci* 2005;46:1208–1218. [PubMed: 15790881]
106. Zhang DQ, McMahon DG. Gating of retinal horizontal cell hemi gap junction channels by voltage, Ca²⁺, and retinoic acid. *Mol Vis* 2001;7:247–252. [PubMed: 11715013]
107. Jorgensen NR, Henriksen Z, Sørensen OH, Eriksen EF, Civitelli R, Steinberg TH. Intercellular calcium signaling occurs between human osteoblasts and osteoclasts and requires activation of osteoclast P2X₇ receptors. *J Biol Chem* 2002;277:7574–7580. [PubMed: 11756404]
108. Guyot A, Hanrahan JW. ATP release from human airway epithelial cells studied using a capillary cell culture system. *J Physiol* 2002;545:199–206. [PubMed: 12433960]
109. Romanello M, D'Andrea P. Dual mechanism of intercellular communication in HOBIT osteoblastic cells: A role for gap-junctional hemichannels. *J Bone Miner Res* 2001;16:1465–1476. [PubMed: 11499869]
110. Bruzzone S, Guida L, Zocchi E, Franco L, De Flora A. Connexin 43 hemichannels mediate Ca²⁺-regulated transmembrane NAD⁺ fluxes in intact cells. *FASEB J* 2000;15:10–12. [PubMed: 11099492]
111. Kamermans M, Fahrenfort I, Schultz K, Janssen-Bienhold U, Sjoerdsma T, Weiler R. Hemichannel-mediated inhibition in the outer retina. *Science* 2001;292:1178–1180. [PubMed: 11349152]
112. Ye ZC, Wyeh MS, Baltan-Tekkok S, Ransom BR. Functional hemichannels in astrocytes: a novel mechanism of glutamate release. *J Neurosci* 2003;23:3588–3596. [PubMed: 12736329]
113. Hofer A, Dermietzel R. Visualization and functional blocking of gap junction hemichannel (connexons) with antibodies against external loop domains in astrocytes. *Glia* 1998;24:141–154. [PubMed: 9700496]

114. Boucher S, Bennett SAL. Differential connexin expression, gap junction intercellular coupling, and hemichannel formation in NT2/D1 human neural progenitors and terminally differentiated hNT neurons. *J Neuro Res* 2003;72:393–404.
115. Contreras JE, Saez JC, Bukauskas FF, Bennett MVL. Gating and regulation of connexin 43 (Cx43) hemichannels. *Proc Nat Acad Sci (USA)* 2003;100:11388–11393. [PubMed: 13130072]
116. Saez JC, Contreras JE, Bukauskas FF, Retamal MA, Bennett MVL. Gap junction hemichannels in astrocytes of the CNS. *Acta Physiol Scand* 2003;179:9–22. [PubMed: 12940934]
117. John S, Cesario D, Weiss JN. Gap junctional hemichannels in the heart. *Acta Physiol Scand* 2003;179:23–31. [PubMed: 12940935]
118. Gomez-Hernandez JM, de Miguel M, Larrosa B, Gonzalez D, Barrio LC. Molecular basis of calcium regulation in connexin-32 hemichannels. *Proc Nat Acad Sci (USA)* 2003;100:16030–16035. [PubMed: 14663144]
119. Jedamzik B, Marten I, Negezahayo A, Ernst A, Kolb HA. Regulation of lens rCx46-formed hemichannels by activation of protein kinase C, external Ca(2+) and protons. *J Membrane Biol* 2000;173:39–46. [PubMed: 10612690]
120. Valiunas V. Biophysical properties of connexin-45 gap junction hemichannels studied in vertebrate cells. *J Gen Physiol* 2002;119:147–164. [PubMed: 11815665]
121. Quist AP, Rhee SK, Lin H, Lal R. Physiological role of gap-junctional hemichannels: extracellular calcium-dependent isosmotic volume regulation. *J Cell Biol* 2000;148:1063–1074. [PubMed: 10704454]
122. Kondo RP, Wang SY, John SA, Weiss JN, Goldhaber JJ. Metabolic inhibition activates a non-selective current through connexin hemichannels in isolated ventricular myocytes. *J Mol Cell Cardiol* 2000;32:1859–1872. [PubMed: 11013130]
123. Contreras JE, Sánchez HA, Eugenín EA, Speidel D, Theis M, Willecke K, Bukauskas FF, Bennett MVL, Sáez JC. Metabolic inhibition induces opening of unapposed connexin 43 gap junction hemichannels and reduces gap junctional communication in cortical astrocytes in culture. *Proc Natl Acad Sci* 2002;99:495–500. [PubMed: 11756680]
124. Plotkin LI, Bellido T. Bisphosphonate-induced, hemichannel-mediated anti-apoptosis through the Src/ERK pathway: a gap junction-independent action of connexin43. *Cell Commun Adhes* 2001;2001:377–382. [PubMed: 12064622]
125. Plotkin LI, Manolagas SC, Bellido T. Transduction of cell survival signals by connexin-43 hemichannels. *J Biol Chem* 2002;277:8648–8657. [PubMed: 11741942]
126. Jiang JX, Cherian PP. Hemichannels formed by connexin 43 play an important role in the release of prostaglandin E2 by osteocytes in response to mechanical strain. *Cell Commun Adhes* 2003;10:259–264. [PubMed: 14681026]
127. Cherian PP, Siller-Jackson AJ, Gu S, Wang X, Bonewald LF, Sprague E, Jiang JX. Mechanical strain opens connexin 43 hemichannels in osteocytes: a novel mechanism for the release of prostaglandin. *Mol Biol Cell* 2005;16:3100–3106. [PubMed: 15843434]
128. Frost HM. Perspectives: A proposal general model of the "mechanostat". *Anat Rec* 1996;244:139–147. [PubMed: 8808388]
129. Oster, G. Cell shape: determinants, regulation and regulatory role. In: Stein, WD.; Bronner, F., editors. *Cell mobility and tissue morphogenesis*. Academic Press; San Diego: 1989. p. 33–61.
130. Cowin SC, Moss-Salentijn L, Moss ML. Candidates for the mechanosensory system in bone. *J Biomed Eng* 1991;113:191–197.
131. Aarden EM, Burger EH, Nijweide PJ. Function of osteocytes in bone. *J Cell Biochem* 1994;55:287–299. [PubMed: 7962159]
132. Burger EH, Klein-nulend J. Mechanotransduction in bone-role of the lacunocanalicular network. *FASEB J* 1999;13:S101–S112. [PubMed: 10352151]
133. Seliger WG. Tissue fluid movement in compact bone. *Anat Record* 1970;166:247–255. [PubMed: 5414695]
134. Knothe-Tate ML, Knothe UPN. Experimental elucidation of mechanical loading-induced fluid flow and its potential role in bone metabolism and functional adaptation. *Am J Med Sci* 1998;316:189–195. [PubMed: 9749561]

135. Knothe-Tate MI, Knothe U. An ex vivo model to study transport processes and fluid flow in loaded bone. *J Biomechanics* 2000;33:247–254.
136. Owan I, Burr D, Turner CH, Qiu JTY, Onyia JE, Duncan RL. Mechanotransduction in bone: osteoblasts are more responsive to fluid forces than mechanical strain. *Am J Physiol* 1997;42:C810–C815. [PubMed: 9316399]
137. Smalt R, Mitchell FT, Howard RL, Chambers TJ. Induction of NO and prostaglandin E2 in osteoblasts by wall stress but not mechanical strain. *Am J Physiol* 1997;273:E751–E758. [PubMed: 9357805]
138. Klein-nulend J, Van Der Plas A, Semeins CM, Ajubi NE, Frangos JA, Nijweide PJ, Burger EH. Sensitivity of osteocytes to biomechanical stress in vitro. *FASEB J* 1995;9:441–445. [PubMed: 7896017]
139. Reich KM, Gay CV, Frangos JA. Fluid shear stress as a mediator of osteoblast cyclic adenosine monophosphate production. *J Cell Physiol* 1990;143:100–104. [PubMed: 2156870]
140. Jacobs CR, Yellowley CE, Davis BR, Zhou Z, Cimbala JM, Donahue HJ. Differential effect of steady versus oscillating flow on bone cells. *J Biomech* 1998;31:969–976. [PubMed: 9880053]
141. Weinbaum S, Cowin SC, Zeng Y. A model for the excitation of osteocytes by mechanical loading induced bone fluid shear stresses. *J Biomed Eng* 1993;27:339–360.
142. Turner CH, Forwood MR. What role does the osteocyte network play in bone adaptation? *Bone* 1995;16:283–285. [PubMed: 7786630]
143. Mullender MG, Huiskes R. Osteocytes and bone lining cells: which are the best candidates for mechano-sensors in cancellous bone? *Bone* 1997;20:527–532. [PubMed: 9177866]
144. Burger EH, Klein-nulend J, Van Der Plas A, Nijweide PJ. Function of osteocytes in bone - their role in mechanotransduction. *J Nutrition* 1995;125:2020S–2023S. [PubMed: 7602386]
145. Knothe, Tate ML. "Whither flows the fluid in bone?" An osteocyte's perspective. *J Biomech* 2003;36:1409–1424. [PubMed: 14499290]
146. Wang L, Fritton SP, Weinbaum S, Cowin SC. On bone adaptation due to venous stasis. *J Biomech* 2003;36:1439–1451. [PubMed: 14499293]
147. Zhang D, Cowin SC, Weinbaum S. Electrical signal transmission and gap junction regulation in a bone cell network: a cable model for an osteon. *Annal Biomed Engin* 1997;25:357–374.
148. Whitfield JF. Primary cilium - is it an osteocyte's strain-sensing flowmeter? *J Cell Biochem* 2003;89:233–237. [PubMed: 12704786]
149. Ajubi NE, Klein-nulend J, Nijweide PJ, Vrijheid-lammers T, Alblas MJ, Burger EH. Pulsating fluid flow increases prostaglandin production by cultured chicken osteocytes-a cytoskeleton-dependent process. *Biochem Biophys Res Comm* 1996;225:62–68.
150. Turner CH, Forwood MR, Otter MW. Mechanotransduction in bone: do bone cells act as sensors of fluid flow? *FASEB J* 1994;8:875–878. [PubMed: 8070637]
151. Nefussi JR, Sautier JM, Nicolas V, Forest N. How osteoblasts become osteocytes: a decreasing matrix forming process. *J Biol Buccale* 1991;19:75–82. [PubMed: 1864862]
152. Knothe Tate ML, Adamson JR, Tami AE, Bauer TW. The osteocytes. *Int J Biochem Cell Biol* 2004;36:1–8. [PubMed: 14592527]
153. Kato Y, Windle JJ, Koop BA, Mundy GR, Bonewald LF. Establishment of an osteocyte-like cell line, MLO-Y4. *J Bone Miner Res* 1997;12:2014–2023. [PubMed: 9421234]
154. Gu G, Nars M, Hentunen TA, Metsikkö &, Väänänen HK. Isolated primary osteocytes express functional gap junctions in vitro. *Cell Tissue Res* 2006;323:263–271. [PubMed: 16175387]
155. Yellowley CE, Li Z, Zhou Z, Jacobs CR, Donahue HJ. Functional gap junctions between osteocytic and osteoblastic cells. *J Bone Miner Res* 2000;15:209–217. [PubMed: 10703922]
156. Rubin J, Rubin C, Jacob CR. Molecular pathway mediating mechanical signaling in bone. *Gene* 2006;367:1–16. [PubMed: 16361069]
157. Donahue HJ. Gap junctional intercellular communication in bone: A cellular basis for the mechanostat point. *Calcif Tissue* 1998;62:85–88.
158. Donahue HJ. Gap junctions and biophysical regulation of bone cell differentiation. *Bone* 2000;26:417–422. [PubMed: 10773579]

159. Ziambaras K, Lecanda F, Steinberg TH, Civitelli R. Cyclic stretch enhances gap junctional communication between osteoblastic cells. *J Bone Miner Res* 1998;13:218–228. [PubMed: 9495514]
160. Van der Molen MA, Donahue HJ, Rubin CT, Mcleod KJ. Osteoblastic networks with deficient coupling: differential effects of magnetic and electric field exposure. *Bone* 2000;27:227–231. [PubMed: 10913915]
161. Saunders MM, You J, Trosko JE, Yamasaki H, Li Z, Donahue HJ, Jacobs CR. Gap junctions and fluid flow response in MC3T3-E1 cells. *Am J Physiol (Cell Physiol)* 2001;281:C1917–C1925. [PubMed: 11698250]
162. Saunders MM, You J, Zhou Z, Li Z, Yellowley CE, Kunze EL, Jacobs CR, Donahue HJ. Fluid flow-induced prostaglandin E2 response of osteoblastic ROS 17/2.8 cells is gap junction-mediated and independent of cytosolic calcium. *Bone* 2003;32:350–356. [PubMed: 12689677]
163. Tang LY, Cullen DM, Yee JA, Jee WSS, Kimmel DB. Prostaglandin E₂ increases the skeletal response to mechanical loading. *J Bone Miner Res* 1997;12:276–282. [PubMed: 9041061]
164. Baylink TM, Mohan S, Fitzsimmons RJ, Baylink DJ. Evaluation of signal transduction mechanisms for the mitogenic effects of prostaglandin E2 in normal human bone cells in vitro. *J Bone Min Res* 1996;11:1413–1418.
165. Keller J, Klammer A, Bak B, Suder P. Effects of local prostaglandin E₂ on fracture callus in rabbit. *Acta Orthop Scand* 1993;64:59–63. [PubMed: 8451949]
166. Jee WSS, Ueno K, Deng YP, Woodbury DM. The effects of prostaglandin E2 in growing rats: increased metaphyseal hard tissue and corticoendosteal bone formation. *Calcif Tissue Int* 1985;37:148–156. [PubMed: 3924371]
167. Reilly GC, Haut TR, Yellowley CE, Donahue HJ, Jacobs CR. Fluid flow induced PGE2 release by bone cells is reduced by glycocalyx degradation whereas calcium signals are not. *Biorheology* 2003;40:591–603. [PubMed: 14610310]
168. Contreras JE, Sanchez HA, Veliz LP, Bukauskas FF, Bennett MV, Saez JC. Role of connexin-based gap junction channels and hemichannels in ischemia-induced cell death in nervous tissue. *Brain Res Brain Res Rev* 2004;2004:290–303. [PubMed: 15572178]
169. Vergara L, Bao X, Cooper M, Bello-Reuss E, Reuss L. Gap junctional hemichannels are activated by ATP depletion in human renal proximal tubule cells. *J Membrane Biol* 2003;196:173–184. [PubMed: 14724743]
170. Vergara L, Bao X, Bello-Reuss E, Reuss L. Do connexin 43 gap junctional hemichannels activate and cause cell damage during ATP depletion of renal-tubule cells? *Acta Physiol Scand* 2003;179:33–38. [PubMed: 12940936]
171. Kim DY, Kam Y, Koo SK, Joe CO. Gating connexin 43 channels reconstituted in lipid vesicles by mitogen-activated protein kinase phosphorylation. *J Biol Chem* 1999;274:5581–5587. [PubMed: 10026174]
172. Bao X, Reuss L, Altenberg GA. Regulation of purified and reconstituted connexin 43 hemichannels by protein kinase C-mediated phosphorylation of serine 368. *J Biol Chem* 2004;279:20058–20066. [PubMed: 14973142]
173. Retamal MA, Cortés CJ, Reuss L, Bennett MV, Sáez J. S-nitrosylation and permeation through connexin 43 hemichannels in astrocytes: induction by oxidant stress and reversal by reducing agents. *Proc Nat Acad Sci (USA)* 2006;103:4475–4480. [PubMed: 16537412]
174. Cherian PP, Siller-Jackson AJ, Burra S, Gu S, Bonewald LF, Sprague E, Jiang JX. Potential role of 5 integrin as a mechanosensor in the opening of hemichannels for the release of prostaglandin in response to mechanical stress. *J Bone Min Res* 2005;20 (suppl 1):S25.
175. Lampe PD, Nguyen BP, Gil S, Usui M, Olerud J, Takada Y, Carter WG. Cellular interaction of integrin alpha3beta1 with laminin 5 promotes gap junction communication. *J Cell Biol* 1998;143:1735–1747. [PubMed: 9852164]
176. Guo Y, Martinez-Williams C, Rannels E. Gap junction-microtubule associations in rat alveolar epithelial cells. *Am J Physiol Lung Cell Mol Physiol* 2003;285:L1213–L1221. [PubMed: 14604851]

177. Guo Y, Martinez-Williams C, Yellowley CE, Donahue HJ, Rannels DE. Connexin expression by alveolar epithelial cells is regulated by extracellular matrix. *Am J Physiol Lung Cell Mol Physiol* 2001;280:L191–L202. [PubMed: 11158997]
178. Rundus VR, Marshall GB, Parker SB, Bales ES, Hertzberg EL, Minkoff R. Association of cell and substrate adhesion molecules with connexin43 during intramembranous bone formation. *Histochem J* 1998;30:879–896. [PubMed: 10100730]
179. Wang N, Ingber DE. Control of cytoskeletal mechanisms by extracellular matrix, cell shape and mechanical tension. *Science* 1993;66:2181–2189.
180. Chen CS, Tan J, Tien J. Mechanotransduction at cell-matrix and cell-cell contacts. *Ann Rev Biomed Eng* 2004;6:275–302. [PubMed: 15255771]
181. Salter DM, Robb JE, Wright MO. Electrophysiological responses of human bone cells to mechanical stimulation: evidence for specific integrin function in mechanotransduction. *J Bone Miner Res* 1997;12:1133–1141. [PubMed: 9200014]
182. Katsum A, Orr AW, Tzima E, Schwartz MA. Integrin in mechanotransduction. *J Biol Chem* 2004;279:12001–12004. [PubMed: 14960578]
183. Hughes DE, Salter DM, Dedhar S, Simpson R. Integrin expression in human bone. *J Bone Min Res* 1993;8:527–533.
184. Bennett JH, Carter DH, Alavi AL, Beresford JN, Walsh S. Patterns of integrin expression in a human mandibular explant model of osteoblast differentiation. *Arch Oral Biol* 2001;46:229–238. [PubMed: 11165569]
185. McCarthy AD, Uemura T, Etcheverry SB, Cortizo AM. Advance glycation endproducts interfere with integrin-mediated osteoblastic attachment to a type-1 collagen matrix. *Int J Biochem Cell Biol* 2004;36:840–848. [PubMed: 15006636]
186. Ostergaard K, Salter DM, Petersen J, Bendtzen K, Hvolris J, Andersen CB. Expression of α and β subunits of the integrin superfamily in articular cartilage from macroscopically normal and osteoarthritic human femoral heads. *Am Rheum Dis* 1998;57:303–308.
187. Plotkin LI, Mathov I, Aguirre JI, Parfitt AM, Manolagas SC, Bellido T. Mechanical stimulation prevents osteocyte apoptosis: requirement of integrins, src kinase and ERKs. *Am J Physiol (Cell Physiol)* 2005;289:633–643.

Figure 1.

Model diagram for the role of hemichannels under fluid flow shear stress in osteocytes. Hemichannels are expressed on the plasma membrane away from cell-cell junction regions. (A) In the absence of mechanical stress, hemichannels remain closed, whereas gap junctions are kept open. (B) Fluid flow shear stress induces the opening of hemichannels (upper panel). PGE₂, possibly ATP and other responding physiological factors are released into canaliculi to mediate biological responses elicited by mechanical stress (lower panel).