MEDICAL PRACTICE

Clinical Problems

Parkinsonism: Natural History and Clinical Features

C. A. PALLIS

British Medical Journal, 1971, 3, 683-690**Introduction**

Parkinsonism is a syndrome, (that is, a constellation of symptoms and signs) characterized by tremor, rigidity, and hypokinesia. The condition may result from several causes, yet no cause is known for the commonest form. This variety of the disorder (widely known in the literature as paralysis agitans or Parkinson's disease) will be referred to throughout these articles as idiopathic Parkinsonism.

Genetically-determined Parkinsonism is now receiving increasing attention.¹⁻⁴ A recent monograph⁵ suggested that there was probably a dominant inheritance with clinical manifestations in 25% of gene carriers.

Forty years ago Parkinsonism of known aetiology was usually a sequel to encephalitis lethargica. New cases of this condition are now very rare and Parkinsonism of known aetiology is today in most instances drug-induced. The administration of phenothiazines⁶ (such as chlorpromazine, perphenazine, and promazine), butyrophenones⁷ (such as haloperidol), rauwolfia alkaloids⁸ (such as reserpine), tetrabenazine,⁹ methyl-dopa,^{10, 11} and procaine¹² have all caused Parkinsonism.

Parkinsonism has also been reported in chronic manganese intoxication^{13, 14} after carbon monoxide poisoning,¹⁵ as a sequel to repeated head injury¹⁶ (that is, as part of a chronic traumatic encephalopathy), as a manifestation of neurosyphilis,¹⁷ or of Behçet's disease involving the central nervous system,^{18, 19} in patients with cerebral tumours (whether these directly affected the basal ganglia²⁰ or merely resulted in mid-brain compression²¹), and in a patient with a syrinx involving the substantia nigra.²² Cerebral atherosclerosis was once

considered a fairly common cause of Parkinsonism,²³ and is still so considered by some,²⁴⁻²⁶ though many neurologists would today consider the association of Parkinsonism and cerebral arteriosclerosis fortuitous.

Several rare neurological conditions may show some or all of the clinical features of Parkinsonism and will be discussed under "differential diagnosis." Among them are Wilson's disease (hepatolenticular degeneration), the "rigid-akinetic" form of Huntington's chorea, the Shy-Drager syndrome, Creutzfeldt-Jakob disease, progressive supranuclear palsy, certain varieties of olivopontocerebellar degeneration, and progressive pallidal atrophy.

Historical

In 1817 James Parkinson²⁷ described six patients afflicted by "the shaking palsy" (paralysis agitans). In a classic essay he recorded many of the features of the condition which subsequently came to bear his name. He emphasized the tremor at rest, noted how it might temporarily subside during the performance of willed movement, described the flexed posture, festinant gait, dysphagia and dysarthria, and commented on the severe constipation that might complicate the disease. He did not mention the facial immobility and omitted specific reference either to the muscular rigidity or to the slowness in the initiation and execution of movement—features which many today would consider essential to a diagnosis of Parkinsonism.

Charcot, who did so much to differentiate conditions characterized by tremor, considered the name "paralysis agitans" a misnomer. He pointed out that it "was odd to give the name 'paralysis' to a condition in which for a long time muscular strength was well preserved"²⁸ and stated that tremor might be lacking, even in well-established cases, thus making the adjective "agitans" equally misleading. He proposed that the condition should be called Parkinson's disease.

That infection of the nervous system might cause the syndrome of tremor, rigidity, and hypokinesia does not seem

to have been recorded before the epidemic of encephalitis lethargica which started in 1917. The word "Parkinsonism" was coined by French neurologists at this time. It gained immediate popularity, achieving a permanent place in the vocabulary of neurologists after the description of "syphilitic Parkinsonism" in 1924¹⁷ and of "arteriosclerotic Parkinsonism" in 1929.²³

Epidemiology

Parkinsonism is common. The determination of its exact prevalence, in different parts of the world and at different times is, however, beset with difficulties. Among the problems encountered in epidemiological studies are the insidious nature of the onset of the disease, the fact that the condition is probably underdiagnosed—many laymen and even some medical practitioners considering its cardinal components to be normal features of ageing—the fact that studies which rely on the pooled diagnoses of many clinicians may be grossly in error (through the inclusion of cases of idiopathic tremor²⁹), and the fact that mortality data in this condition are a very poor source of statistics. A recent study of the death certificates of known cases of Parkinsonism³⁰ showed that in almost 40% of cases the crucial information that they had suffered from Parkinsonism was not even mentioned on the certificate.

Several careful studies,^{4 31 32} however, suggest a prevalence rate of just over 1 per 1,000 for the total population of England and Wales³¹ and of 1.57 per 1,000 for the total population of the U.S.A.⁴ With advancing age there is a steep rise of prevalence rates, which in one study approached 1% of patients aged 50 or over.³²

Postencephalitic Parkinsonism is today a condition of dwindling importance. The epidemic of encephalitis lethargica probably affected over a million people. A recent review³³ has shown that many of those who survived this disease soon developed Parkinsonism; at least one-third within three years and one-half or more within five years. Of all the cases of postencephalitic Parkinsonism 80% had developed their symptoms within 10 years of the infection. But, today, if encephalitis lethargica is still occurring, it is only on a very small scale. Moreover, the condition among survivors has tended to become stabilized. Patients now suffering from postencephalitic Parkinsonism are "long-term survivors representing a dwindling fragment of the devastation which encephalitis lethargica left in its wake."³³ An additional implication is that postencephalitic Parkinsonism will soon disappear altogether. The current preponderance of idiopathic cases will become progressively enhanced.

Others have argued a very different thesis.

Basing themselves on the well-documented steady increase in the mean age of all patients with Parkinsonism,^{34 35} they have suggested that all Parkinsonism is postencephalitic, most cases seen today representing a cohort of survivors of the 1917-26 pandemic. On this assumption they have predicted that Parkinsonism itself (and not just its "postencephalitic" form) would disappear as a major clinical entity by 1980. The increase in mean age is not disputed, but an alternative explanation seems more plausible. Before the pandemic of encephalitis lethargica the mean age of onset of Parkinsonism, recorded in several series, varied between 52.7 and 56.8 years.³⁶ During the 1920's it fell by some 15 years owing to the large numbers of young patients who developed Parkinsonism as a sequel to encephalitis lethargica.³⁷ These cases came to comprise two-thirds of all cases of Parkinsonism seen during this period. Between 1930 and 1942 they still comprised half of all cases seen. The epidemic had left behind it sufficient survivors, showing a sufficiently high incidence of postencephalitic Parkinsonism, seriously to affect epidemiological data. When the epidemic subsided, the mean age of survivors began to rise. It is now approaching what it was before 1917. Postencephalitic Parkinsonism, as seen today, is often non-progressive. The mean age of survivors can be expected to rise still further. It may even come

temporarily to exceed the mean age of patients with idiopathic Parkinsonism, which is a progressive disorder. Eventually the patients with post-encephalitic Parkinsonism will all have died. Parkinsonism, far from disappearing, will then probably revert to its classical form.

Natural History

A knowledge of the natural history of Parkinsonism is essential before the long-term benefits of any therapeutic endeavour—medical or surgical—can be realistically assessed. Such knowledge has until recently been scanty and based largely on impressions.

Parkinsonism induced by drugs is usually, but not invariably, reversible. Idiopathic Parkinsonism is a progressive disease. In an authoritative study³⁸ only a small group of atypical cases of idiopathic Parkinsonism were found to maintain their balance and still to be ambulant after 20 years or more. Of the remainder a quarter were severely disabled or dead within 5 years of first being seen, the figures rising to two-thirds in 10 years and 80% in 15 years. There was no correlation between the rate of progression and such factors as the sex of the patient, the age of onset of the disease, or a positive family history. Various clinical signs might progress at independent rates. Parkinsonism with tremor as the initial symptom progressed more slowly than Parkinsonism presenting in other ways.

The same study stressed the good prognosis among surviving cases of postencephalitic Parkinsonism, where only 25% were severely disabled after 20 years of illness. This confirms the previously mentioned impression^{33 36} that the disease of most postencephalitic patients still alive today seems to have reached a plateau beyond which only very slow progression occurs.

Parkinsonism, regardless of aetiology or the age at which it starts, shortens life appreciably. In this study³⁸ the observed mortality was three times that of the general population of the same age, sex, and racial background. The average patient with "primary Parkinsonism" died nine years after the onset of symptoms, though survival periods differed widely, ranging from 1 to 33 years. In postencephalitic Parkinsonism the mean duration of survival was nearly three times as long, partly reflecting the earlier age of onset of the condition. The commonest causes of death, in all age groups, were arteriosclerotic heart disease, bronchopneumonia, malignant neoplasms, vascular lesions of the central nervous system, and infections of the urinary tract. Of these, bronchopneumonia and urinary tract infections may be directly related to the underlying neurological disorder, the other causes of death being those most commonly encountered in non-parkinsonians, in the same age groups.

Clinical Features

TREMOR

Tremor is both one of the main signs and one of the main symptoms of Parkinsonism. In a very large recent series³⁸ it was the initial complaint in 70% of patients with idiopathic Parkinsonism and in just over half the cases with postencephalitic disease.

Parkinsonian tremor usually affects distal muscle groups, such as those of the hand, forearm, and foot. It beats at a rate of 4-8 c/sec, is characteristically present at rest, and is made worse by nervousness, excitement, or fatigue. It is sometimes best seen while the patient is walking. Commonly starting in one hand, it often affects the ipsilateral lower limb before spreading to the opposite side of the body. Flexion and extension movements characteristically involve the metacarpophalangeal joints of the fingers and thumb. Movements of pronation and supination of the forearm may also be seen. The tremor is not constantly present and varies considerably in amplitude from moment to moment. Tremor of the jaw and tongue is not un-

common, but the head is much less often affected. This may help differentiate Parkinsonism from senile tremor: a tremor confined to the head or predominating there is not Parkinsonism.

Parkinson himself commented on how voluntary movement might momentarily bring tremor under control. He described how an afflicted artist "while his hand and arm is palpitating strongly, will seize his pencil and the motions will be suspended, allowing him to use it for a short period." Very occasionally, a genuine intention tremor is present, in addition to the tremor at rest,³⁸ a feature originally pointed out by Gowers.³⁹ Parkinsonian tremor ceases during sleep and Parkinson was wide of the mark when he stated that "even when exhausted nature seizes a small portion of sleep, the motion becomes so violent as not only to shake the bed hangings but even the floor and sashes of the room."²⁷

RIGIDITY

Rigidity is common in all types of Parkinsonism and contributes to many of the more distressing symptoms of the disorder. Like tremor, it may be predominantly unilateral. It appears early in the muscles of the neck.

Rigidity is detected clinically as resistance to passive movement. It is enhanced by active movement of the contralateral limb and this may provide a useful means for detecting it.⁴⁰ Even the smallest muscles may be affected. It differs from spasticity of corticospinal origin in that it affects agonists and antagonists equally, and it is uniform throughout the whole range of passive movement being tested. The terms "plasticity" and "lead-pipe rigidity" have been coined to describe this type of hypertonus. "Cogwheel rigidity"—a jerking sensation often experienced by the examiner on passively stretching a hypertonic Parkinsonian muscle—is still widely attributed to a combination of rigidity and tremor.²⁵ It may occur, however, in the absence of resting tremor, and its frequency (6-12 c/sec) is higher than that of the usual Parkinsonian tremor.⁴¹ It has been suggested that cogwheel rigidity is an overactive physiological tremor, modified by Parkinsonism in some way we do not as yet understand.

James Parkinson²⁷ had commented that patients with the shaking palsy were "less strict than usual in preserving an upright posture." This is not due, however, to a change in moral standards—or to carelessness or inattention—but to rigidity affecting the trunk and nuchal musculature. Lack of postural control may also play a part and will be discussed later. The head is flexed on the trunk and, at a later stage, the trunk is flexed on the pelvis. The back of the patient's head may fail to touch the pillow as he lies on his bed, or remain in the air if the pillow is suddenly withdrawn. The arms are adducted at the shoulders and flexed at the elbows, wrists, and knuckles. The fingers and thumbs are constantly adducted. The thighs are held close together and the knees slightly flexed. In this posture, as Charcot put it, these patients "seem to move all of a piece, as if their joints were soldered."²⁸

The flexed attitude of patients with Parkinsonism probably contributes to the phenomena of anteropulsion and festinant (hurrying) gait. Other contributory factors are the inability of these patients to control their centre of gravity in the sagittal plane.⁴² If pushed from behind, while standing, they will take several increasingly rapid steps, showing—in James Parkinson's words "an almost invincible propensity to run, when wishing only to walk."²⁷ It has been stated that patients with Parkinsonism "run after their centre of gravity" to prevent themselves from falling. This inability to control the centre of gravity in the sagittal plane probably also contributes to festination.⁴² Quick changes of attitude or of direction are difficult.

Parkinsonian hypertonus, if severe and sustained, may lead to deformities and even to contractures.

HYPOKINESIA

So far as the patient is concerned, hypokinesia is probably the most important component of the Parkinsonian triad, though it may be the least obvious to the observer. It is the most difficult symptom to discuss, for little work has been directed at unravelling its nature. We are not really much further advanced today in our understanding of hypokinesia than when Parkinson emphasized how difficult it was for his patients to achieve "a submission of the limbs to the directions of the will . . . in the performance of the most ordinary offices of life."²⁷

Slowness in the initiation and execution of voluntary motor acts and a general poverty of automatic and associated movements are cardinal features of Parkinsonism.

Muscular rigidity may contribute to these findings, but it has been clear for many years that it cannot be the sole explanation. Stereotactic surgery in Parkinsonism may relieve tremor and rigidity, yet leave the basic disturbance of motor function unaltered or worse.

Patients with Parkinsonism often complain of weakness. Their motor deficit is not, however, associated with those characteristic signs of either an upper or a lower motor neuron lesion, which our conceptual categories lead us to expect if weakness is to be considered "organic." Undoubtedly, however, fine movements of the fingers—particularly involving the interossei—are poorly performed by these patients, even in the absence of rigidity or tremor. Similar "weakness" may affect closure of the eyes, protrusion of the tongue, phonation, and articulation.⁴³

Charcot spoke of "a difficulty in performing movement which depended neither on tremor nor on muscular rigidity" and emphasized that there was slowness in performance rather than genuine reduction of motor power. He would admit only to a "diminution apparente de la force musculaire" in Parkinsonism²⁸ and claimed that experiments with dynamometers showed no real weakness. Later investigators,^{43,44} however, have claimed that repetitive movements (such as squeezing a sphygmomanometer bulb) may show rapid enfeeblement. Rapid alternating movements such as opening or closing the hand—or pronating and supinating the forearm—may be grossly defective, often petering out in front of the examiner.⁴² Yet with encouragement, or if there is extreme motivation, Parkinsonian "weakness" may abruptly improve (kinesia paradoxa).

Hypokinesia probably accounts, in part at least, for a host of Parkinsonian symptoms and signs, such as the mask-like facies, reduced blinking, impaired ocular convergence, and monotonous speech amounting in severe cases to virtual anarthria. Phonation may be involved as much as articulation. Hypokinesia almost certainly accounts for the characteristic early clumsiness of fine finger movements and for the micrographia. It is evident in the defective swinging of the arms while walking and the slow and shuffling gait—and at the same time the unusual stillness of a patient with Parkinsonism sitting in a chair, rarely crossing his legs or folding his arms. In these patients "owing to the hypokinesia in the arms, a walking stick is no help but only an additional danger."⁴⁵ It is rather uncommon to see a Parkinsonian carrying one.

In some patients hypokinesia dominates the whole clinical picture. They become house-bound, or even chair-bound, increasingly incapable of looking after themselves, and utterly dependent upon relatives. They have to be dressed, fed, washed, got out of bed, put into bed again. While seated, they are often unable even to shift out of an uncomfortable position.

Less severely affected sufferers may develop paroxysms of akinesia at a stage of the disease when they are still ambulant. They will have to be helped off the pavement or into trains or buses. This symptom of "the feet sticking to the floor" has, however, also been attributed⁴² to an inability to rock the body sufficiently to take the weight on one foot, thereby enabling the other to be moved forward.

MENTAL DISTURBANCES

Irritability and querulousness are features of many chronic

diseases and are common in conditions such as Parkinsonism, where the patient is so critically dependent on relatives or nursing staff. Obsessional thinking has been frequently commented on in patients after encephalitis. Depression is common in all varieties of Parkinsonism.

When a patient with Parkinsonism develops an acute confusional state, the physician's first thought should be whether medication might be responsible. Toxic psychoses are frequent with most anticholinergic drugs, particularly in the elderly, and may take several days to subside.

Chronic mental deterioration in patients with Parkinsonism is more difficult to interpret.

In postencephalitic cases behavioural disturbances and a variable degree of dementia are common. The controversial questions are whether dementia occurs in idiopathic Parkinsonism, and whether there is such an entity as "arteriosclerotic Parkinsonism" (of which dementia would be a feature). Classical teaching still stresses that the mental state is well preserved in idiopathic Parkinsonism. Parkinson²⁷ spoke of the "senses and intellect being uninjured," though Charcot²⁸ had commented on the fact that in the last period of their disease these patients might become "gâteux" (dotards). Mental sluggishness in idiopathic Parkinsonism is admittedly more often apparent than real: the patients understand normally but respond slowly and at times in words so poorly articulated as to be virtually incomprehensible. My impression, however, is that there is more to it than this. A recent survey of a large group of patients with Parkinsonism, of mixed aetiology, showed that a quarter were a burden on their family not as a result of the Parkinsonism itself, which was usually mild, but as a result of dementia.²⁹ Another study from a famous mental hospital has shown that non-senile organic dementia was significantly more frequent in a group of patients with Parkinsonism than in an age and sex-matched group of other patients with mental symptoms—and that patients with idiopathic Parkinsonism showed symptoms of dementia as often as did those with "arteriosclerotic Parkinsonism."³⁰ These findings suggest that dementia may be a feature of idiopathic Parkinsonism, a fact previously hinted at by others.^{44 47}

"Arteriosclerotic Parkinsonism"

The status of "arteriosclerotic Parkinsonism" is controversial. Does such an entity exist? Or is this merely a combination of cerebral atherosclerosis (with its quota of symptoms and signs) and idiopathic Parkinsonism (of somewhat later onset than usual)?

One observer has claimed²⁴ that in a personal series of 90 cases of Parkinsonism over 20% of patients had "arteriosclerotic Parkinsonism." Nevertheless, other observers, analysing a larger series, have found no more clinically detectable arterial disease in patients with Parkinsonism than in age-matched controls.⁴⁸ This latter observation, if correct, throws serious doubt on the role of atherosclerosis in the production of Parkinsonism. The clinical features attributed to "arteriosclerotic Parkinsonism" by different authorities are, moreover, so variable—if not contradictory⁴⁸—as to raise the question of whether this condition, more often defined in clinical rather than in pathological terms, should not be relegated to the realm of neuromythology.

OCULAR ABNORMALITIES

Ocular signs are common in all varieties of Parkinsonism. They range from the triviality of infrequent blinking to the dramatic features of a full oculogyric attack.

Infrequent blinking is a common feature of all types of Parkinsonism. It is a manifestation of hypokinesia. Combined with increased width of the palpebral fissure it gives many Parkinsonian patients a rather staring appearance. A normal person blinks several times a minute. Patients with Parkinsonism may not blink at all during this period.

Closure of the eyes may be poorly maintained, blepharoclonus being commonly observed on asking the patient to keep his eyes

gently closed. This sign is not, of course, specific for Parkinsonism. Blepharospasm and a simple inability to open the eyes once they have been closed are also encountered, particularly in postencephalitic cases.

Glabellar Tap

An interesting and at times diagnostically helpful sign is the positive glabellar tap. Each time the root of the nose is tapped there is blinking, clinically synchronous with the blow (to avoid reflex blinking from visual threat, the tapping finger should be invisible to the patient). In normal persons blinking ceases after the first few taps. In Parkinsonism it often persists indefinitely, provided the rate of tapping is not over 2 per second. The sign is almost always present in Parkinsonism, often at a very early stage of the disease.⁴⁹ It may be encountered in patients with widespread neurological disease (dementia, encephalitis, sequelae of severe head injury, etc.), but under such circumstances difficulties in differential diagnosis from Parkinsonism are unlikely to be encountered.

A byproduct of rigidity and loss of associated movements may be noted when testing horizontal gaze in patients with Parkinsonism. The average person, when asked to follow a finger sideways, will usually move his head together with his eyes, to the mild annoyance of the examiner. Patients with Parkinsonism obey their examiners to perfection.

Paresis of convergence is common in all varieties of Parkinsonism but particularly in postencephalitic cases, where other forms of ophthalmoplegia may also be seen. In most patients there is a normal range of conjugate lateral gaze although the movements may be "saccadic" (jerky). Vertical upward gaze is sometimes limited. Some patients show a peculiar tendency to blink once or twice before looking sideways to order.⁴³ Nystagmus is not uncommon in postencephalitic Parkinsonism.

Pupillary abnormalities, either unilateral or bilateral, may be seen in postencephalitic Parkinsonism. Argyll Robertson pupils may, very rarely, be encountered. More often there is impairment of the near reflex. A paradoxical pupillary response to light (dilatation) may occasionally be seen.

Oculogyric Crises

Oculogyric crises are significantly more common among women than among men.⁵⁰ They do not occur in the idiopathic type of Parkinsonism. They may last from a few minutes to twelve hours or more, and may occur several times a week, constituting a major symptom in certain cases of postencephalitic Parkinsonism.

Crises may be preceded by fluttering of the lids and by a vertical nystagmus with its quick phase directed upwards. During a crisis the eyes are deviated and "fixed" usually in an upward (or upward and outward) posture. When there is a lateral component to the abnormal posture it is usually away from the side of the body showing the greatest rigidity. Forward gaze may be achieved for a few seconds by a great effort of will, but downward gaze is impossible during an attack.

During the crises the face is flushed. Scleral congestion may be appreciable and lachrymation may occur.⁵⁰ The pupillary responses to light are sluggish. Muscular rigidity may be increased and total akinesia may supervene. Vertigo and headache may be associated features. The blood pressure is often raised and tachycardia is common. Fear may subsequently be described. Emotional disturbances usually accompany severe attacks during which patients often make grunting or moaning sounds.

In postencephalitic Parkinsonism oculogyric attacks tend to occur in patients with obvious evidence of brain-stem lesions, in the form of pupillary abnormalities or disturbances of external ocular movement. In one study⁵⁰ three-quarters of the patients

had associated nystagmus, thought to be of vestibular origin in view of the absence of cerebellar signs. Attacks can be precipitated by caloric vestibular stimulation.⁵¹

Oculogyric crises provoked by phenothiazines differ from the postencephalitic type in that they are nearly always associated with lingual movements, spasm of the neck muscles, head rotation and retraction, and even opisthotonus; these may occur but are rare in postencephalitic crises. In susceptible individuals dyskinesias such as oculogyric attacks are among the earliest neurological manifestations of phenothiazine overdose;⁶ they do not occur in a context of established, drug-induced Parkinsonism.

POSTURAL DISORDERS

Defects of postural fixation⁴² of the head, trunk, or limbs may account for many unusual symptoms and signs encountered in patients with postencephalitic Parkinsonism.

The head may fall forward as when a normal person falls asleep while sitting in a chair. Blindfolding the patient may exacerbate this tendency. The forward drop of the head can sometimes best be demonstrated when the patient is on "all fours." Similar "slumping" forwards or backwards of the whole trunk may occur while the patient is sitting on a stool. If this occurs while he is walking it may immobilize him in flexion. Such postural disturbances of the head or trunk may be found when there is only minimal rigidity.

Similar disturbances may be seen in the limbs. Many of these patients have difficulty in maintaining their arms outstretched while using their hands at shoulder level. The arms droop rapidly. Similar defects of postural fixation in the lower limbs may lead to sagging of the knees. Loss of postural fixation of the body as a whole is best shown by difficulty in resisting the effects of being pushed. A loss of righting reflexes in patients with Parkinsonism may lead to gross physical instability, to a loss of ability to stand unaided, and to a definite tendency to fall "all of a piece"—that is, without the usual defensive reactions.

Whether these disturbances are merely further manifestations of hypokinesia or whether they represent a more specific disorder of postural reflexes, as has been argued,⁴² is at present uncertain.

SKELETAL DEFORMITIES

These are not uncommon, particularly in patients with postencephalitic Parkinsonism. They may result in grossly abnormal appearances of the hands and feet, and add to the numerous problems already confronting these unfortunate patients.

Charcot first pointed out (and illustrated) how abnormal hand postures in "paralysis agitans" might mimic to a nicety those seen in certain cases of rheumatoid arthritis. He stressed, however, that there would be found "neither swelling nor stiffness of joints, nor the bony deposits and creaking seen in nodose rheumatism."²⁸

In the commonest form of the deformity the hand is held in slight ulnar deviation with the fingers flexed at the metacarpophalangeal joint and extended at the interphalangeal joints.

The thumb is adducted, opposed, and extended at both metacarpophalangeal and interphalangeal joints. As the condition progresses the extension at the interphalangeal joints progresses. The terminal phalanges finally sublux and may then go into mild flexion. The condition closely resembles the "intrinsic contractures" and "swan neck" digital deformities well known to orthopaedic surgeons and rheumatologists⁵²⁻⁵⁵ and probably arises on a similar basis—namely, a state of sustained contraction of the intrinsic muscles of the hand. But whereas in rheumatoid arthritis this state of contraction is probably due to primary inflammatory changes in the muscles of the hand,^{54 56} in Parkinsonism it is due to abnormal neuronal drive. This can be shown by the fact that the digital deformities of Parkinsonism may in some cases be abolished by an ulnar nerve block⁵⁷ or by contralateral stereotactic thalamic surgery. Fixed contractures may ultimately develop, however. X-ray films of the hands will show no erosions, even in the presence of gross deformity.

The feet may be inverted in an equinovarus posture and exhibit a painful clawing and crowding of the toes, which will contribute to the patient's locomotor difficulties. This again may be improved by intracranial surgery.

CARDIOVASCULAR ASPECTS

Idiopathic Parkinsonism usually starts after the age of 50. It is therefore not surprising that patients with this disease will commonly be found to have arterial disease as well.

It has been claimed that patients with idiopathic Parkinsonism generally have a lower blood pressure than other individuals of their age and sex.⁵⁸ There seems to be little quantitative evidence for this belief at present and the subject warrants further investigation. What has been established is that the baroreceptor reflexes deteriorate with age^{59, 60} and this alone probably explains the undoubtedly impaired adjustments to postural change encountered in Parkinsonism.⁶¹ Some workers have suggested that Parkinsonism itself may result in impaired cardiovascular reflexes.⁶² Reduced aldosterone output has been described.^{63 64}

ALIMENTARY DISORDER

Symptoms related to the alimentary tract are common in Parkinsonism. An awareness of this fact may spare some patients unnecessary, and at times risky, investigations.

Masticatory disorders are not uncommon, particularly in postencephalitic Parkinsonism,⁶⁵ and are usually attributed to a combination of bradykinesia and rigidity. Dyskinesia of the lips and tongue may occasionally interfere with chewing.⁶⁶ Hitherto rare, they are being seen with increasing frequency since the advent of levodopa.

Disorder of Salivation

The picture of the advanced case, almost constantly drooling saliva, is familiar. Some neurologists state or imply that this is due to "sialorrhoea" or salivary hypersecretion;^{25 43} others attribute the disorder to dysphagia and to mechanical difficulty in coping with a normal amount of saliva. The only study in which the amount of saliva produced by patients with Parkinsonism was actually measured suggested that mechanical factors were paramount.⁶⁷ In the words of Parkinson "the saliva fails of being directed to the back part of the fauces and hence is continually draining from the mouth."²⁷

Dysphagia of mild degree is common in all varieties of Parkinsonism and is usually experienced high up in the throat. Some radiological studies^{68 69} have shown delay in the initiation of swallowing, irregular and jerky epiglottic movements, and stasis of the contrast in the pyriform fossa and valleculae. Others have shown no appreciable disturbance in the pharyngeal phase of swallowing, even in patients complaining of dysphagia.⁷⁰ Spill-over into the bronchial tree may certainly occur, such aspiration probably being more common than is generally realized. We have recently seen two patients with Parkinsonism who had developed lipoid pneumonia due to aspiration of liquid paraffin, taken to cope with their severe constipation.

Despite Parkinson's claim, "perpetual agitation" of the muscles of the tongue and pharynx does not significantly contribute to dysphagia, which is in the main related to deglutitional rigidity and bradykinesia. The inferior pharyngeal constrictor is a voluntary muscle and may expect to be involved in Parkinsonian rigidity. The vagal innervation in man may, however, extend as high as the pharyngo-oesophageal junction. Lesions of the dorsal vagal nuclei (which provides parasympathetic innervation to thoracic and abdominal viscera) are very common in Parkinsonism⁷¹ and may account for some of the dysphagia.

Oesophageal mobility may be abnormal. About 5% of unselected Parkinsonian patients show non-peristaltic segmental oesophageal spasm ("cork-screw oesophagus") on barium swallow,⁷² a change attributed to neuromuscular incoordination of the oesophageal wall.

Heartburn and Constipation

Heartburn is common⁶⁵ and at least a quarter of patients with Parkinsonism (whatever the aetiology or severity) have hiatus hernia of the sliding type and gastro-oesophageal reflux.⁷³ The incidence of hiccup and vomiting is not appreciably increased. The prevalence of peptic ulcer in Parkinsonism is controversial.^{65 73}

Constipation is very common in Parkinsonism, particularly in the more severe cases. It is a major symptom of the disease and a cause of great distress to many patients. Parkinson himself stated that "the bowels . . . in most cases demand stimulating medicines of very considerable power, the expulsion of faeces from the rectum sometimes requiring mechanical aid."²⁷

Over half the patients with Parkinsonism fail to defecate once daily,⁶⁵ compared with 12.5% of an age-matched control group. The cause of this symptom is unknown but is usually attributed to the combined effects of a sedentary life, possible salivary loss, and diminished food and fluid intake. Constipation may be severe, even in patients not receiving anticholinergic medication, though such medication tends to aggravate it. The constipation may lead to megacolon,⁷⁴ large bowel obstruction, and unnecessary laparotomies.

Weight Loss

Three-quarters of patients with Parkinsonism are below their optimum weight⁴⁴ and some lose weight rapidly. Charcot²⁸ described a severe generalized muscle wasting in the advanced stages of the disease, but this is now thought to be due to defective calorie intake. Rigidity and tremor of the hands, and problems of chewing and swallowing all contribute to make eating a laborious and time-consuming exercise. This should be remembered before patients with Parkinsonism are submitted to extensive investigations for suspected occult cancer.

OTHER SYMPTOMS AND SIGNS

Many patients with postencephalitic Parkinsonism have a greasy skin and seborrhoea.^{75 76} A greatly increased secretion of hard wax into the external auditory meatus may also occur.⁵⁰

Intolerance to warmth was described as a feature of "Parkinson's disease" by Charcot.²⁸ Attacks of paroxysmal sweating, requiring a change of pyjamas, or of sheets, may occur in patients with postencephalitic Parkinsonism, even in the winter months. During such attacks the face may become flushed and there may be profuse salivation, slight fever, and considerable tachycardia (up to 150/min).⁵⁰

Bladder symptoms are not uncommon in Parkinsonism, but there seem to have been no major studies of the subject using modern cystometrographic techniques. Several older texts speak of "detrusor tetany" in Parkinsonism. Prostatic hypertrophy of mild degree, which would probably have been fairly well tolerated in a normal person, may give rise to serious symptoms in patients with Parkinsonism.⁴⁴ Anticholinergic medication in such patients may precipitate acute retention.

Sensory symptoms are common in all forms of Parkinsonism. Patients may complain of tightness, stiffness, and considerable discomfort in the limbs and experience sudden urges to be moved. These symptoms are probably related to the hypokinesia.

Diagnosis

UNUSUAL PRESENTATIONS

The average case of moderately advanced Parkinsonism presents no diagnostic problems. The flexed posture, expressionless face, lack of blinking, and asymmetrical distal tremor in a context of generally diminished mobility form a clinical constellation which is immediately recognizable. If to these are added dysarthria, nuchal rigidity, cogwheeling of the limbs, and a shuffling or festinant gait the diagnosis is inescapable. In certain circumstances, however, diagnostic difficulties may arise:

(1) In its very early stages the condition is often missed. Mild hypokinesia may be attributed to "just old age" and tremor to neurosis, particularly when it comes on—as it sometimes does—after some severe emotional stress. The earliest evidence of Parkinsonism may be discovered during examination for some completely unrelated condition or following some surgical operation, when problems may arise with postoperative physiotherapy. Retardation from severe depression or hypothyroidism may occasionally cause diagnostic problems.

(2) If the tremor is inconspicuous and the rigidity predominantly one-sided the slowly progressive unilateral disability may lead the unwary to suspect cerebral tumour. Most neurologists will have seen patients with Parkinsonism who have been investigated with brain scans and E.E.G.s (not to mention more unpleasant methods) to exclude intracranial space-occupying lesions.

(3) In elderly patients, with as yet undiagnosed Parkinsonism, vague aches and pains, particularly if felt mainly in the back, shoulders, or hips may lead to misdiagnosis of senile osteoporosis or degenerative arthritis, until the underlying rigidity is detected. Again, this will be most likely when tremor is minimal.

In the diagnosed case the joints may again come under unjustified suspicion. Some patients, severely limited by rigidity and bradykinesia, will experience distressing urges to move, which, of course, they find it difficult or impossible to comply with. Such patients may complain bitterly of discomfort in the affected limbs. The deformities of advanced Parkinsonism have already been referred to and their similarity to those of rheumatoid arthritis stressed. Rheumatoid arthritis and Parkinsonism may, of course, coexist.

(4) In patients with known vascular disease—and in particular in patients who have already had a stroke—the development of Parkinsonism may be missed altogether, and a progressive neurological deficit may be attributed to cerebral arteriosclerosis alone. Dysarthria and dysphagia will be wrongly attributed to "pseudo-bulbar palsy." When such "pseudo-bulbar palsy" is not associated with either a brisk jaw-jerk or signs of emotional release (uncontrolled laughter or weeping) it is always worth bearing in mind the possible co-existence of idiopathic Parkinsonism.

Another confusing situation is where a stroke obliterates the evidence of pre-existing Parkinsonism. A minor degree of tremor or rigidity in the non-paralysed limbs may then provide the only clue. As the paralytic effects of the stroke recede, the diagnostically revealing tremor may reappear on the originally affected side.

(5) Unexpected falls may occasionally be the presenting symptom of Parkinsonism. Although in some cases there may be symptoms and signs of associated vertebrobasilar ischaemia, falling is often solely attributable to the loss of righting reflexes which may be an obvious feature in certain cases.

(6) Gross oedema of the feet may lead to a suspicion of heart failure or even of thrombosis of the inferior vena cava. Such oedema will often clear rapidly when the legs are raised. It is usually the result of prolonged immobility interfering with venous return from the legs, though in some cases hypoproteinaemia may be playing a minor part.

Differential Diagnosis

Aetiological diagnosis may be difficult. Relevant points are the presence or absence of a family history of similar disorder, a history of ingestion of drugs or exposure to toxic hazards, a past history of encephalitis lethargica, the age of the patient at the onset of symptoms; the rate of progression; the exact nature of any involuntary movements seen; and whether the neurological

disorder consists solely of tremor, rigidity, and hypokinesia, or whether these are only part of a wider neurological deficit.

Postencephalitic Parkinsonism can often be distinguished from idiopathic Parkinsonism. There may be a clear-cut history of past encephalitis,* a characteristic onset early in life, characteristic non-progression (or only very slow progression) over two or more decades, and characteristic residua such as tics, behaviour disorder, pallidalia, severe skeletal deformities, appreciable seborrhoea, or oculogyric spasms. In practice, with the progressive decline of postencephalitic Parkinsonism, the opportunities for such differentiation are themselves rapidly declining.

The differential diagnosis of idiopathic Parkinsonism from other neurological disorders is mainly a matter of prognosis, though there are therapeutic implications to a diagnosis of hepatolenticular degeneration. Essential tremor is a common and benign disorder but most of the other conditions discussed carry a very bad prognosis. They are all much rarer than idiopathic Parkinsonism. Though some of these disease may, at some stage or other of their evolution, show features of Parkinsonism there will usually be sufficient clinical clues to lead the physician to recognize that the case is "atypical" and that he is dealing with some other condition.

In practice one will far more often be confronted with the problem of deciding whether a given patient has Parkinsonism or not, than of deciding to which rare nosological entity his Parkinsonism is attributable.

ESSENTIAL TREMOR

Essential tremor is a chronic, benign, monosymptomatic disorder, of unknown aetiology, which progresses very slowly over several decades. The peak incidence is in early adult life and there is often a positive family history. The tremor varies widely in amplitude and frequency, at first being noticeable only on movement²⁵⁻²⁷ but later also appearing at rest. It affects the head much more often than does Parkinsonian tremor, but nearly always spares the legs. There are no other abnormal neurological signs (such as rigidity or hypokinesia) and significant incapacity does not develop. A similar tremor appearing in old age is referred to as "senile tremor." If there is a positive family history, the condition is described as "familial tremor."

HEPATOLENTICULAR DEGENERATION (WILSON'S DISEASE)

When "Parkinsonism" develops during adolescence or in early adult life, doctors should suspect and seek to exclude hepatolenticular degeneration. At least in its early stages, the rigidity in this condition may resemble that of Parkinsonism. The involuntary movements tend to be more varied. Choreic grimacing and jerking, tremors, athetoid and dystonic writhing, and a very coarse "flapping" movement of the outstretched hands may all be seen. Wilson's disease should be considered when involuntary movements, at rest suggestive of Parkinsonism, come to resemble those of multiple sclerosis when the patient is asked to put his finger to his nose. Dysarthria and euphoria are often early features. Mental deterioration is a feature of the more advanced cases.

Useful help in clinching the diagnosis will be provided by such findings as corneal scleral pigmentation (the Kayser-Fleischer ring) and a bluish discoloration of the nail beds (both due to the deposition of copper), combined with clinical and/or laboratory evidence of hepatic disease and abnormalities of serum copper, caeruloplasmin, and urinary amino acids (in both the patient and some of his relatives).

*A past history of a "flu-like" illness is insufficient. It can usually be attributed to flu. A part "flu-like" illness lasting 10 days or more is, however, suggestive.

HUNTINGTON'S CHOREA

Huntington's chorea, with its familial occurrence of dementia, hypotonia, and involuntary movements of choreic type, is usually easy to recognize. In its classical form it is unlikely to be misdiagnosed as Parkinsonism. Rigid and akinetic forms of the disease occur, however,⁷⁸⁻⁷⁹ are probably more common than is generally recognized, and may give rise to considerable diagnostic difficulties.

The disease may manifest itself from the onset in a rigid form and some cases show an akinesia quite out of proportion to the rigidity. Chorea in these cases may not be a prominent feature, and tends to be modified into athetosis or even, ultimately, into involuntary movements resembling torsion-dystonia. Expressionless facies, infrequent blinking, forward tilting of the head and trunk, flexion of the forearms and wrists, loss of normal associated arm movements when walking, and festinant gait have all been described in these "rigid-akinetic" variants of Huntington's disease.

Rigid cases of Huntington's chorea tend to run in families and are often misdiagnosed as hepatolenticular degeneration or Parkinsonism. Errors will occur most frequently when the crucial family history is not available. The most important clues distinguishing this form of Huntington's chorea from Parkinsonism are the early age of the onset of the condition—the mean age of onset in 64 rigid cases was 22.2 years⁷⁹ and the nature of any involuntary movement exhibited. Tremor, which is not uncommon, will be characteristically accentuated on movement, and classical "pill-rolling" or inhibition of the tremor on exertion will not be seen.

SHY-DRAGER SYNDROME

The association of mild Parkinsonism with orthostatic hypotension, anhidrosis, and an atonic bladder with incontinence—the so-called Shy-Drager syndrome—presents a dramatic picture,⁸⁰⁻⁸¹ which once seen is unlikely to be forgotten. Other features of autonomic dysfunction have been described in this condition, such as impotence, loss of tone in the rectal sphincter, and a fixed cardiac rate. There may be other signs of disorder of the nervous system such as mild cerebellar incoordination, fasciculations, and amyotrophy.

Orthostatic hypotension, which may eventually be of sufficient severity to prevent the patient from even sitting in a chair, is usually the first problem, the symptoms of Parkinsonism only developing slowly and after a variable interval. In one reported case the Parkinsonian features did not appear for three years.⁸²

The condition is progressive and the outlook almost uniformly bad. Shy-Drager syndrome should be thought of in any patient with Parkinsonism presenting with syncopal attacks or postural dizziness. It is more likely, however, that the patient will first have been seen by a cardiologist, the orthostatic hypotension recognized, and a neurological opinion sought concerning minor striatal or cerebellar signs. It is not known at present what proportion of patients with severe, primary orthostatic hypotension go on to develop the full picture of the Shy-Drager syndrome.

PROGRESSIVE SUPRANUCLEAR PALSY (STEELE-RICHARDSON-OLSZEW-SKI SYNDROME)

Within the last few years a chronic degenerative disorder of unknown aetiology has been recognized which has probably long been mistakenly labelled "arteriosclerotic Parkinsonism." The disorder occurs in elderly people and seems to have a universally bad prognosis, the mean life expectancy from the onset of symptoms being about six years.⁸³

The condition, now usually known as "progressive supranuclear palsy,"⁸⁴⁻⁸⁵ is characterized clinically by mask-like facies, hypokinesia, spastic dysarthria and dysphagia, falls, a profound increase of tone of the axial and nuchal muscles (often resulting in an extended posture of the head), mild to moderate dementia, and a diagnostically crucial abnormality of ocular motility. There is a paralysis of vertical conjugate gaze, at first

for willed movements only, but subsequently also for following movements and the upward movement of the eyeballs which occurs on closing the lids. At a later stage, lateral conjugate gaze is also affected, the loss again first involving voluntary movements and at a later stage reflexly induced or following movements.

Tremor is not a feature of this condition, which is unlikely to be missed if ocular movements are tested routinely in every case of "Parkinsonism."

CREUTZFELDT-JAKOB DISEASE

This disease, which may occasionally show striatal features, broadens our vista of pathogenetic agents capable of causing Parkinsonism. It is the second chronic degenerative disease of the human nervous system (after kuru) to have been transmitted to chimpanzees⁸⁶ and is probably due to a "slow virus."

For a long time the disease was thought to be a slowly progressive disorder characterized by Parkinsonism, dementia with pyramidal signs, and sometimes wasting of the hands and feet.²⁵ The clinical spectrum covered by the name Creutzfeldt-Jakob disease has been broadened considerably in recent years⁸⁶⁻⁸⁸ and now includes patients presenting with more rapidly progressive dementia, mutism, and myoclonus—or even cerebellar ataxia.⁸⁹ The symptoms in all these patients have been found to arise on a similar pathological substratum. Patients previously described as suffering from "spongiform encephalopathy"⁹⁰ are now thought to have Creutzfeldt-Jakob disease.

Creutzfeldt-Jakob disease should be suspected when Parkinsonian features are seen in the context of a progressive and widespread neurological disorder, involving any of the fore-mentioned symptoms or signs. The presence of associated cerebral arteriosclerosis may, for a while, delay recognition.

OLIVOPONTocerebellar Atrophy

Occasional patients with olivopontocerebellar atrophy, which usually presents as a progressive cerebellar ataxia, may exhibit generalized rigidity, expressionless facies, tremor at rest (in addition to intention tremor), drooling of saliva, and dementia.⁹¹ A dominant mode of inheritance has been recorded in the affected kinships. The positive family history and associated dementia and cerebellar signs help differentiate this rare condition from more common forms of Parkinsonism.

PROGRESSIVE PALLIDAL ATROPHY

In the absence of encephalitis or of the administration of drugs, Parkinsonism in a child is extremely rare. Progressive pallidal atrophy is occasionally the pathological substratum of cases of juvenile Parkinsonism,⁹²⁻⁹³ though "pure" pallidal atrophy more often results in choreo-athetotic disorders or torsion dystonia.⁹⁴

Parkinsonism—Neuropathology

P. D. LEWIS

British Medical Journal, 1971, 3, 690-692

Experimental studies, often of great value in elucidating disorders of central nervous function, have been of limited use in studying Parkinsonism. The full Parkinsonian syndrome cannot be reproduced in animals, though if the midbrain of monkeys is damaged bilaterally to cause loss of cells in the substantia nigra, tremor and hypokinesia may appear.⁹⁵⁻⁹⁶ The relation of the deep grey masses of the cerebrum to involuntary movement is well known, and the importance of the substantia nigra in this respect is suggested by the many anatomical studies in various species revealing its connexion with the corpus striatum. Nigrostriatal fibres have been shown by conventional histological means,⁹⁷⁻⁹⁹ and a nigrostriatal pathway has been convincingly demonstrated by formalin-induced fluorescence. This technique can be used to show the localization of dopamine in microscope sections, and reveals high concentrations of this catecholamine both in nigral cell bodies and in striatal nerve endings.¹⁰⁰ Since experimental lesions of the brainstem which

produce loss of cells in the substantia nigra reduce the concentration of dopamine in the corpus striatum,¹⁰¹ it is not unreasonable to infer that the striatal depletion of dopamine in Parkinsonism¹⁰² is the result of nigral damage. That lesions of the nigrostriatal pathway are responsible for Parkinsonism is indeed borne out by the neuropathological findings.

Until the end of the last century lack of neurophysiological knowledge and clinicopathological correlations allowed only guesses at the nature of Parkinsonism. James Parkinson himself thought that it originated in a damaged cervical spinal cord and lower brainstem, while preservation of intellectual function suggested to him that the cerebral hemispheres were spared.⁹⁷ In contrast, William Gowers, noting—ten years before the concept of an extrapyramidal pathway had been formulated—that Parkinsonism was a purely motor disorder, believed it to arise from a disturbance of the cerebral cortex.⁹⁹

The midbrain and substantia nigra were first implicated in 1893.¹⁰³ Blocq and Marinesco's case, a man of 38 with left-sided Parkinsonian signs, was shown at necropsy to have a circumscribed lesion 2.5 cm in diameter replacing the right half of the substantia nigra and adjacent structures. The suggestion that lesions of the nigra might be responsible for idiopathic Parkinsonism was made soon afterwards,¹⁰⁴ but neuropathologists, concentrating their efforts on the basal ganglia, seem to have ignored this region for more than twenty years. In 1919 Tretiakoff examined the brains of nine patients with Parkinsonism and observed a variety of degenerative features and reduction of