Published in final edited form as:

Methods Enzymol. 2006 ; 419: 285–321.

Stem Cells in the Lung

Xiaoming Liu, Ryan R. Driskell, and John F. Engelhardt

Abstract

The lung is composed of two major anatomically distinct regions—the conducting airways and gas-exchanging airspaces. From a cell biology standpoint, the conducting airways can be further divided into two major compartments, the tracheobronchial and bronchiolar airways, while the alveolar regions of the lung make up the gas-exchanging airspaces. Each of these regions consists of distinct epithelial cell types with unique cellular physiologies and stem cell compartments. This chapter focuses on model systems with which to study stem cells in the adult tracheobronchial airways, also referred to as the proximal airway of the lung. Important in such models is an appreciation for the diversity of stem cell niches in the conducting airways that provide localized environmental signals to both maintain and mobilize stem cells in the setting of airway injury and normal cellular turnover. Because cellular turnover in airways is relatively slow, methods for analysis of stem cells *in vivo* have required prior injury to the lung. In contrast, *ex vivo* and *in vitro* models for analysis of airway stem cells have used genetic markers to track lineage relationships together with reconstitution systems that mimic airway biology. Over the past decades, several widely acceptable methods have been developed and used in the characterization of adult airway stem/progenitor cells. These include localization of label-retaining cells (LRCs), retroviral tagging of epithelial cells seeded into xenografts, air–liquid interface cultures to track clonal proliferative potential, and multiple transgenic mouse models. This chapter reviews the biologic context and use of these models while providing detailed methods for several of the more broadly useful models for studying adult airway stem/progenitor cell types.

Introduction

The adult lung is lined with numerous distinct types of epithelial cells in various anatomical regions, progressing proximally in the order trachea → bronchi → bronchioles → alveoli. The conducting airways (including the trachea, bronchi, and bronchioles) constitute a minority of the surface area in the human lung, amounting to approximately 0.25 m², whereas the alveolar regions of the lung make up approximately 100 m². In a normal steady state airway, epithelial cell proliferation in the adult lung is much lower compared with highly proliferative compartments in the gut, skin, and hematopoietic system. As in other adult tissues and organs, stem cells in the adult lung are a subset of undifferentiated cells with the capacity to maintain multipotency in the context of the physiologic domain in which they reside. In this context, adult lung stem cells give rise to transient amplifying (TA) progenitor cells capable of abundant self-renewal and regeneration of specific cell lineages in the lung. According to their position within the pulmonary tree, several cell types in the lung have been suggested to act as stem/progenitor cells in response to injury, and to serve to effect local repair (Emura, 2002; Engelhardt, 2001; Otto, 2002). Basal cells, Clara cells, and cells that reside in submucosal glands have been shown to function as progenitors or stem cells in the conducting airway of mice (Borges *et al.*, 1997; Borthwick *et al.*, 2001; Engelhardt, 2001; Engelhardt *et al.*, 1995; Hong *et al.*, 2004a). In addition, a subset of variant Clara cells residing within neuroepithelial bodies (NEBs) (Hong *et al.*, 2001; Peake *et al.*, 2000; Reynolds *et al.*, 2000a,b) or bronchoalveolar duct junctions (Giangreco *et al.*, 2002; Kim *et al.*, 2005) has been considered stem cells in bronchioles. In contrast, alveolar epithelial type II cells (AEC II) are thought to function as the progenitor of the alveolar epithelium, based on their capacity to replicate and to give rise to terminally differentiated alveolar type I cells (AEC I) (Mason *et al.*, 1997). More

recently, “side population” (SP) cells, exhibiting Hoechst dye efflux properties similar to those of hematopoietic stem cells, have been isolated and identified as a putative adult stem cell population for alveolar regions of the adult lung (Giangreco *et al.*, 2004; Summer *et al.*, 2003). These SP cells appear to have both mesenchymal and epithelial potential.

Despite these advances in the study of lung stem cells, a single lung stem cell that can give rise to multiple epithelial lineages in both the proximal and distal airways of the lung has not been identified. To date, the concept of a pluripotent stem cell for all regions of the adult lung has not been widely accepted, based on a developing concept that local regional niches are required to control both the phenotype and expansion of stem cells across vast distances of biologically distinct airways. Thus, stem cell biology in the adult lung is most widely investigated in distinct regions of tracheobronchial, bronchiolar, and alveolar (AEC II) epithelium (Berns, 2005; Bishop, 2004; Emura, 1997, 2002; Engelhardt, 2001; Liu *et al.*, 2004; Mason *et al.*, 1997; Neuringer and Randell, 2004; Otto, 2002).

Most of our current understanding of stem cells in the adult lung originates from research using classic lung injury models and/or epithelial reconstitution models. These models will likely continue to be the most useful approaches to identify stem cell populations and to understand the regulation of stem cell fates in the adult lung. However, as the molecular understanding of genes that control lung stem cell populations unravels, improved genetic approaches to study stem cell phenotypes in their niches will likely emerge in combination with more traditional approaches of analysis. This chapter first reviews the diversity of cell types in the adult lung. We then summarize the current existing knowledge and describe classic experiments used to characterize potential stem cell populations and stem cell niches in the adult lung. Last, we describe several methods for characterizing stem cells in the adult airway and provide detailed protocols for stem cell labeling, the use of lung injury models to mobilize airway stem cells, and the reconstitution of airway epithelia and submucosal glands, using *in vitro* and *ex vivo* models.

Anatomical and Cellular Diversity of Adult Lung

The adult lung can be functionally and structurally divided into three epithelial domains: the proximal cartilaginous airway (trachea and bronchi), distal bronchioles (bronchioles, terminal bronchioles, and respiratory bronchioles) and gas-exchanging airspaces (alveoli). Each of these domains has historically been classified on the basis of morphologic criteria that define the unique type of airway epithelium within each domain of the lung (Fig. 1). Consequently, defining functional criteria for the diversity of epithelial cell types in the airway has been a major focus.

In the murine proximal conducting airway and major bronchi, basal, Clara, and ciliated cells are predominant cell types, although less frequent pulmonary neuroendocrine cells (PNECs) also reside in this domain. The nonciliated, columnar Clara cells comprise the majority of the distal bronchiolar epithelium in mice, whereas the alveolar epithelium is lined by squamous alveolar type I pneumocytes (AEC I) and cuboidal type II pneumocytes (AEC II). Unique to the proximal cartilaginous airways are submucosal glands (SMGs) that are contiguous with surface airway epithelium (Fig. 1). Although less abundant in mice and confined to the trachea, SMGs are a major secretory structure located in the interstitium beneath the cartilaginous airway of other nonrodent species (Choi *et al.*, 2000; Liu *et al.*, 2004; Plopper *et al.*, 1986; Widdicombe *et al.*, 2001). SMGs are composed of an interconnecting network of serous and mucous tubules that secrete antibacterial factors, mucus, and fluid into the airway lumen. In contrast, the alveolar epithelium of adult lung is highly similar among mammalian species (Ten Have-Opbroek and Plopper, 1992).

Most of the current studies on stem cell biology in the adult airway have employed rodent animal models. To this end, it is important to note the anatomical and cellular differences between rodent and human airways. In mice, Clara cells reside throughout the tracheobronchial and bronchiolar epithelium and are the predominant secretory cell type in airways of this species. In contrast, Clara cells are limited to the bronchioles of human airways, and goblet cells are the predominant secretory cell type in the human tracheobronchial airway. In contrast to humans, the predominant secretory cell type in the rat conducting airway is the serous cell. However, goblet cells in mice and rats can be induced by specific cytokine stimuli and/or injury (Liu *et al.*, 2004). A summary of cellular differences in the conducting airway epithelium and SMGs between rodents (mouse and rat) and primates (monkey and human) is listed in Table I. Despite the notable differences between rodents and primates in the cell biology of the airway epithelium and the abundance of SMGs in the conducting airway, studies in mice have played significant roles in investigating stem cell biology in proximal and distal airways because of their versatile genetics.

Stem Cell Phenotypes and Niches in Adult Lung

Studies in stem cell biology have demonstrated that cell fate and the maintenance of stem cell populations are regulated by their local anatomical and chemical microenvironment, or niche. Niches are discrete microenvironments of specialized cell types, matrix, and diffusible factors such as cytokines and growth factors, which are critical for maintaining stem cells and promoting appropriate cell fate and migration decisions (Watt and Hogan, 2000). In the adult lung, candidate stem cell populations have been identified that are restricted to the tracheal submucosal gland ducts, neuroepithelial bodies (NEBs) of the bronchi and bronchioles, and bronchoalveolar duct junction (BADJ) of the terminal bronchioles, suggesting distinct regional stem cell niches in the adult lung (Borthwick *et al.*, 2001; Engelhardt, 2001; Engelhardt *et al.*, 1995; Giangreco *et al.*, 2002; Hong *et al.*, 2001, 2004a; Kim *et al.*, 2005).

Because stem cells divide infrequently, injury to the lung has been necessary to study lung stem cell phenotypes and their niches in the airway. This slow-cycling characteristic has allowed for the use of DNA-labeling techniques with detectable nucleotide analogs to track stem cells *in situ*. Bromodeoxyuridine (BrdU) is one such common nucleotide analog that is classically used to track label-retaining cells (LRCs) after a prolonged “washout” period that dilutes the label within the more rapidly cycling transient-amplifying (TA) cells.

Two methods of injury used to amplify mobilization of the stem cell compartment in the proximal airway and to identify LRCs have included intratracheal instillation of povidone-iodine or inhalation of SO₂ (Borthwick *et al.*, 2001). Intratracheal instillation of povidone-iodine or inhalation of SO₂, followed by *in vivo* BrdU administration, labels airway epithelial cells throughout the entire trachea at early time points (Borthwick *et al.*, 2001). However, longer time points postlabeling revealed LRC accumulation in submucosal gland (SMG) ducts of mouse trachea, suggesting that this region provides a protective niche for stem cells (Borthwick *et al.*, 2001; Engelhardt, 2001). An example of this LRC patterning after naphthalene injury in mouse trachea is shown in Fig. 2. The ability of glandular cell types to repopulate the surface airway epithelium of the trachea was also studied by subcutaneous transplantation of epithelium-denuded tracheas into recipient mice (Borthwick *et al.*, 2001). By 28 days after transplantation, outgrowth of glandular cells led to repopulation of a ciliated surface airway epithelium, suggesting that gland or gland duct cells could regenerate the tracheal epithelium.

It is commonly thought that basal cells have the capacity to produce all the major cell phenotypes found in mouse trachea, including basal, ciliated, goblet, and granular secretory cells (Hong *et al.*, 2004b; Schoch *et al.*, 2004). In humans, where the proximal airway is more pseudostratified, parabasal cells (which are located just above the basal cells, and may also be

considered intermediate cells) are thought to contribute more to the proliferative cell population than basal cells, based on higher-level expression of the proliferation marker MIB-1 (Ki-67) (Boers *et al.*, 1998). Hence, parabasal cells (or intermediate cells) are thought to act as a TA cell population in the proximal airway that is derived from a subset of basal stem cells. Studies using human bronchial xenografts and clonal analysis with retroviral vectors also support this concept (Engelhardt *et al.*, 1995).

Clara cells have long been thought to act as progenitor cells of the airway. This type of cell resides throughout all levels of the mouse conducting airways, but is confined to bronchiolar airways in humans. A number of studies have suggested that a subset of Clara cells are stem cells in the bronchiolar epithelium of mice. Lung injury studies using naphthalene depletion of Clara cells revealed that a subset of variant Clara cells shows multipotent differentiation and the ability to regenerate the bronchiolar epithelium. This population of Clara cells is cytochrome *P*-450 2F2 (CYP2F2) negative, resides in discrete pools associated with NEBs and at BADJs. Further characterization of these cells has revealed that they express stem cell antigen (Sca-1), and have the ability to efflux Hoechst dye (Reynolds *et al.*, 2000a,b).

Pulmonary neuroendocrine cells (PNECs) reside within NEBs at cartilage–intercartilage junctions in the proximal airway and airway junctions in the bronchioles. These cells or group of cells are thought to play a role in lung development and postnatal lung maintenance through the release of neuropeptides. After naphthalene injury, proliferative cells that express either Clara cell secretory protein (CCSP) or the PNEC marker calcitonin gene-related peptide (CGRP) accumulate within NEBs (Hong *et al.*, 2001). To exclude the possibility that CGRP-expressing PNECs are the potential stem cell compartment in bronchioles, studies were performed with transgenic mice expressing the thymidine kinase gene under the direction of the CCSP promoter. Expression of thymidine kinase specifically in Clara cells allowed for the specific ablation of naphthalene-induced replicating CCSP-expressing cells. Cell-specific ablation was facilitated by administration of the prodrug ganciclovir, which is converted by thymidine kinase to a toxic nucleotide analog that prevents DNA replication and hence kills cells. Under these conditions that ablated all CCSP-expressing Clara cells after naphthalene injury, hyperplasia of CGRP-expressing PNECs occurred; however, a ciliated epithelium could not be regenerated. Consequently, these studies demonstrated that CGRP-expressing PNECs are not a stem/progenitor cell population in the distal airway. Rather, PNECs are thought to provide a niche that regulates expansion of a CCSP-expressing stem cell population in mouse distal airways (Hong *et al.*, 2001).

More recent studies have also found, in another region in the terminal bronchioles, that is, the bronchioalveolar duct junctions, bronchioalveolar stem cells (BASCs) that accumulate LRCs. BASCs were previously termed DPCs (double-positive cells that express both SP-C and CCSP) (Giangreco *et al.*, 2002; Kim *et al.*, 2005). These cells appear to be resistant to bronchiolar and alveolar damage, proliferate during epithelial cell renewal *in vivo*, and are thought to function to maintain the bronchiolar Clara cell and alveolar cell populations of the distal lung. BASCs also possess characteristics of regional stem cells, such as self-renewal and multipotency in clonal assays (Kim *et al.*, 2005).

In the gas-exchanging regions of the lung, alveolar type II cells (AEC II) have been considered the stem/progenitor cell of the alveolar epithelium, based on their ability to repopulate both AEC II and AEC I after injury (Giangreco *et al.*, 2002; Kim *et al.*, 2005; Reynolds *et al.*, 2004). Studies using a rat lung injury model have suggested that there may be four AEC II groups, based on their expression of markers. The subpopulation of AEC II that is E-cadherin negative, proliferative, and expresses high levels of telomerase activity was considered a stem cell candidate for alveolar epithelium (Reddy *et al.*, 2004).

In contrast to insights regarding candidate stem cells in the respiratory epithelium of adult lung, much less information is available regarding stem cells in the vascular compartment of the lung. One study demonstrated that an extremely small “spore-like” cell population in the lungs of adult sheep and rats with low oxygen demand can generate lung-like alveolar tissue *in vitro* (Vacanti *et al.*, 2001). Side population (SP) cells isolated from the lung have also been shown to express stem cell markers indicative of epithelial and mesenchymal lineages. SP cells isolated from whole lung, using Hoechst 33342 efflux and other stem cell markers, comprise 0.03–0.07% of mouse lung cells and are Sca-1 antigen positive, lin negative, heterogeneous for CD45, and express the vascular marker CD31 (Giangreco *et al.*, 2004; Summer *et al.*, 2003). On the basis of the expression of hematopoietic marker CD45, lung SP cells were further subdivided into hematopoietic (CD45-positive) and non-hematopoietic (CD45-negative) subpopulations. Nonhematopoietic SP cells express markers of epithelial and mesenchymal cells, share some characteristics with airway stem cells, and are currently under further investigation (Giangreco *et al.*, 2004). Consequently, there is not a single unique approach to isolate stem cell populations from the intact adult lung; however, isolation of BASCs and SP cells from the lung has been described (Giangreco *et al.*, 2004; Kim *et al.*, 2005). We have summarized in Table II the current knowledge of epithelial cell types in the conducting and respiratory airways, their specific cellular markers, their potential as stem/progenitor cells, and their potential lineage relationship.

In Vivo Injury Models of the Lung

Overview

Stem cells are slow cycling in the setting of normal tissue turnover, giving rise to transient amplifying (TA) progenitor cells that retain the capacity to replicate and impart most of the tissue renewal in the presence of injury. However, unlike stem cells that have unlimited proliferative capacity, TA cells are eventually incapable of proliferation and become terminally differentiated (TD) cells. Injury models have played key roles in mobilizing stem cell compartments in the lung. Different epithelial cell types in distinct areas of the lung are sensitive to various agents and have been used in lung injury models to investigate stem cells in the adult lung (Table III). These include hyperoxic agents (Smith, 1985), radiation (Theise *et al.*, 2002), SO₂ or polidocanol (Borthwick *et al.*, 2001; Randell, 1992), bleomycin (Bigby *et al.*, 1985; Izibicki *et al.*, 2002), elastase (Dubaybo *et al.*, 1991), and naphthalene (Kim *et al.*, 2005; Van Winkle *et al.*, 1999). Detailed methods for polidocanol and naphthalene injury are described later, and the others are briefly introduced here.

Exposing the lung to high concentrations of oxygen (O₂) or oxidant gases such as nitrogen dioxide (NO₂) or ozone (O₃) has been used to generate hyperoxic lung injury models. Cell death from hyperoxic injury may occur through either apoptotic or nonapoptotic pathways, possibly via free oxygen radicals. The biochemical, cellular, and morphologic characterizations of hyperoxic lung injury have been extensively studied (Smith, 1985). Mice show mild damage after 3 days of exposure to high concentrations of O₂, damage that consists of alveolar septal thickening and increases in alveolar macrophages. Extensive damage can be induced by continuous prolonged exposure to O₂, resulting in the destruction of alveolar walls, proteinaceous exudates in alveoli, and large numbers of interstitial and alveolar polymorphonuclear leukocytes (PMNs) (Smith, 1985).

Inhalation exposure of mammals to NO₂ has also been used extensively as a model of controlled acute pulmonary injury and repair. Prolonged exposure of rodents to NO₂ at 20–30 ppm results in mild emphysema and a partially reversible decrease in lung elastin and collagen content. The mechanism by which NO₂ damages AEC I is thought to be oxidation of unsaturated fatty acids of the cell membranes (Evans *et al.*, 1981). Subacute NO₂ exposure (20 ppm NO₂ for 28 days) in mice also results in swelling of the ciliary shaft, focal loss of cilia, and the formation

of compound cilia in the airway epithelium. Such ciliary lesions appear to be reversible when NO₂ exposure is stopped (Ranga and Kleinerman, 1981). In addition to the mouse NO₂ injury lung models, rat (Meulenbelt *et al.*, 1992a), rabbit (Meulenbelt *et al.*, 1994) and sheep (Januszkiewicz and Mayorga, 1994) NO₂ injury lung models have also been described.

An alternative mode of lung injury includes radiation. Mice exposed to 1200 cGy of total body irradiation have lungs that appear hypocellular, with a breakdown of capillaries within alveolar septae and extravasation of erythrocytes into alveolar spaces within 3 days of exposure to radiation. Irradiation-induced alveolar epithelial damage is quickly repaired by AEC II proliferation followed by differentiation in AEC I (Theise *et al.*, 2002). The notion that bone marrow-derived stem cells have the ability to contribute to repair of alveolar epithelium was also suggested by the transplantation of retrovirally tagged green fluorescent protein (GFP) bone marrow cells into irradiated mice (Grove *et al.*, 2002).

Another mode of injury to alveolar regions of the lung includes bleomycin. Bleomycins are a family of compounds with antibiotic and antitumor activity that have the major side effect of pulmonary toxicity. Intratracheal instillation of bleomycin induces alveolar injury in the lung (Hay *et al.*, 1991). The cellular toxicity of bleomycin is based on its bithiazole component that partially intercalates into the DNA helix and causes oxidative damage to DNA. However, lipid peroxidation caused by bleomycin may also account for alveolar damage and subsequent pulmonary inflammation. Bleomycin lung injury is characterized by pulmonary fibrosis due to increased production of collagen and other matrix components in the lung (Bigby *et al.*, 1985; Hay *et al.*, 1991; Izbicki *et al.*, 2002). The AEC I are particularly sensitive to bleomycin and are the first to be injured, whereas AEC II have a more variable sensitivity. Bleomycin-resistant AEC II undergo metaplasia in the presence of the drug (Izbicki *et al.*, 2002). More details regarding the generation of bleomycin injury for the characterization of stem cells have been described by Kim and colleagues studying the BASC niche at the BADJ (Kim *et al.*, 2005).

Sulfur dioxide (SO₂), a common pollutant in the air responsible for many diseases of the respiratory system, has been used to induce injury in the proximal airways of animal models. Various species of animals have different sensitivities to SO₂—mice are most sensitive, whereas rats are the most tolerant. Inhalation of SO₂ has been used in pulmonary epithelial injury models of various species including mice, rats, sheep, guinea pigs, and swine. SO₂-induced lung injury in the rat and sheep has also been proposed as a model for bronchitis (Borthwick *et al.*, 2001; Lamb and Reid, 1968). SO₂ is soluble in water, and therefore is easily absorbed into the wet mucous membranes of the airway, causing the greatest damage to the trachea and large bronchi, whereas the bronchioles are invariably spared (Asmundsson *et al.*, 1973). This feature makes inhalation of SO₂ an important injury model for the proximal airway.

Another chemical used to induce injury in the proximal airway of mice includes polidocanol, a surface-active detergent clinically used to enhance absorption of small proteins/peptides. Other surface-active agents such as polyoxyethylene 9 lauryl ether (Laureth-9), sodium glycocholate, and Triton X-100 have also been shown to induce lung injury after intratracheal instillation (Borthwick *et al.*, 2001). Direct intratracheal instillation of 10 µl of 2% polidocanol in phosphate-buffered saline (PBS) to the mouse trachea causes widespread denudation of the airway epithelium by 24 h postinjury (Fig. 3D). Using this polidocanol injury model, Borthwick and colleagues have identified the existence of stem cell niches in SMGs of the proximal mouse airway (Borthwick *et al.*, 2001). A detailed method of this approach is described in a later section.

Naphthalene, a chemical agent that is toxic to Clara cells, can induce damage to both proximal and distal airways of mice. This injury method has been used to identify two distinct stem cell

niches associated with NEBs in the proximal airway and BADJ (Giangreco *et al.*, 2002; Hong *et al.*, 2001; Reynolds *et al.*, 2000a,b). Naphthalene is metabolized by the 2F2 subtype of cytochrome *P*-450, resulting in the production of a cytotoxic epoxide. Inhalation or intraperitoneal injection of naphthalene produces a dose-dependent cytotoxicity to Clara cells, whereas cells deficient in CYP2F2 are spared (Buckpitt *et al.*, 1992; West *et al.*, 2001). The bronchiolar epithelium of mice is a particularly sensitive site for naphthalene toxicity, given its high content of Clara cells (Giangreco *et al.*, 2002; Hong *et al.*, 2001). At higher doses of naphthalene (275 mg/kg), however, significant injury to the tracheobronchial epithelium can also be achieved (Fig. 2A). As discussed in more detail previously, a subpopulation of variant CCSP-expressing Clara cells deficient in CYP2F2 has been suggested as the principal stem/progenitor cell in the bronchiolar epithelium that facilitates repair after administration of naphthalene (Giangreco *et al.*, 2002; Hong *et al.*, 2001; Kim *et al.*, 2005; Peake *et al.*, 2000; Reynolds *et al.*, 2000a,b). Detailed methods for naphthalene-induced mouse lung injury and the identification of LRCs by BrdU labeling are presented in the following section.

Mouse Naphthalene Injury and BrdU-Labeling Protocol

1. Weigh the appropriate amount of naphthalene (cat. no. N143-500; Fisher Scientific, Barrington, IL) directly into a 15-ml polypropylene conical tube; make fresh on the day of use.
2. Calculate the volume of sterile corn oil (Mazola; density of corn oil, 0.9185 g/ml) needed to achieve the desired concentration of naphthalene (27.5 mg/ml). In general, the concentration of naphthalene used is 27.5 mg/ml; however, the dose can be adjusted to achieve different levels of injury to the proximal and/or distal airways. To generate a 27.5-mg/ml naphthalene stock sufficient for a 10-mouse experiment, 110 mg of naphthalene is weighed into a conical tube followed by the addition of 3.65 g of corn oil (4 ml $\times$ 0.9185 g/ml).
3. Naphthalene is difficult to dissolve. Vortex the solution at full speed for several minutes, and continue mixing on a tube rotator for at least 20 min at room temperature. Make sure that all the naphthalene has dissolved by verifying that the solution is clear of solid naphthalene.
4. Load the naphthalene solution into a 1-ml syringe, using an 18-gauge needle. After loading, remove the 18-gauge needle and use a 26-gauge needle for intraperitoneal injection.
5. Weigh the mice to the nearest 0.1 g to calculate the volume dose. It is particularly important that the dosing be exact for reproducible injury. In addition, it is important to inject the mice before 10:00 A.M. to take advantage of the minimal glutathione levels at this time of day. Sex differences exist in the extent of lung injury at a given dose of naphthalene (female mice are more sensitive than males), so make sure all animals are of the same sex.
6. Administering light anesthesia to the mice will help in obtaining an accurate weight and will also make the injection easier. Intraperitoneal injections of the freshly made naphthalene solution described previously are performed at 10 μ l/g body weight (i.e., 250 μ l for a 25-g mouse given a total dose of 275 mg/kg). Naphthalene-injured animals will present a ruffled appearance with crusted eyes at approximately 20–36 h postinjury.
7. To identify LRCs, mice can be injected intraperitoneally with a 80- to 100-mg/kg body weight dose of BrdU (Roche, Indianapolis, IN)–PBS solution every 24 h, beginning 6 h after the naphthalene injection, for five consecutive days. Multiple injections (at least three) are required to label a significant percentage of LRCs.

8. Maximal injury of airway epithelium is typically observed 2 days after naphthalene injection, and newly regenerated airway epithelium can be seen by 1 week after injury (Fig. 2B). To localize LRC stem cell populations, mice are killed at various time points, such as 21, 42, and 90 days after injury and BrdU labeling.
9. Harvest the trachea and/or lung and remove blood by washing the organs in precooled PBS. The tissues can be either fixed in 10% formalin for paraffin embedding and sectioning or freshly frozen in optimal cutting temperature (O.C.T.) compound embedding medium (Tissue-Tek; Sakura Finetek USA, Torrance, CA).
10. BrdU incorporation is detected by immunofluorescence or immunohistochemical staining of 10- μ m frozen sections or 6- μ m paraffin sections, respectively.
11. Bring frozen sections to room temperature for 10–20 min to dry. Fix the sections in 4% paraformaldehyde in PBS for 20 min before processing for epitope retrieval.
12. Use an antigen-unmasking solution (cat. no. H-3300; Vector Laboratories, Burlingame, CA) to retrieve epitopes as specified by the manufacturer.
13. Rinse the sections in three changes of PBS (2 min for each), and then block nonspecific binding of immunoglobulin by incubating the sections in 5% rabbit serum (the same species to which the secondary antibody was produced) in PBS at room temperature for 30–60 min.
14. Directly apply a 1:100 dilution of mouse anti-BrdU antibody (cat. no. 1–299–946; Roche) in 5% rabbit serum–PBS to sections and incubate at room temperature for 1 h.
15. Rinse the sections in three changes of PBS (3 min for each) before incubating with a 1:200 dilution of fluorescein isothiocyanate (FITC)- conjugated rabbit anti-mouse IgG in 1.5% rabbit serum–PBS (cat. no. 315096045; Jackson ImmunoResearch, West Grove, PA) at room temperature for 45–60 min.
16. Rinse the sections in three changes of PBS (3 min for each) and mount them in VECTASHIELD mounting medium containing 4', 6-diamidino-2-phenylindole (DAPI, cat. no. H-1200; Vector Laboratories). Evaluate BrdU staining under a fluorescence microscope (Fig. 2C).

Mouse Polidocanol Injury and BrdU-Labeling Protocol

1. Tare the balance to zero with a 15-ml polypropylene conical tube and weigh 200 mg of polidocanol (cat. no. P-9641; Sigma, St. Louis, MO) in the tube. Add sterile PBS to bring the total volume up to 10 ml, making a 2% polidocanol (w/v) solution. Rotate the tube to ensure the polidocanol is completely dissolved.
2. Anesthetize the mice with ketamine (80 mg/kg) and xylazine (10 mg/kg) in sterile PBS. The dose of anesthesia should not be so deep as to hinder respiration, so it is advised to adjust this dose to the strain of mouse being used.
3. To facilitate intubation of mice for delivery of polidocanol into the trachea, hang the mouse on a self-made wire rack with a suture loop around the incisor teeth (Fig. 3A and B). A small flashlight can be placed under the neck of the mouse to aid in visualization of the tracheal orifice if needed.
4. Open the mouth of the mouse and pull its tongue out and down, using flat jaw forceps. Remove the needle from a 0.55-mm Angiocath (BD Angiocath; BD Medical, Sandy, UT) and insert the plastic catheter into the trachea.

5. Use a 25-gauge needle to apply 20 μ l of 2% povidone-iodine into the trachea through the catheter. Leave the catheter in the trachea for 1 min after delivering the solution to prevent immediate expulsion into the mouth.
6. After the instillation, intraperitoneal injection of BrdU may be performed as described previously.
7. Massive injury to the tracheal epithelium is seen 24 h postinjury, and the epithelium is completely regenerated in 1–2 weeks (Fig. 3D and E). LRC location can also be performed as described for the naphthalene model.

Ex Vivo Epithelial Tracheal Xenograft Model to Study Stem Cell Expansion in Proximal Airway

Overview

The *ex vivo* epithelial xenograft model is an approach that has historically been used to study progenitor-progeny relationships in the adult airway (Duan *et al.*, 1998; Engelhardt *et al.*, 1991, 1992, 1995; Presente *et al.*, 1997; Sehgal *et al.*, 1996). This model seeds isolated airway epithelial cells onto graft tracheas that have been denuded of all endogenous airway epithelia by freeze–thawing. After the airway stem/progenitor cells are seeded, the tracheal grafts are implanted subcutaneously into immunocompromised hosts, such as *nu/nu* or severely compromised immunodeficient (SCID) mice. A fully differentiated epithelium regenerates approximately 3–4 weeks after transplantation (Filali *et al.*, 2002).

A major advantage of this approach is that it can be applied to airway epithelia from multiple species. This model has been used study airway stem cell biology and/or lineage, using purified or enriched specific epithelial cell types to reconstitute the denuded tracheal grafts. These studies have demonstrated the ability of basal and secretory cells to reconstitute a fully differentiated surface airway epithelium in multiple species (Hook *et al.*, 1987; Inayama *et al.*, 1988; Randell, 1992). Rabbit basal cells purified by centrifugal elutriation have the ability to repopulate epithelium containing basal, ciliated, and goblet cells after reconstitution of a tracheal xenograft (Inayama *et al.*, 1988). In addition, seeding of enriched rabbit Clara cells into xenografts has demonstrated that this cell type has a limited capacity for differentiation into Clara and ciliated cells (Hook *et al.*, 1987). Although enrichment of a given cell type is greater than 90–95% in most instances, and can reach up to 98% purity after cell sorting with a combination of cellular surface markers and light scattering, one major limitation of such a cell enrichment approach is the potential for small levels of contamination with unidentified airway stem cells.

To circumvent the limitations of cell enrichment approaches, retrovirus-mediated gene transfer has been used to genetically tag epithelial stem/progenitor cells and to follow lineage relationships, using histochemical markers (Engelhardt *et al.*, 1992, 1995). Using this approach, primary human airway cells can be infected with a variety of retroviral vectors before reconstituting the xenograft airway epithelium (Fig. 4A). Histochemical staining for marker transgenes such as β -galactosidase and/or alkaline phosphatase can then be used to visualize clonal expansion of stem/progenitor cells in the reconstituted airway epithelium (Fig. 4B). Using two independent viral vectors with two independent marker transgenes (i.e., β -galactosidase and/or alkaline phosphatase), studies have demonstrated that transgene-expressing clones indeed arise from a single retrovirally infected progenitor/stem cell (Engelhardt *et al.*, 1995). In reconstituted xenografts with retrovirally tagged epithelial cells, the number of cells in each clone (i.e., size) can be used as an index of proliferative capacity, with the notion that clones arising from adult stem cell fractions would have a larger proliferative capacity. Phenotyping of clones (i.e., the types of transgene-expressing cells in

each clone: basal, ciliated, goblet, or intermediate cells) can be used to assess the variety of progenitor cell types in the airway with either limited or pluripotent capacity for differentiation. These studies have suggested that a diverse repertoire of progenitor cells likely exists in the adult human proximal airway, with differing capacities for proliferation and differentiation. These studies demonstrated that a stem/progenitor population with the ability to differentiate into SMGs and all surface airway epithelial cell types exists in the adult human airway (Engelhardt *et al.*, 1995). This cell type is thought to be a candidate stem cell in the proximal airway.

The protocol for the generation of human bronchial xenografts, described later, is also suitable for studies with other species such as rat, with modifications of cell culture conditions (Engelhardt *et al.*, 1991).

Human Bronchial Xenograft Protocol

1. Dissection of human airway tissue:

- a. The bronchial/lung cassette should be chilled immediately after removal from the donor in cold physiologic saline and should remain chilled during dissection. For ideal aseptic and safety conditions, the dissection is best done in a laminar flow hood. Handling human tissue and primary airway cultures may expose the user to blood-borne pathogens, and suitable health protection measures should be followed.
- b. The proximal airways are dissected from the lung on ice and placed immediately in chilled sterile Ca^{2+} - and Mg^{2+} -free PBS containing penicillin (50 U/ml), streptomycin (50 $\mu\text{g/ml}$), tobramycin (80 $\mu\text{g/ml}$), ceftazidime (100 $\mu\text{g/ml}$), imipenem/cilastatin (Primaxin, 100 $\mu\text{g/ml}$), and amphotericin B (5.0 $\mu\text{g/ml}$). Remove the blood, mucus, and extraneous connective tissues from the airways and cut the specimen into suitably sized half-rings and place them into 50-ml polypropylene conical tubes with the preceding antibiotic solution. Invert the tubes for several minutes to wash the tissue, and repeat four times by moving the tissue from one conical tube to a fresh tube with new antibiotic-containing cold PBS.
- c. It is important to avoid exposing specimens to drastic changes in temperature during processing and dissection to prevent epithelial degradation. In addition, segments of the proximal airway (trachea and bronchi) may be separately processed because their epithelial characteristics may vary.
- d. After washing in antibiotic solution, airway tissue is placed in a 50- or 15-ml conical tube filled with medium A (described at the end of this section) to dissociate airway epithelial cells. It is important that the conical tubes be filled to the top with medium A and tightly capped to avoid changes in pH caused by alteration in CO_2 concentration over time. Airway tissues are incubated in medium A at 4° for 24–96 h. Nasal turbinates require 24–48 h for dissociation whereas tracheobronchial tissues require 48–72 h (depending on the desired degree of separation into single cells versus cell clumps). However, it should be noted that dissociation times longer than 72 h will decrease cell viability, and cell clumps appear to proliferate more rapidly in culture.

2. Harvesting airway epithelial cells:

- a. Add 10% fetal bovine serum (FBS) into the dissociation solution to inhibit the Pronase in medium A while transferring the solutions and tissue to a

larger or greater number of conical tubes filled approximately one-third full. Invert the tube(s) several times vigorously to help dissociate the cells from the airway tissue. Mild shaking is helpful to increase yields, but avoid excessive force that may damage cells (i.e., do not shake hard enough to cause the FBS to foam).

- b.** After shaking, allow 1 min for airway tissue to settle to the bottom of the tube (liberated airway cells will remain in suspension). Pipette the medium containing the airway cells into a fresh tube on ice, and place the remaining tissue into a 100-mm tissue culture plate.
- c.** Using the blunt side of a scalpel, scrape the surface of the airways to remove the remaining epithelial cells. Rinse airway samples with 10% FBS–Ham’s F12 medium, combine the washed cells into 50-ml conical tubes, and centrifuge the tubes at 120g for 5 min at 4°.
- d.** Resuspend the cell pellets in medium B (described at the end of this section) and transfer the cell suspension to 100-mm Primaria tissue culture dishes (cat. no. 353803; BD Biosciences Discovery Labware, Bedford, MA).
- e.** Incubate the suspension in a 5% CO₂ incubator at 37° for a minimum of 1–2 h to allow fibroblasts within the cell suspension to attach to the plastic surface. Airway epithelial cells will not attach rapidly to the plastic surface without collagen coating.
- f.** Collect the nonattached cells by centrifugation, resuspend them in medium C (described at the end of this section), and count them with a hemacytometer.

3. Culture of airway epithelial cells:

- a.** Plate $1\text{--}2 \times 10^6$ of the preceding airway cells per 100-mm collagen-coated tissue culture dish (BD Biosciences Discovery Labware) in 10 ml of airway culture medium C. Incubate the cells overnight in a 5% CO₂ incubator at 37°.
- b.** On the day after plating, aspirate the culture medium and unattached cells. Wash the dish containing adherent cells with prewarmed Ham’s F12 medium and refeed the cells with fresh medium C. Culture the cells for an additional 48 h.
- c.** At 72 h postplating, feed the cells with medium D (containing lower levels of antibiotics and no amphotericin B). Typically, cells are ready for cryopreservation, passaging, or transplantation into xenograft models by 5 days postplating (~80% confluency). Care should be taken not to allow cells to become more than 80% confluent or they will begin to differentiate and lose their capacity for subculturing. Unpassaged fresh cultures of airway epithelia also give consistently better reconstitution of epithelium in the xenograft model.
- d.** If primary airway cells are to be expanded (typically cells can be expanded once without the loss of the ability to differentiate in a xenograft model), they should be treated with 0.1% trypsin–EDTA for 1–3 min at 37° followed by neutralization with an equal volume of trypsin inhibitor buffer (recipe described at the end of this section). Cells should be closely monitored during trypsinization and harvested immediately once they are released by gentle tapping of the plate. Cell suspensions are centrifuged to remove trypsin and washed once in medium D, followed by plating at a 1:5 dilution.

- e. For cryo preservation, epithelial cell pellets are resuspended in 4° medium D with 10% dimethyl sulfoxide (DMSO) and 10% FBS and aliquoted in 2-ml cryogenic vials for slow freezing at -80°, overnight. Slow freezing can be performed in isopropanol-containing cryopreservation containers (Nalgene; Nalge Nunc International, Rochester, NY). Cells are then moved to liquid nitrogen storage. Typically, one subconfluent 100-mm plate of cells is aliquoted per vial, and when subcultured placed into five 100-mm plates.
4. Retroviral infection of primary airway epithelial cells (optional): Retroviral infection of airway epithelial cells before seeding into tracheal xenograft has been useful in studying airway stem/progenitor cell biology (Engelhardt *et al.*, 1995) (Fig. 4A and B). Briefly, freshly isolated primary airway epithelial cells on the second day postseeding (approximately 10% confluency) can be incubated in the presence of serum containing conditioned retroviral or lentiviral producer supernatant (10 ml of retroviral supernatant per 100-mm plate of primary cells) for 2 h in the presence of Polybrene (2 µg/ml; Sigma). After each retroviral infection, the cells are washed twice with F12 medium before the addition of hormonally defined medium D. Cells can be infected up to three times on sequential days. Typically, viral titers of 1×10^6 colony-forming units/ml are capable of transducing primary airway cells at an efficiency of 10–30% after three serial infections. Serum-free concentrated stocks of vesicular stomatitis virus glycoprotein-pseudotyped retrovirus or lentivirus can also be used to achieve higher transduction efficiencies (our unpublished data).
5. Preparation and construction of xenograft cassettes: A xenograft cassette is assembled by connecting tubing and adapters fitted to the length of the donor rat trachea and secured with silk sutures as illustrated in Fig. 4C (materials and vendors are described in the caption to Fig. 4). Preassembled parts of the cassette are placed in a 100-mm tissue culture plate, sealed in a sterilization pouch, and gas sterilized before anchoring tracheal xenografts.
6. Preparation of denuded rat tracheas:
 - a. Rats are killed by CO₂ asphyxiation and pinned to a Styrofoam bed. Tracheas are excised from the pharynx to the carina under sterile conditions, using 70% ethanol to clean the site of incision. Place the tracheas in separate 2-ml screw cap tubes and place them on ice immediately after excision.
 - b. After all tracheas have been harvested, denude the tracheas of all viable epithelium by freeze–thawing them three times: freezing at -80° and thawing at room temperature.
 - c. After the third round of freeze–thawing, clean the tracheas of excessive fat and cut them to size (typically from the first to thirteenth cartilage ring). Rinse each tracheal lumen with 10 ml of precooled modified Eagle’s medium (MEM).
 - d. Pair tracheas of similar length in the same tube. The denuded tracheas can be stored at -80° for prolonged periods of time before transplantation.
7. Seeding tracheas with primary airway epithelial cells:
 - a. Airway epithelial cells cultured in 100-mm dishes (at ~80% confluence) are harvested with trypsin and trypsin inhibitor as described earlier, resuspended in medium D at a concentration of $1\text{--}2 \times 10^6$ cells per 20 µl, and kept on ice.
 - b. Ligate the rat tracheas to the adaptor *e* attached to tubing *b* (as shown in Fig. 4C) with securely tied triple-knotted sutures, and loop the sutures around the tubing and trachea a total of three times.

- c. Inject 20–25 μ l of medium containing $1\text{--}2 \times 10^6$ cells into the open end of the rat trachea, using a micropipettor under sterile conditions. Insert the pipette tip as deeply as possible into the trachea and slowly withdraw the pipette as cells are injected into the lumen of the trachea (taking care not to allow cells to leak out).
 - d. Ligate the remaining open end of the rat trachea to adapter *e'* attached to tubing *c* as shown in Fig. 4C. Secure the length of the rat trachea by stretching it to physiologic length and clamping the tubing *b* and *a* with a hemostat. Tie the remaining two sutures as shown in Fig. 4C to secure the tracheas to adapter *e'* and adapter *e'* to tubing *d*.
 - e. Place the cell-seeded xenograft cassettes into a 100-mm tissue culture dish with 1–2 ml of medium D overlaid on top of the trachea to keep it in moist.
 - f. Incubate the cassette in a 5% CO₂ incubator at 37° for 1–2 h to equilibrate the pH before proceeding to transplantation.
 - g. For transport, use a small, airtight, sterile container equilibrated in the incubator with the dishes of xenografts to maintain CO₂ before transplantation.
8. Transplanting xenografts into *nu/nu* mice:
- a. Anesthetize male *nu/nu* athymic mice (Harlan, Indianapolis, IN) by intraperitoneal injection of ketamine (80 mg/kg) and xylazine (10 mg/kg) in sterile PBS.
 - b. Place the mice on a sterile drape after they are anesthetized. Clean the surgical incision sites with povidone-iodine followed by 70% ethanol. Make four incisions as shown in Fig. 4D (center panel), two small incisions on the neck of the mouse (~0.16 cm) with just enough width to pass the tubing, and two larger incisions on the flanks of the mouse (~1.0 cm).
 - c. Separate the skin from the muscle by blunt dissection and place the xenograft cassette subcutaneously by tunneling the distal end of cassette (tubing *c*) under the skin toward the tail of the mouse, and the proximal end of the cassette (tubing *a*) out of the small incision behind the neck, using forceps to guide the xenograft tubing (Fig. 4D).
 - d. After transplantation, use two or three staples to close each of the largest incisions and an additional staple to anchor each xenograft to the skin at the loop of tubing *c*. No staples are necessary for the exit port of tubing *a*. Then transfer the mouse to a sterile cage and monitor until it is awake.
 - e. Irrigate the xenografts weekly with 1 ml of Ham's F12 medium, using a Surflo winged infusion set (Fisher Scientific) with an 0.75-in., 21-gauge needle to remove excess secretions in the first 3 weeks after transplantation. Beginning 3 weeks posttransplantation, irrigate the xenografts twice per week. Stratified xenograft epithelium usually is reconstituted by 4–6 weeks after transplantation (sEngelhardt *et al.*, 1991, 1992, 1995).

Medium A

Modified Eagle's medium (MEM), Ca²⁺ and Mg²⁺ free

Penicillin (100 U/ml)

Streptomycin (100 µg/ml)
Tobramycin (80 µg/ml)
Ceftazidime (100 µg/ml)
Primaxin (100 µg/ml)
Amphotericin B (5.0 µg/ml)
Deoxyribonuclease I [DNase I (100 µg/ml), cat. no. DN25; Sigma]
Pronase (1.5 mg/ml) (cat. no. 1459643; Roche)
Make fresh and keep at 4°.

Medium B

5% FBS in Dulbecco's modified Eagle's medium (DMEM)–Ham's F12 (1:1 ratio)
1% MEM nonessential amino acids solution
Penicillin (100 U/ml)
Streptomycin (100 µg/ml)
Tobramycin (80 µg/ml)
Ceftazidime (100 µg/ml)
Primaxin (100 µg/ml)
Amphotericin B (5.0 µg/ml)

Medium C

DMEM–Ham's F12 (1:1 ratio)
15 mM HEPES
3.6 mM Na₂CO₃
Penicillin (100 U/ml)
Streptomycin (100 µg/ml)
Tobramycin (80 µg/ml)
Ceftazidime (100 µg/ml)
Primaxin (100 µg/ml)
Amphotericin B (0.25 µg/ml)
Clonetics BEGM SingleQuots kit (Cambrex Bio Science Walkersville, Walkersville, MD)
Combine all components of the SingleQuots kit with the preceding ingredients, resulting in the following final concentrations: insulin (10.0 µg/ml), cholera toxin (1.0 µg/ml), bovine

pituitary extract (40 µg/ml), human epidermal growth factor (hEGF, 1.0 µg/ml), epinephrine (1.0 µg/ml), transferrin (20.0 µg/ml), and retinoic acid (0.0001 µg/ml).

Medium D

BEGM (bronchial epithelial growth medium)

BEGM SingleQuots kit Penicillin (50 U/ml)

Streptomycin (50 µg/ml)

Tobramycin (40 µg/ml)

Ceftazidime (50 µg/ml)

Primaxin (50 µg/ml)

Trypsin Inhibitor Buffer

Trypsin inhibitor type I-S: from soybean (cat. no. T-6522; Sigma)

Ham's F12 medium

Combine trypsin inhibitor with Ham's F12 medium at 1 mg/ml. Filter through a 0.22-µm pore size filter, aliquot, and store at −20°.

***In Vitro* Colony-Forming Efficiency Assay to Characterize Stem Cell Populations in Conducting Airway Epithelium**

Overview

Adult airway stem cells have the ability to generate highly proliferative TA cells by asymmetric cell division. This feature allows stem cells to undergo multiple rounds of clonal proliferation in the setting of injury/repair that can be measured using an *in vitro* colony-forming efficiency (CFE) assay in air–liquid interface (ALI) airway epithelial cell cultures (Randell, 1992) (Fig. 5). This CFE assay is similar to that used in the tracheal xenograft model with recombinant retroviral markers to assess clonal expansion of airway stem/progenitor cells (Engelhardt *et al.*, 1995; Zepeda *et al.*, 1995), but with the added advantage of being easier to carry out. Disadvantages of this *in vitro* CFE assay include less efficient differentiation of ciliated and goblet cell types in the ALI culture as compared with the tracheal xenograft method. Hence, this *in vitro* CFE assay has been used primarily to assess the proliferative potential of stem/progenitor cell populations, and less frequently to study progenitor–progeny relationships in the airway epithelium.

Using this ALI culture system, Schoch and colleagues evaluated the clonal growth potential of murine tracheal epithelial cells, using CFE as an index. They defined a subset of basal cells in mouse tracheal epithelium with the capacity to generate large colonies in ALI culture, suggesting they are derived from stem or TA cells (Schoch *et al.*, 2004). They diluted a single-cell suspension of mouse tracheal epithelial cells derived from β-galactosidase-expressing ROSA26 mice into non-ROSA26 tracheal epithelial cells and placed this mixture into the ALI culture model. After 3 weeks of ALI culture, 5-bromo-4-chloro-3-indolyl-β-D-galactopyranoside (X-Gal) staining revealed ROSA26 LacZ-positive colonies within the polarized airway epithelium. They observed that 1.7% of the tracheal epithelial cells formed colonies of varying size, with 0.1% of the clones forming large colonies (Fig. 5C–E). This subset of cells with larger proliferative potential was suggested as a possible stem or early TA cell population. Previous studies from the same group have demonstrated that high keratin 5

(K5) promoter activity exists in specific niches in the mouse trachea that correspond to the location of BrdU-labeled LRCs, thought to be stem cells (Borthwick *et al.*, 2001). Transgenic mice harboring a K5 promoter-driven EGFP transgene expressed EGFP in most basal cells of the body, including a subset of tracheal basal cells apparently localized to positions similar to previously identified stem cell niches. Further studies using the same mouse model and FACS-facilitated purification of K5 promoter-driven EGFP-positive tracheal epithelial cells revealed an overall colony-forming efficiency 4.5-fold greater than that of EGFP-negative cells. In addition, these K5 promoter-driven EGFP-positive tracheal epithelial cells retained the ability to generate 12-fold larger colonies than EGFP-negative cells, consistent with the notion that K5-positive basal cells contain a proximal airway stem/progenitor cell subpopulation (Borthwick *et al.*, 2001; Schoch *et al.*, 2004). The following protocol presents the details of this CFE assay, using the mouse ALI culture model, and is based on the studies by Schoch and colleagues (2004).

***In Vitro* Mouse Tracheal Epithelial Cell Colony-Forming Efficiency Assay Protocol**

1. Isolation of mouse tracheal epithelial cells: Four- to 8-week-old ROSA26 mice (B6.129S4-*Gt(ROSA)26*; Jackson Laboratory, Bar Harbor, ME) and non-ROSA26 mice (Harlan, Indianapolis, IN) are killed by CO₂ asphyxiation and pinned to a Styrofoam bed.
 - a. Excise the tracheas from the pharynx to the bronchial main branches under sterile conditions and place the tracheas in ice-cold Ham's F12 medium containing penicillin (100 U/ml), streptomycin (100 µg/ml), and amphotericin B (Fungizone, 1.0 µg/ml) (Ham's F12 pen–strep–Fungizone). Isolate ROSA26 and non-ROSA26 tracheal epithelial cells separately.
 - b. Open the tracheas longitudinally after the muscle and fat have been removed from the tracheas. Incubate the tracheas in the preceding Ham's F12 pen–strep–Fungizone medium containing Pronase (1.5 mg/ml; Roche Molecular Biochemicals, Indianapolis, IN) for 18–24 h at 4°, occasionally inverting the tube several times during dissociation (Davidson *et al.*, 2000; Liu *et al.*, 2006; You *et al.*, 2002).
 - c. Add FBS to the digestion tube at a final concentration of 10%. Dissociate the tracheal epithelial cells by inverting the tube 10–20 times and transfer the cell suspension to a new tube. Wash the tracheas with Ham's F12 pen–strep–Fungizone medium twice and pool the cell suspensions. Collect the cells by centrifugation at 500g for 10 min at 4°.
 - d. Resuspend the cell pellets in Ham's F12 medium containing penicillin (100 U/ml), streptomycin (100 µg/ml), Fungizone (1.0 µg/ml), crude pancreatic DNase I (0.5 mg/ml) (Sigma-Aldrich, St. Louis, MO), and bovine serum albumin (BSA, 10 mg/ml), and incubate them on ice for 5 min. Wash the cells with Ham's F12 pen–strep–Fungizone medium and centrifuge the cells at 500g for 5 min at 4°.
2. The cell pellet is then resuspended in Ham's F12 pen–strep–Fungizone medium with 10% FBS and cells are incubated in tissue culture plates (Primaria; BD Biosciences Discovery Labware) for 2 h in 5% CO₂ at 37° to adhere fibroblasts. Nonadherent epithelial cells are collected by centrifugation and resuspended in 5% FBS–BEGM medium (Cambrex Bio Science Walkersville). Total cell yields are then counted with a hemacytometer. BEGM medium is made by adding reagents from one BEGM

SingleQuots kit to 500 ml of 50% DMEM–50% Ham's F12 medium supplemented with 1% pen–strep, Fungizone (0.25 µg/ml), 15 mM HEPES, and 3.6 mM Na₂CO₃.

3. Preparation of ALI culture membranes: Primary airway epithelial cells will not attach to an ALI culture membrane support that is not coated with collagen. Use the following protocol to prepare ALI culture membranes.
 - a. Mix 30 mg of human placental collagen type IV (cat. no. C-7521; Sigma) with 50 ml of deionized water and 100 µl of glacial acetic acid in a 100-ml sterile beaker.
 - b. Cover the holding beaker with Parafilm and stir moderately at 37° until collagen strands are dissolved (~15–30 min).
 - c. Dilute the collagen stock 1:10 with deionized sterile water and filter sterilize the solution through a 0.22-µm pore size membrane. The diluted sterile collagen stock (60 µg/ml) is the working solution for coating the plastic and membrane surfaces.
 - d. Insert 12-mm-diameter Millicell-PCF membrane inserts (cat. no. PIHP 01250; Millipore, Bedford, MA) in a 24-well plate and apply 0.3–0.5 ml of collagen working solution on the apical surface of the inserts.
 - e. Collagen coat the surface for a minimum of 18 h at room temperature. Remove the liquid collagen from the surface and air dry the membrane surface in a laminar flow hood. Once dried, rinse both sides of the membrane support three times with sterile PBS or DMEM to remove all traces of free collagen. Redry the membrane inserts in a laminar flow hood and store at 4° (they will be stable for several months).
4. Seeding cells on the membrane:
 - a. Dilute an experimentally enriched subpopulation of ROSA26 tracheal epithelial cells into non-ROSA26 cells at an appropriate ratio. The ratio must be determined empirically to allow for outgrowth of isolated clones and will depend on the proliferative index of the cells being analyzed. Alternatively, one can also use retrovirally tagged cells expressing a detectable histochemical marker (i.e., EGFP, alkaline phosphatase, or β-galactosidase) for seeding onto membranes.
 - b. Seed a total of $1\text{--}2.5 \times 10^5$ cells/cm² in a 0.2-ml volume of 5% FBS–BEGM on the surface of a Millicell insert (about $1\text{--}1.5 \times 10^5$ cells per 12-mm insert). The volume should be sufficient to ensure a uniform distribution of cell settling on the membrane surface.
 - c. Apply 0.3–0.5 ml of 5% FBS–BEGM in the basal compartment of the insert to immerse the membrane without floating the insert; make sure that the membranes are level to ensure uniform cell attachment during the first 12 h.
 - d. Leave the plates containing the inserts undisturbed for a minimum of 18–24 h in an 8–9% CO₂ incubator at 37°. The higher CO₂ level has been shown to increase successful achievement of confluence (Karp *et al.*, 2002).
5. Establishment of ALI culture:
 - a. The day after seeding, wash the membranes with prewarmed PBS to remove unattached cells and refeed them with fresh 5% FBS–BEGM. The medium is changed every 2 days until the transmembrane resistance (R_t) is greater than $\sim 1000 \Omega/\text{cm}^2$ (this usually occurs 2–5 days after seeding). The electrical

resistance across the membrane is measured with a Millicell-ERS system ohm meter (cat. no. MERS 000 01; Millipore).

- b. Remove the medium inside the insert (apical) and refeed the outside (basal) chamber with BEGM without FBS to establish an ALI. Cells are refeed with BEGM every 2–3 days. A polarized and differentiated airway epithelium will form ~10–15 days after moving the epithelium to an ALI.
6. X-Gal staining and evaluation: About 3 weeks (21 days) after seeding, the mouse ALI cultures should be ready for evaluation of CFE or other assays as appropriate. Depending on the purposes of the study, the membrane with polarized mouse tracheal epithelial cells can be either freshly embedded in O.C.T. compound for frozen sectioning, or directly fixed for *en face* immunostaining or X-Gal staining as previously described (Schoch *et al.*, 2004). Figure 5C–E depicts X-Gal-stained clonal expansion of ROSA26 mouse tracheal epithelial progenitor cells in this model.

Models to Study Stem/Progenitor Cells of Airway Submucosal Glands

Overview

Airway submucosal glands (SMGs) are major secretory structures that reside in the cartilaginous airways of many mammalian species (Liu *et al.*, 2004). SMGs play important roles in both innate immunity of the lung (Dajani *et al.*, 2005) and cell biology of the proximal airways (Borthwick *et al.*, 2001; Engelhardt, 2001; Engelhardt *et al.*, 1995). As discussed in more detail in a previous section, SMGs serve as a protective niche for surface airway epithelial stem cells (Borthwick *et al.*, 2001; Engelhardt, 2001). Furthermore, the pluripotent progenitor cells that exist in the human tracheobronchial surface airway epithelium (i.e., cells with the capacity to differentiate into ciliated, secretory, intermediate, and basal cells) also have a developmental capacity for SMGs (Engelhardt *et al.*, 1995). Given that adult surface airway epithelial stem/progenitor cells have the capacity to develop SMGs, the biology that controls the morphogenesis of these structures in the airway is highly relevant to defining stem cell characteristics in the airway.

The development of three-dimensional culture substrates has enabled the creation of model systems to study certain features of SMG development, using a system that is far less complex than tracheal xenografts. Epithelial invasion of the extracellular matrix (ECM) is an important aspect of lung development and SMG morphogenesis. Infeld and colleagues initially developed an *in vitro* model for early airway gland development by culturing human tracheobronchial epithelial cells on a floating collagen gel substrate that contained fetal lung fibroblasts (Infeld *et al.*, 1993). Similar studies by other groups have also demonstrated the capacity of human airway epithelial cells to invade a collagen gel matrix (Emura *et al.*, 1996; Jacquot *et al.*, 1994). Together with technical advances in the isolation and culture of mouse tracheal epithelial cells, we have adapted these protocols to evaluate bud formation, tubulogenesis, and branching morphogenesis of airway epithelia, using the collagen gel matrix model (Fig. 6). Although such models cannot reproduce the native cellular diversity found in SMGs, they will be useful in defining fundamental epithelial–mesenchymal interactions required for airway gland morphogenesis.

In Vitro Invasion and Tubulogenesis Assay Using Mouse Tracheal Epithelial Cells

1. Collagen gel contraction:
 - a. Primary mouse embryonic fibroblast cells (PMEFs): PMEFs are generated according to a previously described general protocol (Wassarman and

DePamphilis, 1993). PMEFs generated from ICR or C57/BL6 mice can be used for this model with no differences in experimental outcome. Culture PMEFs in DMEM supplemented with 10% FBS. It is worth noting that early passages of PMEFs may give better results in gel contraction than later passage cells. However, for the most consistent results the same passages of PMEFs should always be used to generate contracted collagen cells.

- b. PMEFs are harvested by trypsinization, and the total cell number is quantified with a hemacytometer counting chamber.
 - c. To generate the collagen gel matrix support on which epithelial cells are seeded, add the following reagents, in order, for each 60-mm dish at room temperature: 3.5 ml of fibroblast culture medium, 1.8 ml of FBS, 0.3 ml of 0.1 N NaOH, $3-5 \times 10^5$ fibroblasts in 1.0 ml of medium, and 9.5 mg of rat tail collagen type I (BD Biosciences Discovery Labware) in 2.7 ml of H₂O to a final volume of 9.3 ml. Mix the solution gently by inverting the tube. The mixture should be light pink (if not, add 0.1 N NaOH to adjust the pH to turn the phenol red pink).
 - d. Apply ~9 ml of collagen gel to a 60-mm dish (or ~5 ml for a 35-mm dish) and incubate the gel at room temperature for 10 min before moving the dish into a 5% CO₂ incubator at 37° for 1 h. Add a small amount of fibroblast culture medium to the gel and run a sterile pipette around the outside rim of the gel to allow contraction. Continue incubation in 5% CO₂ at 37° and feed twice per week with fibroblast culture medium. The gel will be fully contracted in 2–4 weeks.
2. Isolation of mouse tracheal epithelial cells: Use the same protocol described in the previous section for *in vitro* CFE assays.
 3. Mouse tracheal epithelial cell seeding:
 - a. After gel contraction, fresh isolated or cultured mouse airway cells are resuspended in 5% FBS–BEGM at a density of $3-5 \times 10^7$ cells/ml. BEGM is made by adding reagents from one BEGM SingleQuot kit to 500 ml of 50% DMEM–50% Ham's F12 medium supplemented with 1% pen–strep, Fungizone (0.25 µg/ml), 15 mM HEPES, and 3.6 mM Na₂CO₃.
 - b. Remove the medium from the fibroblast gel and apply 10 µl of the cell solution to the top of each gel while attempting to evenly spread the cells across the gel. Incubate the gels in 5% CO₂ at 37° for 30 min before adding a small amount of 5% FBS–BEGM around the edge of the gel. Continue incubation in 5% CO₂ at 37° overnight to allow adequate cell adhesion. Take care not to add so much medium that it covers the top of the gel and washes the cells off the surface.
 - c. The following morning, add enough 5% FBS–BEGM to cover the top of the gel and let the gel sit for 3 days before placing the gels into 30-mm diameter Millicell-PCF membrane inserts (cat. no. PIHP 03012; Millipore) in a 6-well dish to establish an air–liquid interface (ALI). After the ALI is established, feed the cells from the bottom of the inserts with BEGM (without FBS). The apical side of the gel should be exposed to air. Bud formation and tubulogenesis of airway epithelial cells can be evaluated 1–2 weeks after establishment of the ALI (Fig. 6).

Acknowledgements

The authors gratefully acknowledge NIDDK research funding in the area of this review (DK47967) and the editorial assistance of Ms. Leah Williams.

References

- Asmundsson T, Kilburn KH, McKenzie WN. Injury and metaplasia of airway cells due to SO₂. *Lab Invest* 1973;29:41–53. [PubMed: 4728356]
- Berns A. Stem cells for lung cancer? *Cell* 2005;121:811–813. [PubMed: 15960966]
- Bigby TD, Allen D, Leslie CG, Henson PM, Cherniack RM. Bleomycin- induced lung injury in the rabbit: Analysis and correlation of bronchoalveolar lavage, morphometrics, and fibroblast stimulating activity. *Am Rev Respir Dis* 1985;132:590–595. [PubMed: 2412475]
- Bishop AE. Pulmonary epithelial stem cells. *Cell Prolif* 2004;37:89–96. [PubMed: 14871239]
- Boers JE, Ambergen AW, Thunnissen FB. Number and proliferation of basal and parabasal cells in normal human airway epithelium. *Am J Respir Crit Care Med* 1998;157:2000–2006. [PubMed: 9620938]
- Borges M, Linnoila RI, van de Velde HJ, Chen H, Nelkin BD, Mabry M, Baylin SB, Ball DW. An achaete-scute homologue essential for neuroendocrine differentiation in the lung. *Nature* 1997;386:852–855. [PubMed: 9126746]
- Borthwick DW, Shahbazian M, Krantz QT, Dorin JR, Randell SH. Evidence for stem-cell niches in the tracheal epithelium. *Am J Respir Cell Mol Biol* 2001;24:662–670. [PubMed: 11415930]
- Buckpitt A, Buonarati M, Avey LB, Chang AM, Morin D, Plopper CG. Relationship of cytochrome P450 activity to Clara cell cytotoxicity. II Comparison of stereoselectivity of naphthalene epoxidation in lung and nasal mucosa of mouse, hamster, rat and rhesus monkey. *J Pharmacol Exp Ther* 1992;261:364–372. [PubMed: 1560380]
- Castleman WL, Dungworth DL, Tyler WS. Intrapulmonary airway morphology in three species of monkeys: A correlated scanning and transmission electron microscopic study. *Am J Anat* 1975;142:107–121. [PubMed: 804810]
- Choi HK, Finkbeiner WE, Widdicombe JH. A comparative study of mammalian tracheal mucous glands. *J Anat* 2000;197:361–372. [PubMed: 11117623]
- Dajani R, Zhang Y, Taft PJ, Travis SM, Starner TD, Olsen A, Zabner J, Welsh MJ, Engelhardt JF. Lysozyme secretion by submucosal glands protects the airway from bacterial infection. *Am J Respir Cell Mol Biol* 2005;32:548–552. [PubMed: 15746432]
- Davidson DJ, Kilanowski FM, Randell SH, Sheppard DN, Dorin JR. A primary culture model of differentiated murine tracheal epithelium. *Am J Physiol Lung Cell Mol Physiol* 2000;279:L766–L778. [PubMed: 11000138]
- Duan D, Sehgal A, Yao J, Engelhardt JF. Lef1 transcription factor expression defines airway progenitor cell targets for *in utero* gene therapy of submucosal gland in cystic fibrosis. *Am J Respir Cell Mol Biol* 1998;18:750–758. [PubMed: 9618379]
- Dubaybo BA, Crowell LA, Thet LA. Changes in tissue fibronectin in elastase induced lung injury. *Cell Biol Int Rep* 1991;15:675–686. [PubMed: 1742801]
- Emura M. Stem cells of the respiratory epithelium and their *in vitro* cultivation. *In Vitro Cell Dev Biol Anim* 1997;33:3–14. [PubMed: 9028828]
- Emura M. Stem cells of the respiratory tract. *Paediatr Respir Rev* 2002;3:36–40. [PubMed: 12065180]
- Emura M, Ochiai A, Hirohashi S. *In vitro* reconstituted tissue as an alternative to human respiratory tract. *Toxicol Lett* 1996;88:81–84. [PubMed: 8920720]
- Engelhardt JF. Stem cell niches in the mouse airway. *Am J Respir Cell Mol Biol* 2001;24:649–652. [PubMed: 11415927]
- Engelhardt JF, Allen ED, Wilson JM. Reconstitution of tracheal grafts with a genetically modified epithelium. *Proc Natl Acad Sci USA* 1991;88:11192–11196. [PubMed: 1763032]
- Engelhardt JF, Yankaskas JR, Wilson JM. *In vivo* retroviral gene transfer into human bronchial epithelia of xenografts. *J Clin Invest* 1992;90:2598–2607. [PubMed: 1281842]

- Engelhardt JF, Schlossberg H, Yankaskas JR, Dudus L. Progenitor cells of the adult human airway involved in submucosal gland development. *Development* 1995;121:2031–2046. [PubMed: 7635050]
- Evans MJ, Cabral-Anderson LJ, Dekker NP, Freeman G. The effects of dietary antioxidants on NO₂-induced injury to type 1 alveolar cells. *Chest* 1981;80:5–8. [PubMed: 7249739]
- Filali M, Zhang Y, Ritchie TC, Engelhardt JF. Xenograft model of the CF airway. *Methods Mol Med* 2002;70:537–550. [PubMed: 11917549]
- Giangreco A, Reynolds SD, Stripp BR. Terminal bronchioles harbor a unique airway stem cell population that localizes to the bronchoalveolar duct junction. *Am J Pathol* 2002;161:173–182. [PubMed: 12107102]
- Giangreco A, Shen H, Reynolds SD, Stripp BR. Molecular phenotype of airway side population cells. *Am J Physiol Lung Cell Mol Physiol* 2004;286:L624–L630. [PubMed: 12909587]
- Griffiths MJ, Bonnet D, Janes SM. Stem cells of the alveolar epithelium. *Lancet* 2005;366:249–260. [PubMed: 16023517]
- Grove JE, Lutzko C, Priller J, Henegariu O, Theise ND, Kohn DB, Krause DS. Marrow-derived cells as vehicles for delivery of gene therapy to pulmonary epithelium. *Am J Respir Cell Mol Biol* 2002;27:645–651. [PubMed: 12444022]
- Hansell MM, Moretti RL. Ultrastructure of the mouse tracheal epithelium. *J Morphol* 1969;128:159–169. [PubMed: 5793184]
- Hay J, Shahzeidi S, Laurent G. Mechanisms of bleomycin-induced lung damage. *Arch Toxicol* 1991;65:81–94. [PubMed: 1711838]
- Hong KU, Reynolds SD, Giangreco A, Hurley CM, Stripp BR. Clara cell secretory protein-expressing cells of the airway neuroepithelial body microenvironment include a label-retaining subset and are critical for epithelial renewal after progenitor cell depletion. *Am J Respir Cell Mol Biol* 2001;24:671–681. [PubMed: 11415931]
- Hong KU, Reynolds SD, Watkins S, Fuchs E, Stripp BR. Basal cells are a multipotent progenitor capable of renewing the bronchial epithelium. *Am J Pathol* 2004a;164:577–588. [PubMed: 14742263]
- Hong KU, Reynolds SD, Watkins S, Fuchs E, Stripp BR. *In vivo* differentiation potential of tracheal basal cells: Evidence for multipotent and unipotent subpopulations. *Am J Physiol Lung Cell Mol Physiol* 2004b;286:L643–L649. [PubMed: 12871857]
- Hook GE, Brody AR, Cameron GS, Jetten AM, Gilmore LB, Nettesheim P. Repopulation of denuded tracheas by Clara cells isolated from the lungs of rabbits. *Exp Lung Res* 1987;12:311–329. [PubMed: 3582283]
- Inayama Y, Hook GE, Brody AR, Cameron GS, Jetten AM, Gilmore LB, Gray T, Nettesheim P. The differentiation potential of tracheal basal cells. *Lab Invest* 1988;58:706–717. [PubMed: 3379917]
- Infeld MD, Brennan JA, Davis PB. Human fetal lung fibroblasts promote invasion of extracellular matrix by normal human tracheobronchial epithelial cells *in vitro*: A model of early airway gland development. *Am J Respir Cell Mol Biol* 1993;8:69–76. [PubMed: 8417759]
- Izbicki G, Segel MJ, Christensen TG, Conner MW, Breuer R. Time course of bleomycin-induced lung fibrosis. *Int J Exp Pathol* 2002;83:111–119. [PubMed: 12383190]
- Jacquot J, Spilmont C, Burlet H, Fuchey C, Buisson AC, Tournier JM, Gaillard D, Puchelle E. Glandular-like morphogenesis and secretory activity of human tracheal gland cells in a three-dimensional collagen gel matrix. *J Cell Physiol* 1994;161:407–418. [PubMed: 7962124]
- Januszkiewicz AJ, Mayorga MA. Nitrogen dioxide-induced acute lung injury in sheep. *Toxicology* 1994;89:279–300. [PubMed: 8023332]
- Jeffery PK. Morphologic features of airway surface epithelial cells and glands. *Am Rev Respir Dis* 1983;128:S14–S20. [PubMed: 6881701]
- Karp PH, Moninger TO, Weber SP, Nesselhauf TS, Launsbach JL, Zabner J, Welsh MJ. An *in vitro* model of differentiated human airway epithelia: Methods for establishing primary cultures. *Methods Mol Biol* 2002;188:115–137. [PubMed: 11987537]
- Kim CF, Jackson EL, Woolfenden AE, Lawrence S, Babar I, Vogel S, Crowley D, Bronson RT, Jacks T. Identification of bronchioalveolar stem cells in normal lung and lung cancer. *Cell* 2005;121:823–835. [PubMed: 15960971]

- Lamb D, Reid L. Mitotic rates, goblet cell increase and histochemical changes in mucus in rat bronchial epithelium during exposure to sulphur dioxide. *J Pathol Bacteriol* 1968;96:97–111. [PubMed: 5667859]
- Langley-Evans SC, Phillips GJ, Jackson AA. Sulphur dioxide: A potent glutathione depleting agent. *Comp Biochem Physiol C Pharmacol Toxicol Endocrinol* 1996;114:89–98. [PubMed: 8760604]
- Liu X, Driskell RR, Engelhardt JF. Airway glandular development and stem cells. *Curr Top Dev Biol* 2004;64:33–56. [PubMed: 15563943]
- Liu X, Yan Z, Luo M, Engelhardt JF. Species-specific differences in mouse and human airway epithelial biology of rAAV transduction. *Am J Respir Cell Mol Biol* 2006;34:56–64. [PubMed: 16195538]
- Mason RJ, Williams MC, Moses HL, Mohla S, Berberich MA. Stem cells in lung development, disease, and therapy. *Am J Respir Cell Mol Biol* 1997;16:355–363. [PubMed: 9115744]
- Meulenbelt J, Dormans JA, Marra M, Rombout PJ, Sangster B. Rat model to investigate the treatment of acute nitrogen dioxide intoxication. *Hum Exp Toxicol* 1992a;11:179–187. [PubMed: 1352113]
- Meulenbelt J, van Bree L, Dormans JA, Boink AB, Sangster B. Biochemical and histological alterations in rats after acute nitrogen dioxide intoxication. *Hum Exp Toxicol* 1992b;11:189–200. [PubMed: 1352114]
- Meulenbelt J, van Bree L, Dormans JA, Boink AB, Sangster B. Development of a rabbit model to investigate the effects of acute nitrogen dioxide intoxication. *Hum Exp Toxicol* 1994;13:749–758. [PubMed: 7857694]
- Neuringer IP, Randell SH. Stem cells and repair of lung injuries. *Respir Res* 2004;5:6. [PubMed: 15285789]
- Otto WR. Lung epithelial stem cells. *J Pathol* 2002;197:527–535. [PubMed: 12115868]
- Pack RJ, Al-Ugaily LH, Morris G, Widdicombe JG. The distribution and structure of cells in the tracheal epithelium of the mouse. *Cell Tissue Res* 1980;208:65–84. [PubMed: 6248229]
- Peake JL, Reynolds SD, Stripp BR, Stephens KE, Pinkerton KE. Alteration of pulmonary neuroendocrine cells during epithelial repair of naphthalene- induced airway injury. *Am J Pathol* 2000;156:279–286. [PubMed: 10623676]
- Plopper CG, Hill LH, Mariassy AT. Ultrastructure of the nonciliated bronchiolar epithelial (Clara) cell of mammalian lung. III A study of man with comparison of 15 mammalian species. *Exp Lung Res* 1980;1:171–180. [PubMed: 7227344]
- Plopper CG, Mariassy AT, Wilson DW, Alley JL, Nishio SJ, Nettesheim P. Comparison of nonciliated tracheal epithelial cells in six mammalian species: Ultrastructure and population densities. *Exp Lung Res* 1983;5:281–294. [PubMed: 6662075]
- Plopper CG, Weir AJ, Nishio SJ, Cranz DL, St George JA. Tracheal submucosal gland development in the rhesus monkey, *Macaca mulatta*: Ultrastructure and histochemistry. *Anat Embryol (Berl)* 1986;174:167–178. [PubMed: 3740452]
- Presente A, Sehgal A, Dudus L, Engelhardt JF. Differentially regulated epithelial expression of an Eph family tyrosine kinase (fHek2) during tracheal surface airway and submucosal gland development. *Am J Respir Cell Mol Biol* 1997;16:53–61. [PubMed: 8998079]
- Randell SH. Progenitor–progeny relationships in airway epithelium. *Chest* 1992;101:11S–16S. [PubMed: 1541185]
- Ranga V, Kleiner J. A quantitative study of ciliary injury in the small airways of mice: The effects of nitrogen dioxide. *Exp Lung Res* 1981;2:49–55. [PubMed: 7346269]
- Reddy R, Buckley S, Doerken M, Barsky L, Weinberg K, Anderson KD, Warburton D, Driscoll B. Isolation of a putative progenitor subpopulation of alveolar epithelial type 2 cells. *Am J Physiol Lung Cell Mol Physiol* 2004;286:L658–L667. [PubMed: 12922980]
- Reynolds SD, Giangreco A, Power JH, Stripp BR. Neuroepithelial bodies of pulmonary airways serve as a reservoir of progenitor cells capable of epithelial regeneration. *Am J Pathol* 2000a;156:269–278. [PubMed: 10623675]
- Reynolds SD, Hong KU, Giangreco A, Mango GW, Guron C, Morimoto Y, Stripp BR. Conditional clara cell ablation reveals a self-renewing progenitor function of pulmonary neuroendocrine cells. *Am J Physiol Lung Cell Mol Physiol* 2000b;278:L1256–L1263. [PubMed: 10835332]
- Reynolds SD, Giangreco A, Hong KU, McGrath KE, Ortiz LA, Stripp BR. Airway injury in lung disease pathophysiology: Selective depletion of airway stem and progenitor cell pools potentiates lung

- inflammation and alveolar dysfunction. *Am J Physiol Lung Cell Mol Physiol* 2004;287:L1256–L1265. [PubMed: 15298853]
- Schoch KG, Lori A, Burns KA, Eldred T, Olsen JC, Randell SH. A subset of mouse tracheal epithelial basal cells generates large colonies *in vitro*. *Am J Physiol Lung Cell Mol Physiol* 2004;286:L631–L642. [PubMed: 12959927]
- Sehgal A, Presente A, Engelhardt JF. Developmental expression patterns of CFTR in ferret tracheal surface airway and submucosal gland epithelia. *Am J Respir Cell Mol Biol* 1996;15:122–131. [PubMed: 8679216]
- Smith LJ. Hyperoxic lung injury: Biochemical, cellular, and morphologic characterization in the mouse. *J Lab Clin Med* 1985;106:269–278. [PubMed: 2993457]
- Souma T. The distribution and surface ultrastructure of airway epithelial cells in the rat lung: A scanning electron microscopic study. *Arch Histol Jpn* 1987;50:419–436. [PubMed: 3439845]
- Summer R, Kotton DN, Sun X, Ma B, Fitzsimmons K, Fine A. Side population cells and Bcrp1 expression in lung. *Am J Physiol Lung Cell Mol Physiol* 2003;285:L97–L104. [PubMed: 12626330]
- Suzuki M, Machida M, Adachi K, Otake K, Sugimoto T, Hayashi M, Awazu S. Histopathological study of the effects of a single intratracheal instillation of surface active agents on lung in rats. *J Toxicol Sci* 2000;25:49–55. [PubMed: 10736790]
- Ten Have-Opbroek AA, Plopper CG. Morphogenetic and functional activity of type II cells in early fetal rhesus monkey lungs: A comparison between primates and rodents. *Anat Rec* 1992;234:93–104. [PubMed: 1416101]
- Theise ND, Henegariu O, Grove J, Jagirdar J, Kao PN, Crawford JM, Badve S, Saxena R, Krause DS. Radiation pneumonitis in mice: A severe injury model for pneumocyte engraftment from bone marrow. *Exp Hematol* 2002;30:1333–1338. [PubMed: 12423687]
- Vacanti MP, Roy A, Cortiella J, Bonassar L, Vacanti CA. Identification and initial characterization of spore-like cells in adult mammals. *J Cell Biochem* 2001;80:455–460. [PubMed: 11135375]
- Van Winkle LS, Johnson ZA, Nishio SJ, Brown CD, Plopper CG. Early events in naphthalene-induced acute Clara cell toxicity: Comparison of membrane permeability and ultrastructure. *Am J Respir Cell Mol Biol* 1999;21:44–53. [PubMed: 10385592]
- Wassarman, PM.; DePamphilis, ML. *Guide to Techniques in Mouse Development*. Academic Press; San Diego, CA: 1993.
- Watt FM, Hogan BL. Out of Eden: Stem cells and their niches. *Science* 2000;287:1427–1430. [PubMed: 10688781]
- West JA, Pakehham G, Morin D, Fleschner CA, Buckpitt AR, Plopper CG. Inhaled naphthalene causes dose dependent Clara cell cytotoxicity in mice but not in rats. *Toxicol Appl Pharmacol* 2001;173:114–119. [PubMed: 11384213]
- Widdicombe JH, Chen LL, Sporer H, Choi HK, Pecson IS, Bastacky SJ. Distribution of tracheal and laryngeal mucous glands in some rodents and the rabbit. *J Anat* 2001;198:207–221. [PubMed: 11273045]
- You Y, Richer EJ, Huang T, Brody SL. Growth and differentiation of mouse tracheal epithelial cells: Selection of a proliferative population. *Am J Physiol Lung Cell Mol Physiol* 2002;283:L1315–L1321. [PubMed: 12388377]
- Zepeda ML, Chinoy MR, Wilson JM. Characterization of stem cells in human airway capable of reconstituting a fully differentiated bronchial epithelium. *Somat Cell Mol Genet* 1995;21:61–73. [PubMed: 7541563]

FIG. 1.

Cellular diversity of epithelial cell types in the human adult lung. When discussing progenitor/stem cells in adult airways, one must always consider the diversity of cell phenotypes that exist in spatially distinct epithelia of the lung. Three main levels of conducting airways exist in the lung, including the trachea, bronchi, and bronchioles. Predominant cell types in the human pseudostratified, columnar tracheal, and bronchial epithelia include basal (B), intermediate (I), goblet (G), and ciliated (Ci) cells; less abundant nonciliated and neuroendocrine cells are not shown. Submucosal glands are also present only in the cartilaginous airways of the trachea and bronchi. Predominant cell types in the human columnar bronchiolar epithelia include Clara cells (Cl) and ciliated cells; less abundant neuroendocrine cells are not shown. Predominant cell types in the gas-exchanging alveolar airspaces include alveolar epithelial type II cells (AEC II), alveolar epithelial type I cells (AEC I), and capillary endothelial cells of the capillary networks (cn). Mt, mucous tubule; St, serous tubule; SAE, surface airway epithelium; Cd, collecting duct.

FIG. 2.

Naphthalene-induced mouse tracheal epithelial injury and the subacute proliferative response. Hematoxylin and eosin (H&E)-stained sections of proximal trachea 1 day (A) and 7 days (B) after intraperitoneal injection of naphthalene (275 mg/kg). Naphthalene treatment induced complete ablation of tracheal epithelium (arrows in A). By 7 days postinjury, a regenerated tracheal epithelium is observed (arrows in B). (C) BrdU labeling of mice during the first week of naphthalene injury and harvested at 28 days postinjury. Immunofluorescence detection of BrdU reveals high levels of incorporation in epithelia of submucosal glands and infrequent labeling of cells in proximal surface tracheal epithelium.

FIG. 3.

Polidocanol-induced injury of the mouse tracheal epithelium. Anesthetized mice are hung on a self-made rack by placing the upper jaw incisor teeth within a suture before placing a plastic catheter into the trachea as shown in (A) (*front view*) and (B) (*top view*). The rack is made with a >3-mm metal wire fixed on a wooden board with a screw. After intratracheal instillation of 20 μ l of 2% polidocanol, tracheas are harvested and sectioned. (C–E) H&E-stained sections from (C) control uninjured trachea, (D) injured trachea at 1 day, and (E) injured trachea at 14 days. Arrows mark injury to the surface airway epithelium at 1 day after polidocanol treatment in (D) as compared with injured controls (C) and regenerated epithelium (E). SAE, surface airway epithelium; SMG, submucosal gland.

FIG. 4.

Methods for generating tracheal xenografts to study clonal expansion. (A) Schematic methods for generating proximal airway epithelial xenograft models. Primary airway epithelial cells are cultured *in vitro* and may be infected with integrating recombinant viral vectors (lentivirus or retrovirus) before transplantation of epithelia into denuded rat tracheas. Fully differentiated epithelium is obtained by 4 weeks posttransplantation (B, basal cells; Ci, ciliated cells; I, intermediate cells; G, goblet cells). (B) Colony-forming efficiency (CFE) of stem/progenitor cells can be evaluated by detecting virally expressed transgenes in the reconstituted surface airway epithelium (SAE) (arrows). In this example a recombinant retroviral vector expressing the β -galactosidase transgene was used and detected by X-Gal staining. (C) Schematic view

of various components of the xenograft cassette. The denuded rat trachea is connected to a sterile tubing cassette by a series of sutures as illustrated: *a*, 1-in. Silastic tubing (cat. no. 602-175; Dow Corning, Midland, MI); *b*, 3/4-in. Silastic tubing (cat. no. 602-175; Dow Corning); *c*, 1.75-in. Silastic tubing (cat. no. 602-175; Dow Corning); *d*, 1.25-in. Teflon tubing (cat. no. 9567-K10; Thomas Scientific, Swedesboro, NJ); *e* and *e'*, adapter (0.8-mm barb-to-barb connector; cat. no. 732-8300; Bio-Rad, Hercules, CA). A chrome wire plug (0.0035-in.-diameter Chromel A steel wire; Hoskins Manufacturing, Hamburg, MI) is also used. (D) Subcutaneous transplantation of the xenograft cassettes in *nu/nu* athymic mice. *Left*: A subcutaneous view of the transplanted xenograft cassette. *Middle*: The four incisions (arrows) made on the back of the recipient *nu/nu* mouse before transplantation. The xenograft cassette is guided subcutaneously with forceps, so that one port exits through the back of the neck and the other port exits through the main incision. Surgical staples are then used to close the main incisions (marked by open arrowheads) and one additional staple is used to anchor the xenograft tubing near the tail end of the mouse (solid arrowheads). *Right*: The resultant xenograft cassette 1 week postsurgery, when the proximal staples are removed. The staples marked by solid arrowheads in the middle panel are used to maintain the position of the cassette and to prevent subcutaneous migration (it is necessary to leave these staples in for at least 2 weeks).

FIG. 5.

In vitro air-liquid interface model of colony-forming efficiency (CFE) assays, using ROSA26 mouse tracheal epithelial cells. (A) Schematic view of the ALI culture model of primary airway epithelial cells growing on collagen-coated Millipore cell culture inserts housed in a 24-well tissue culture plate. (B) Scanning electron micrograph of fully differentiated mouse airway epithelial cultures (scale bar: 10 μ m). (C) *En face* micrograph of X-Gal-stained ALI cultures seeded with a 1:50 mixture of ROSA26 to non-ROSA26 mouse tracheal epithelial cells. X-Gal-positive colonies of various sizes are seen. (D) Higher magnification of an individual X-Gal-positive colony from (C). (E) H&E-stained section of a small X-Gal-positive colony from (C).

FIG. 6.

Mouse tracheal epithelial cell tubulogenesis assay. Bud formation and tubulogenesis are two early events important in submucosal gland development and can be studied with a collagen gel matrix cultured at the air–liquid interface. Depicted here are various stages of gland-like structures formed from mouse tracheal epithelium (*top*) and schematic representations of various stages of tubulogenesis characteristic of early submucosal gland morphogenesis (*bottom*).

Summary of Predominant Epithelial Cell Types and SMGs in the Conducting Airways Between Rodents and Primates

Species	Basal	Intermediate	Goblet	Serous	Clara	NCC	Ciliated	SMG	References
Mouse	T,B	—	— ^a	—	T,B,Br	—	T,B,Br	T	Hansell and Moretti, 1969; Pack <i>et al.</i> , 1980; Widdicombe <i>et al.</i> , 2001
Rat	T,B	T,B	— ^a	T,B	Br	T,B	T,B,Br	T	Plopper <i>et al.</i> , 1983; Souma, 1987; Widdicombe <i>et al.</i> , 2001
Monkey	T,B	T,B	T,B	NA	NA	T,B	T,B,Br	T,B	Castleman <i>et al.</i> , 1975; Plopper <i>et al.</i> , 1983; Plopper <i>et al.</i> , 1986
Human	T,B	T,B	T,B	—	Br	T,B	T,B,Br	T,B	Jeffery, 1983; Plopper <i>et al.</i> , 1980

^aSummary for rodents applies to pathogen-free animals; the abundance of goblet cells may increase in the setting of infection or cytokine stimulation.
NA: Unknown or under investigation; T: Trachea; B: Bronchi; Br: Bronchioles; NCC: Nonciliated, nonsecretory columnar cells; SMG: Submucosal glands.

TABLE II
Epithelial Cell Types and Identified Potential Progenitor Cell Types in Mouse Adult Lung

Cell type	Markers	Progenitor potential	Daughter cells	Pollutant sensitivity	References
Basal	Cytokeratin 5, Cytokeratin 14	Yes	Basal, PNEC? Secretory, Mucous, Ciliated	NA	Hong <i>et al.</i> , 2004a; Schoch <i>et al.</i> , 2004
Clara	CCSP, Cyp450 2F2	Yes	Clara, PNEC? Mucous, Ciliated, AEC I, II	Naphthalene	Hong <i>et al.</i> , 2001; Kim <i>et al.</i> , 2005; Reynolds <i>et al.</i> , 2000b
Ciliated	Tubulin IV	NA	NA	NO ₂ , SO ₂	Borthwick <i>et al.</i> , 2001
Mucous	Mucin 5AC, Mucin 5B	Yes	Basal, Mucous, Ciliated	Polidocanol	Engelhardt <i>et al.</i> , 1995; Hook <i>et al.</i> , 1987
AEC I	Aquaporin 5	No	NA	Bleomycin	Emura, 1997
AEC II	Lamellar body, SpA, SpB, SpC, Aquaporin 1	Yes	AEC I, AEC II	Bleomycin	Griffiths <i>et al.</i> , 2005; Reddy <i>et al.</i> , 2004
PNEC	CGRP	No?	Clara?	NA	Hong <i>et al.</i> , 2001; Peake <i>et al.</i> , 2000
SP	Sca-1, CD31 Hoechst 33342	Yes	NA	NA	Giangreco <i>et al.</i> , 2004; Summer <i>et al.</i> , 2003

NA: Unknown or under investigation.

TABLE III

Some of the Pollutants or Regents Used to Generate Lung Injury Models

Pollutants	Injury methods	Injury mechanism or targets	References
O ₂ , O ₃ , NO ₂ SO ₂	Inhalation Inhalation	Hyperoxic Conducting airway	Meulenbelt <i>et al.</i> , 1992b; Smith, 1985 Asmundsson <i>et al.</i> , 1973; Langley-Evans <i>et al.</i> , 1996
Polidocanol Naphthalene Bleomycin	Intratracheal instillation Intraperitoneal injection Intratracheal instillation	Surface of airway and alveoli Clara cells Alveolar Type I cells, alveolar Type II cells	Borthwick <i>et al.</i> , 2001; Suzuki <i>et al.</i> , 2000 Reynolds <i>et al.</i> , 2000b; West <i>et al.</i> , 2001 Bigby <i>et al.</i> , 1985
Elastase Radiation	Intratracheal instillation Exposure	Alveolar walls Exposed regions	Dubaybo <i>et al.</i> , 1991 Theise <i>et al.</i> , 2002