MINERVA

“Body stuffers” are drug dealers who conceal drugs in their mouths so they can quickly swallow them if they’re approached by the police. This differs from body packers (“mules”), who smuggle packages of drugs in their bowels. In Hamburg, anyone suspected of having swallowed drugs in order to hide them is taken to the Institute of Legal Medicine, where drugs are retrieved by pharmacologically inducing vomiting. Since 2001 this method has successfully recovered drugs in two thirds of suspected cases. Most of the offenders were African men aged 16 to 25 (*Journal of Forensic and Legal Medicine* 2007;14:96-8).

Despite our hunger for all things digital, especially when they’re free, Minerva was amused to discover that the demand for the printed directory of UK Self Help Groups and Support Organisations at £20 a pop continues to grow (see www.ukselfhelp.info/order). The website (www.ukselfhelp.info), containing the most up to date information, remains free of charge.

Amid the latest rounds of changes to medical postgraduate training in the UK, the quality of our future surgeons is to be determined by a move away from apprentice-style training to one of more ordered education. One surgical specialist registrar observes that “it says much for our island mentality that although we belong to the European Union, we consider European surgical training to be distinct from our own” (*Bulletin of the Royal College of Surgeons of England* 2007;89:12-4).

If a slump in blood pressure after lunch increases the risk of coronary events in older people, a postprandial cup of coffee might perhaps reduce it. Analysis of 426 cardiovascular deaths in more than 6500 people with no baseline history of cardiovascular disease found that the over 65s with a habitual higher intake of caffeinated drinks had a lower relative risk of cardiovascular disease and mortality than those who habitually drank less caffeine. The protective effect was dose related, but it was found only in those who weren’t severely hypertensive (*American Journal of Clinical Nutrition* 2007;85:392-8).

A randomised double blind pilot study of long chain omega 3 essential fatty acid supplementation in people who recurrently self harm reports that, at just 12 weeks, the active group had greater improvements in scores for depression, suicidal ideation, and daily

A 65 year old woman was seen in the dermatology clinic with a large friable nodule on her fingertip; it was removed by curettage. Histology confirmed the diagnosis of pyogenic granuloma, a lobulated proliferation of capillary vessels. These common benign tumours occur predominantly on the fingers, lips, face, and gingiva. In most cases no definite cause is found, although minor trauma is often suspected. This lesion may have resulted from a scratch sustained in the garden a week earlier.

K J Hunter SpR, T M Finch consultant, department of dermatology, Solihull Hospital, Solihull B91 2JL katie.hunter@nhs.net

stress. Scores for hostility, impulsivity, and aggression remained the same in both groups. The unanswered question is whether insufficient dietary intake of these essential fatty acids is a reversible risk factor for self harm (*British Journal of Psychiatry* 2007;190:118-22).

Rest alone for osteochondritis dissecans of the knee sometimes fails. To improve blood flow and stability in the articular cartilage and its underlying bone, a recent innovation is to arthroscopically transplant “plugs” of osteochondral cells taken from the non-weight bearing area of the patient’s own knee joint into the unstable fragments. Using magnetic imaging to track 12 knees treated in this way, surgeons found that in all 12 the interface between the avascular fragment and the subchondral bone had disappeared by three months, with no complications arising from the donor site (*American Journal of Sports Medicine* 2007;35:216-22).

Fewer than 50% of strokes and transient ischaemic attacks were reported as being a probable stroke when reported to the Australian emergency services in a study of stroke and the decision to call for an ambulance. Unprompted recognition of a stroke was associated with facial droop and a previous history of stroke or transient ischaemic attacks, and the factors associated with making the call within an hour of the symptoms starting were speech problems, a family history of stroke, and the event being witnessed by another person (*Stroke* 2007;38:361-6).

Mothers with drug habits put not just their own parenting skills at risk but they pose problems for their children’s future functioning as parents, according to a longitudinal study in *Pediatrics* (2007;119:444-51). The results suggest that interventions aimed at reducing drug use in one generation could have knock-on effects on the self esteem of not only the second generation but also the third. Put another way, if the grandchildren’s self esteem is low, multigenerational family therapy may be helpful.

Quite a few adults with epilepsy can actually predict their own seizures before they happen (*Neurology* 2007;68:262-6). The ones who do this best, according to complex mathematical modelling, are young people with high rates of seizures. The researchers didn’t explore the premonitory symptoms reported to be present in up to half of patients with epilepsy, but it may be these patients who are the successful predictors and therefore the potential candidates for prophylactic treatment for seizures.

Snorting cocaine may cause nasal problems, but ironically it’s still the local anaesthetic of choice for surgeons performing septoplasties, who administer it by inserting drug-dipped sponges into the nostrils. A randomised controlled trial of cocaine 4% solution versus tetracaine 2% solution concludes that tetracaine is significantly better at pain reduction than cocaine, and that the use of cocaine should now be limited (*Journal of Laryngology and Otology* 2007;121:130-3).