Correspondence

Letters to the Editor should not exceed 500* words

Antidepressant Drugs and Liver Damage

SIR,—It is very disturbing that, despite being repeatedly asked to do so in this column, no British or American general physician or pathologist seems to be able to tell us how they diagnose between infective hepatitis and “nardil” (phenelzine), “drazine” (phenoxypropazine), or “marsilid” (iproniazid) jaundice, either with or without a full post mortem to help them. Yet their evidence has resulted in the banning of iproniazid in the U.S.A., and the unnecessary panic withdrawal of phenoxypropazine in this country; and now the Witts Subcommittee has published its first pamphlet through the Committee on the Safety of Drugs warning doctors that patients may occasionally die of jaundice if they prescribe these drugs.

Dr. Gerald Garmany's common sense has surely let him down when he uses it to bolster his belief in “largactil” (chlorpromazine) jaundice (25 January, p. 237). Here again, early on, general physicians and pathologists somewhat foolishly rushed in and described its special liver pathology. But although one saw very many cases of so-called chlorpromazine jaundice when the drug was first used, one practically never sees them now, perhaps because there is less infective hepatitis about, and despite the fact that we are now using chlorpromazine in truly enormous doses in very many more patients than then.

I have sent Dr. P. H. A. Willcox evidence of his father's belief that the barbiturates also caused fatal jaundice (25 January, p. 237). I was indeed a very young doctor, but happened at the time to be medical superintendent of St. Mary's Hospital; and later on, when at the Maudsley Hospital, I remember a very highly placed psychiatrist being very worried indeed that he might be blamed at an adjourned inquest because a patient had died of jaundice whom he had been treating with barbiturates. And what about all the general physicians who went on insisting that arsenic injections for syphilis and general paralysis of the insane could cause fatal jaundice until the proper sterilization of syringes diminished the risk of passing on infective hepatitis?

Finally, may I beg Sir Derrick Dunlop and Professor L. J. Witts to agree to appoint at least one officially named psychiatrist to their committees on the safety of drugs, as requested by the Royal Medico-Psychological Association, and stop relying on secret psychiatric advice? There are well over 600 psychiatric consultants now responsible for the treatment of all the patients in nearly half the total hospital beds in this country, and of thousands of psychiatric out-patients as well. If we really did take the out-of-date advice of the Witts Subcommittee in their recent pamphlet on the monoamine-oxidase inhibitor drugs, it would neither help the patients nor ourselves. It also seems, as a page 694

result of it, that we are now much more likely to be censured or sued in the courts, whether we deliberately do accept or very wisely refuse to follow some of the detailed advice given. The history and dangers of general physicians advising psychiatrists on how to use their drugs is repeating itself with a vengeance.—I am, etc.,

WILLIAM SARGANT.

Department of Psychological Medicine,
St. Thomas's Hospital,
London S.E.1.

SIR,—As the contact man for a gang of smugglers in these sometimes United States I am highly disturbed by the recent correspondence in the *British Medical Journal* concerning the possible withdrawal of “marsilid” (iproniazid) from the British market. This drug was “voluntarily” withdrawn by the manufacturer on this side of the ocean after the big-bad-wolf Food and Drug Administration huffed and puffed a bit.

In a review¹ of the subject carried out at that time based on the prescription audit from April 1957 to February 1958, inclusive, approximately 380,000 persons received iproniazid. Dr. Eugene Jolly² of the Food and Drug Administration estimated an incidence of 1 case of jaundice in every 10,000 patients. Considering the usual incidence of infectious jaundice in such a population (approximately 1 in 5,000) it appeared that iproniazid was an excellent drug for protection against jaundice.

A number of small groups of iproniazid-users have now arranged for friends to purchase supplies for them when they visit England and by disguising the drug with such misleading notations as “for la grippe” or “use in case of diarrhoea” they have been able to get it past the customs. If the source of supply in England is dried up we may soon find ourselves in underground commerce with the Russians, since the drug is still manufactured there as “iprozid.”

To save some factifuging letter-writer the trouble I am delighted in advance to acknowledge that I am the Kline who with Loomer and Saunders introduced marsilid as a specific against depression back in 1957. Since that time I have had a number of patients on the medication for whom none of the newer mono-amine oxidase inhibitor drugs adequately substitute. Let me therefore throw in my lot with other chauvinists such as William Don Quixote Sargent, who hold that truth lies in reality rather than with an overly sensitive and overly sensitized Government bureau.—I am, etc.,

NATHAN S. KLINE.

Research Faculty,
Rockland State Hospital,
Orangeburg, New York.

REFERENCES

- ¹ Floody, R. J., Dixon, R. E., and Mattia, V. D., *Dis. nerv. Syst.*, 1958, 19, 541.
- ² Jolly, E. R., *ibid.*, 19, 548.

SIR,—A recent report in a national newspaper suggested that the toxic effects of certain drugs in common usage might be overlooked by some medical practitioners due to the lack of time available to read the mass of literature on medical products deposited through the letter-box. As examples, two recent reports of deaths following the use of “nardil” (phenelzine) in combination with other drugs were mentioned. One of the reports resulted from a post mortem which I performed, and in order to avoid further misinterpretation I feel that a brief description of the case is justified.

A 55-year-old woman, who had been given treatment for G.P.I. and depression on four previous admissions since 1957, was readmitted with severe depression and a history that she had “collapsed” on three occasions that day. Examination did not reveal any physical abnormality apart from extensor plantars and a blood-pressure of 180/110. She died within 24 hours of admission. At the post mortem the only abnormal finding was cerebral oedema. No macroscopic or microscopic evidence of cerebral syphilis was found. The organs and a blood sample were sent for analysis, which showed the presence of little more than therapeutic levels of imipramine, meprobamate, “largactil” (chlorpromazine), and amylobarbitone. The amylobarbitone had been prescribed on admission, but the only other drugs prescribed in the past two months had been phenelzine 25 mg. three times a day and “librium” (chlordiazepoxide) 10 mg. three times a day. It was presumed that the patient had recently taken the imipramine, meprobamate, and chlorpromazine from supplies remaining after previous courses of these drugs.

My attention was drawn to the earlier reports of death following the administration of phenelzine in combination with other drugs. I agreed that the combination of phenelzine and imipramine was a possible significant fact in this case, but that the evidence was insufficient; and I do not feel that the case should be used to illustrate the alleged toxic effects of the combination of the two drugs. A greater degree of proof is required.—I am, etc.,

Department of Pathology, A. M. BARLOW.
Royal Infirmary,
Huddersfield, Yorks.

SIR,—At a recent inquest a verdict of misadventure caused by poisoning due to a combination of “tryptizol” (amitriptyline) and “nardil” (phenelzine) tablets having been taken was recorded.

The fact that both types of tablet were taken grossly in excess of the prescribed dose was not apparently considered to have any real bearing on the case, giving the impression that a normal therapeutic dose of these drugs taken together is inevitably fatal. This is not the view of many members of the medical profession, nor is it substantiated by the number of recorded deaths due to this drug combination.

Forensic tests showed the presence of amitriptyline in the body but no trace of phenelzine. Apparently there was no quantitative analysis done. The pathologist gave the cause of death as due to the combined action of these drugs, as he knew of no reported deaths due to amitriptyline alone