uterine haemorrhage in several instances of chorioncarcinoma. Toxic effects with this form of treatment are usually slight or absent, and it lacks the finality of hysterectomy on the patient's child-bearing capacity.

The risks and unpleasantness of systemic chemotherapy are not to be underestimated, but when it offers a substantial expectation of cure and of further normal pregnancies it would be unfortunate if serious delays in its application result from any impression that it is a fate worse than death.—I am, etc.,

K. D. BAGSHAWE.

Charing Cross Hospital Medical
School,
Fulham Hospital,
London W.6.

REFERENCES

- ¹ Lipp, R. G., Kindschi, J. D., and Schmitz, R., *Amer. J. Obstet. Gynec.*, 1962, **83**, 1644.
- ² Bagshawe, K. D., and Garnett, E. S., *Brit. J. Radiol.*, 1963, **36**, 673.
- ³ Bagshawe, K. D., *Brit. med. J.*, 1963, **2**, 1303.

Buccal Oxytocin

SIR,—Some of the comments in the article entitled "Drugs Acting Upon the Uterus" (9 May, p. 1234) require amplification.

There is no evidence that buccal oxytocin is absorbed irregularly. Elstein and Payling-Wright¹ make the following observations: "In view of the relatively large doses of oxytocin in the tablets and the uncertainty of the rate of absorption, oxytocin assays were attempted further to assess the safety of the preparation. Observations were made on blood samples taken before and during the administration of buccal pitocin, using the milk ejection method of Tindal and Yokoyama (1962).² No oxytocin was demonstrated in the initial samples, but the second yielded 20 μ -units/ml. blood, a figure which falls at the lower limit of sensitivity of the test."

There is little knowledge on the subject of absorption because of the difficulty of detecting oxytocin in the blood specifically. As pointed out in another section of the article there are wide variations in uterine sensitivity to oxytocin, and this precludes the consideration of uterine activity as a measure of the absorption rate.

The dosage regimen quoted has been superseded by entirely different scales. At present in the United Kingdom the first dose is 100 units, tablets are given half-hourly, and the dose increased hourly until it reaches 600 units. This method is followed unless adequate response is obtained at a lower dosage. Finally, precise control is possible and tocography by Elstein and Payling-Wright¹ shows 50% reduction in uterine activity 15 minutes after the tablets are withdrawn. This is an identical result to that obtained by Sica-Blanca and Sala,³ when stopping an intravenous infusion of oxytocin.—I am, etc.,

H. C. MASHETER.

Department of Clinical Investigation,
Parke Davis & Co.,
Hounslow, Middlesex.

REFERENCES

- ¹ Elstein, M., and Wright, H. P., *J. Obstet. Gynaec. Brit. Cweth.*, 1963, **70**, 1005.
- ² Tindal, J. S., and Yokoyama, A., *Endocrinology*, 1962, **71**, 196.
- ³ Sica-Blanca, Y., Sala, N. L., *Oxytocin*, ed. R. Caldeyro-Barcia and H. Heller, 1961, p. 127. Pergamon Press, Oxford.

Quinoline Derivatives in Dermatology

SIR,—For the past thirty years dermatologists have found topically applied quinoline derivatives useful as antiseptics in cases of pyoderma, fungous infection, and infected eczema, and in more recent years they have been used in combination with topical steroids.

However, antibiotic/corticosteroid combinations have become more widely used in the treatment of infected eczema, with neomycin being the most commonly used antibiotic. There is good reason to believe that this antibiotic can no longer be used with complete safety, as there have been several references recently to patients developing skin sensitization and also reports of neomycin-resistant strains of staphylococci.¹⁻³ It would seem therefore that the quinoline antiseptics should regain their previous range of usefulness.⁴ Corticosteroid preparations containing in addition 8-hydroxyquinoline antiseptics may now be the applications of choice in the treatment of infected eczemas, especially in view of their antifungal as well as antibacterial action.⁵ It would further seem desirable to use one of the newer, more potent fluorinated-hydrocortisone derivatives rather than the parent corticosteroid.

The iodine content in the iodochlorhydroxyquinolone derivative can cause an occasional contact or sensitization reaction which may not be suppressed even when this substance is used in combination with hydrocortisone. Additionally, much more effective and less liable to stain and sensitize than the iodochlorhydroxyquinolone-hydrocortisone combination is a halquinol (chlorhydroxyquinolone)-triamcinolone acetate cream and ointment which have been available for clinical trial during the past six months. So far I have not noted any untoward reaction in using them in a hundred cases of infected and non-infected atopic and discoid eczemas, seborrhoeic folliculitis, otitis externa, paronychia, and intertrigo, and under polyethylene in the treatment of psoriasis. One patient amongst nineteen psoriatics so treated did develop a folliculitis of the legs; this is less than would be anticipated if triamcinolone acetate or other steroid application had been used alone under polyethylene in treating psoriasis.

Chlorhydroxyquinolone - triamcinolone cream (now known as "remiderm") has also proved valuable in refractory cases of intertrigo of groin and natal cleft where there is a superimposed monilial infection.—I am, etc.,

Hove, Sussex.

PATRICK HALL-SMITH.

REFERENCES

- ¹ *Brit. med. J.*, 1963, **2**, 981.
- ² Robertson, J. J., *Lancet*, 1963, **2**, 333.
- ³ Jacobs, S. I., and Willis, A. T., *ibid.*, 1963, **2**, 459.
- ⁴ Peterkin, G. A. C., *Practitioner*, 1961, **187**, 446.
- ⁵ Vickers, H. R., *ibid.*, 1964, **192**, 639.

Inherited Characters in Twins

SIR,—Mr. J. M. Anderson (23 May, p. 1384) is, of course, perfectly correct in pointing out that discordances in inherited characters have been observed in identical twins. However, armadillos aside, such discordances are rare.

The occurrence of a thymoma with hypogammaglobulinaemia in an identical twin which I reported (2 May, p. 1159) was the first known such association. The twins were identical in all parameters measured other than serum proteins. In addition, the patient lived to the age of 63 years with no previous evidence suggestive of antibody deficiency. I feel that these two facts strongly support an acquired aetiology.

The purpose in reporting such rare conditions is to draw attention to the subject in the hope that further cases will be found, and establish its significance.—I am, etc.,

Brompton Hospital,
London S.W.3.

SIMON GODFREY.

Aflatoxin

SIR,—Your leading article (8 February, p. 325) states that "even now nobody knows whether it [aflatoxin] is commonly found in the diet in countries where liver cancer is common." You may be interested to learn that a random selection of cereals bought on one visit to the local market here during the last rainy season yielded not only toxic aspergilli but a toxin-producing strain of *Penicillium islandicum*, already shown by Japanese workers to produce a hepatotoxin.¹ The grains included sorghum and millets, much used in the making of local beers, and the covergrass "tef" (*Eragrostis abyssinica*), used to make the staple "injera," a flat sour bread cooked at a low temperature after two or three days' fermentation. I was intrigued to read a few pages further on (p. 336) that some of the liver cancers reported in the Mengo survey "may have been cholangio-cellular tumours"; could they have been like the bile-duct proliferation reported in cattle fed on toxic groundnuts?²

Now, while native beers and primary liver cancer run together from here to the Cape, we have in Ethiopia, it seems, significantly less of the widespread protein malnutrition than is reported among our southern neighbours. Since protein lack as a possible causative agent of liver disease has already received a good deal of study elsewhere, Ethiopia seems a good starting-point for a new approach.

Accordingly it is hoped to pursue the mycological survey of local grains with outside help, concurrently with a closer examination of brewing, baking, harvesting, and storage practices. I would be grateful meanwhile for any relevant information any reader of this letter could send me at this address.—I am, etc.,

A. COADY.

Princess Tsahai Memorial Hospital,
Addis Ababa, Ethiopia.

REFERENCES

- ¹ Uruguchi, K., *Jap. J. exp. Med.*, 1961, **31**, 19.
- ² Clegg, F. G., and Bryson, H., *Vet. Rec.*, 1962, **74**, 992.

Infection with Animal Helminths

SIR,—The article by Dr. A. W. Woodruff and others (18 April, p. 1001) and the subsequent correspondence in the *Sunday Times* draws timely attention to the hazard which domestic dogs and cats may constitute to human beings; this applies particularly to children under 5 years of age in surroundings contaminated by canine or feline faeces.