patients; improvement in swallowing occurred within 36 hours and complete relief of dysphagia by the fourth day. Rapid healing of the oesophagus occurred, and it was confirmed that the radiological appearances return completely to normal within 10 to 14 days (Kaufman *et al.*, 1960). Amphotericin B is also effective (Egeblad, 1961), but would seem to possess little advantage over nystatin in the treatment of gastro-intestinal moniliasis.

The simultaneous administration of nystatin during broad-spectrum antibiotic therapy (Newcomer, Wright, and Sternberg, 1954-5) reduces intestinal overgrowth by candida and should be considered when treating infections in patients who are particularly prone to develop monilial infections.

Summary

Two cases of painful retrosternal dysphagia due to monilial oesophagitis are described. Oesophageal involvement was the

presenting and predominant feature of the monilial infection, there being little evidence of thrush elsewhere. The value of radiology in assisting in diagnosis is emphasized.

Our thanks are due to Dr. E. R. Cullinan and Dr. K. O. Black, under whose care these patients were admitted, for permission to publish details of these cases.

REFERENCES

- Andrén, L., and Theander, G. (1956). *Acta radiol. (Stockh.)*, **46**, 571.
 Bernay, P., and Houver, A. (1961). *Arch. Mal. Appar. dig.*, **50**, 233.
 Browne, S. G. (1954). *Lancet*, **1**, 393.
 Eban, R. E., and Symers, D. A. (1959). *J. Fac. Radiol. (Lond.)*, **10**, 164.
 Egeblad, M. (1961). *Dan. med. Bull.*, **8**, 163.
 Kaufman, S. A., Scheff, S., and Levene, G. (1960). *Radiology*, **75**, 726.
 Kaufmann, H. J. (1958). *New Engl. J. Med.*, **258**, 1143.
 Marsh, A. P. (1959). *Amer. J. Roentgenol.*, **82**, 1063.
 Newcomer, V. D., Wright, E. T., and Sternberg, T. H. (1954-5). *Antibiot. Ann.*, p. 686.
 Woods, J. W., Manning, I. H., and Patterson, C. N. (1951). *J. Amer. med. Ass.*, **145**, 207.

Medical Memoranda

Temporary Feeding-gastrostomy— A New Gastrostomy Catheter

Brit. med. J., 1964, **1**, 822

Gastric decompression with planned replacement of fluid and electrolyte, although of established value in the management of patients having operations on the gastro-intestinal tract, has several disadvantages. Intravenous infusion alarms some patients, and discomfort caused by the naso-gastric tube is complained of by many. The risk of post-operative chest complications may be increased in two ways: the presence of a tube in the nasopharynx discourages coughing; the infusion and aspiration equipment anchor the patient to his bed and limit active movement. Intravenous therapy may cause painful vein-thrombosis. Finally, injury to the larynx and stricture of the lower oesophagus have been attributed to gastric intubation.

It is not surprising, therefore, that some surgeons no longer use gastric aspiration after Polya gastrectomy. However, many consider aspiration advisable after Billroth I gastrectomy and

Studies made in this department during the past two years have provided the following information. Firstly, Ross *et al.* (1963) have shown that the mean duration of inhibition of small-bowel motility after surgical vagotomy is as little as 10 hours. Secondly, in tests performed at 18 hours after vagotomy, Cox (1962) has shown that there is already a substantial absorption of monosaccharides, electrolytes, and water. These results support the view that post-operative duodenal or jejunal feeding is a satisfactory substitute for intravenous feeding. Finally, in an investigation to be reported, Ross has found complete inhibition of gastric motor activity for at least 36 hours after vagal section, evidence which is in support of the view that gastric aspiration is advisable during the first two days after this operation.

In the light of these findings, a request was made to Messrs. C. F. Thackray Ltd., Park Street, Leeds 1, to devise a catheter similar to the Foley catheter which would provide a means of gastric aspiration and also permit duodenal feeding after vagotomy with pyloroplasty, or jejunal feeding after vagotomy with gastrojejunostomy. A variety of feeding-gastrostomy tubes have been made for our evaluation. The final product, whose specifications are now given and which has been used by us for six months, is here illustrated.

The three-way F.G. catheter is made of plastic and is fitted with a 20/30 ml. balloon. Two pairs of opposing eyes on the distal side of the balloon facilitate gastric aspiration. The feed-tube protrudes approximately 11 in. (28 cm.) beyond the catheter end and has four eyes spirally arranged at the tip. The catheter is supplied sterile.

after vagotomy, although it should be noted that Hendry (1962) has claimed excellent results from a regimen of restricted fluids without use of gastric suction.

For almost five years it has been our practice to use a Foley catheter as a temporary gastrostomy in patients who have had vagotomy with a drainage procedure. The catheter is inserted through a small stab incision in the anterior wall of the stomach, drawn through an avascular part of the omentum, and finally withdrawn from the abdominal cavity through a small stab incision in the left epigastrium. This exit wound is usually dry within 24 hours of removing the catheter on the seventh day after operation. Gastric decompression by this means has proved to be a safe procedure and one which obviates the discomforts of a naso-gastric tube. However, fluid and electrolyte replacement may be required in some patients.

Patients unfortunate enough to have had previous experience of gastric surgery have acclaimed this form of post-operative management and recalled the discomfort associated with conventional naso-gastric suction. Intravenous therapy is unnecessary and replacement of electrolytes and fluids via the intestine simplifies the problem of maintaining the composition and distribution of the body fluids. Restriction to bed is unnecessary. A controlled study to determine the incidence of post-operative chest complications in patients with and without nasogastric suction has been planned.

ANDREW W. KAY,
Professor of Surgery, Royal Infirmary, Sheffield.

REFERENCES

- Cox, A. G. (1962). *Lancet*, **2**, 1075.
 Hendry, W. G. (1962). *Brit. med. J.*, **1**, 1736.
 Ross, B., Watson, B. W., and Kay, A. W. (1963). *Gut*, **4**, 77.