Correspondence

Letters to the Editor should not exceed 500 words.

Work After a "Coronary"

SIR,—Many readers who have to advise patients after a myocardial infarct will be grateful for the sound general principles outlined in your excellent leading article on "Work after a 'Coronary'" (19 September, p. 703).

We share their gratitude but are sure that the sentence, "Avoidance of foods with a high content of fat or cholesterol is also sound advice . . ." should not be allowed to stand without critical comment. In fact we know of no conclusive evidence to support this advice to patients who have already developed the overt clinical features of ischaemic heart disease. Few formal long-term studies of the effects of such a diet have been made and none has shown that the prognosis is improved. Moreover, so far as drugs are concerned, you yourselves point out that "none of the cholesterol-lowering agents has been shown to have a favourable clinical effect." This is also true for diet; recently Ball and his colleagues¹ have reported a four-year study of the effect of a low-fat diet. They showed clearly that, despite reduction of cholesterol levels, the diet had no demonstrable effect in lowering the relapse rate after myocardial infarction.

We presume that the line of thought in your leader is that the correlations which have been shown to exist between a high-fat diet and ischaemic heart disease in population studies can be interpreted as indicating that the high fat diet is a cause of the ischaemic heart disease. This assumption has not been substantiated and there are certainly many other possible and important aetiological factors in coronary atherosclerosis. It should also be pointed out that, even if the assumption were true, it still does not follow that the institution of a diet low in fat or cholesterol would be of any value to patients whose atherosclerosis has advanced to the stage of angina or myocardial infarction.

Clinicians concerned with the management of patients have to be wary of the dual approach revealed in population studies. It is all too easy to assume false correlations; pre-supposition that low-fat diets are of value in the management of ischaemic heart disease is one unfortunate result.

We recognize that many proved advances in medicine have been based initially on unproved hypotheses. It is, however, wise to realize just how unproved dietetic restrictions are in the management of ischaemic heart disease. It is perhaps appropriate to remember Dr. Armand Vincent's patient² who said: "We are prevented from dying; we are not helped to live." Your leading article recognizes this criticism of medicine. It emphasizes the need to encourage patients to resume a normal way of life after myocardial infarction. It is a pity to spoil it by advocating "hope-it-may-do-good" dietetic measures when these are unsubstantiated and may make

our patients miserable and prevent them living in the fullest sense.—We are, etc.,

R. M. MARQUIS.

M. F. OLIVER.

Department of Cardiology,
The Royal Infirmary,
Edinburgh 3.

REFERENCES

- 1 Ball, K. P., McAllen, P. M., Pilkington, T. R. E., Sharland, D. E., Sowry, G. S. C., and Wilkinson, P., *Proc. IV Europ. Congr. Cardiol.*, 1964, p. 24.
- 2 Vincent, A., *Journées d'étude*, 1950. Paris, Xth year, No. 13. "Pour une médecine humaine."

SIR,—It is interesting that, at this time, you can still include in your leading article of 19 September (p. 703) the recommendation that persons with ischaemic heart disease should reduce their intake of fat, and exclude any suggestion that they should take less sugar.

There has never been any direct evidence for the hypothesis that fat consumption has anything to do with causing the disease; indeed, recent evidence points increasingly against it. For example, studies in East Africa have shown that cardiac ischaemia is rare both in the Samburu¹ and in the Masai,² who consume very large amounts of fat, almost all of it from meat and milk. Results have recently been published of a prospective study in American men, in which no difference in fat intake was found between those who developed ischaemic heart disease and those who did not.³

On the other hand, there is now quite impressive evidence that a high intake of sugar (sucrose) may be an important factor in producing coronary disease. Firstly, epidemiological studies in which populations are compared show a rather better relationship of the incidence of the disease with sugar consumption than with fat consumption.^{4,5}

Secondly, the increasing incidence of the disease that is found in many countries has followed an increasing consumption of sugar rather than of fat. This was demonstrated in Israel amongst Yemeni immigrants,⁶ and more recently in the United States.⁷ The latter study has shown that in the last 70 years there has been only a 12% rise in fat consumption, and a simultaneous increase in the ratio of polyunsaturated to saturated fats. During the same period, in which the number of people dying of coronary thrombosis has increased very considerably, the consumption of sugar has more than doubled.

Thirdly, although studies with individual subjects have shown no difference in fat intake between those with and those without cardiac ischaemia, they have shown considerable differences in sugar intake.⁸ On average, people that have recently had a first known myocardial infarction have been taking twice as much sugar as control subjects, mostly in their tea and coffee. This is the first time that any difference in diet has

been found between individuals with and without the disease.

It will no doubt be several years before we know whether the reduction of dietary sugar will improve the prognosis of those who have suffered their first coronary attack. In the meantime, however, we might reasonably suggest that people at risk should be advised to cut their intake of sugar. One knows of no evidence whatever that this could be harmful, whereas there is increasing suspicion that continued high intake of sugar could be harmful.⁹—I am, etc.,

Department of Nutrition,
Queen Elizabeth College,
London W.8.

JOHN YUDKIN.

REFERENCES

- 1 Shaper, A. G., Jones, M., and Kyobe, J., *Lancet*, 1961, 2, 1324.
- 2 Mann, G. V., Shaffer, R. D., Anderson, R. S., and Sandstead, H. H., *J. Atheroscler. Res.*, 1964, 4, 289.
- 3 Paul, O., Lepper, M. H., Phelan, W. H., Dupertuis, G. W., MacMillan, A., McKean, H., and Park, H., *Circulation*, 1963, 28, 20.
- 4 Yudkin, J., *Lancet*, 1957, 2, 155.
- 5 — *ibid.*, 1964, 2, 4.
- 6 Cohen, A. M., Bavy, S., and Poznanski, R., *ibid.*, 1961, 2, 1399.
- 7 Antar, M. A., Ohlson, M. A., and Hodges, R. E., *Amer. J. clin. Nutr.*, 1964, 14, 169.
- 8 Yudkin, J., and Roddy, J., *Lancet*, 1964, 2, 6.
- 9 — *ibid.*, 1963, 1, 1335.

Misleading Exfoliative Cytology

SIR,—I was glad to see Dr. J. Elizabeth Macgregor being a little more realistic about negative smears in the presence of clinical carcinoma of the cervix (19 September, p. 754). There can be little doubt, as she says, that biopsy is required for any cervix which has the appearance of malignancy, and I wish she had gone a little further and said that smears in such cases are chiefly of academic interest and not necessary to the acceleration of the patient's diagnosis and treatment.

I would like her to reconsider her last sentence where she says that cervical and vaginal smears should be examined before curettage, when body carcinoma is suspected. I have turned her sentence round to make the position a little clearer, for I believe it imperative that where body carcinoma is suspected diagnostic curettage should be arranged—once more, as for clinical cancer of the cervix, *whatever* the smear report.—I am, etc.,

Department of Obstetrics
and Gynaecology,
University of Edinburgh.

A. F. ANDERSON.

SIR,—I was very pleased to see the letter from Dr. J. Elizabeth Macgregor of the Department of Midwifery and Gynaecology at Aberdeen (19 September, p. 754). I recently had a "neurotic" patient who was seen by a first-class consultant gynaecologist and reassured on three occasions, especially after the last time when a cervical smear was taken, that there was no cancer present. However, I regret to say that some two months later she had a total hysterectomy for a carcinoma of the body of the uterus.

While I would obviously agree with cervical smears I do feel that some of the obses-