intercostal muscles to provide either an adequate cough mechanism or augmented respiratory excursion.

(2) In cases of cord section between C.2 and C.6, upper trapezius fibres to operate voluntary finger flexion. In cases in which shoulder girdle and arm muscles have been spared but hand function is deficient, it is hoped that this defect may thus be remedied on a controlled voluntary basis.

(3) Ultimately perhaps, by cross over from anterior deltoid fibres to hip flexors of opposite side, to provide a method of walking.—I am, etc.,

SHANE WATSON.

National Spinal Injuries Centre,
Stoke Mandeville Hospital,
Aylesbury, Bucks.

Eliminating Cervical Cancer

SIR,—Your editorial on the above subject (5 December, p. 1409) will command wide support, but in present economic circumstances statements of objectives do not of themselves achieve objectives. The responsibility for the laboratory examination of cervical smears and the provision of gynaecological services associated with these laboratory services is that of regional hospital boards. The laboratory resources, both in staff and buildings, that can immediately be provided by the boards obviously cannot meet the requirements of a complete programme whereby at regular intervals all women who should have an examination can be assured of the necessary facilities being provided.

Since the National Health Service must, like any other social service, be limited in the share that it can command of the national resources in money and manpower, it follows that the development of laboratory resources can eventually only be at the expense of development in other fields. It is therefore imperative that laboratory resources for cervical cytology, both in buildings and manpower, should, as they are developed, be utilized in an orderly and ordered fashion so that they can be put to the best use for the greatest number of women in accordance with agreed and predetermined programmes.

The Association of County Medical Officers has given consideration to this matter and has come to the conclusion that the best use cannot be made of existing and expanding facilities if specimens are going to be submitted in accordance with no predetermined pattern from general practitioners, family planning clinics, and other sources. In coming to this conclusion, the Association does not in any way suggest that specimens sent from these sources should not be examined, but it is of the opinion that if the best use is to be made of resources in short supply, then some ordered programme is essential. It is of the opinion that such a programme can only be developed by the use of existing local health authority resources, including health education, as to the meaning of cervical cytology and the arranging for health visitors, district nurses, and other officers who provide direct services to the public to take part in bringing to the notice of women in the appropriate age-groups the extent and meaning of the services that are being developed for the examination of cervical smears.

The development of these services on the lines suggested requires a close association with the developing services provided by regional hospital boards, and whilst it would be imperative, both initially and at various stages of development, to consult local medical committees, it is the view of the Association that the only way in which an effective programme can be worked out to take full advantage of expanding hospital facilities is for the local health authorities to be charged with the task of providing agreed programmes in consultation with the regional hospital boards and their associated hospital management committees. This view of the Association has been communicated to the County Councils Association with the purpose of asking the Minister of Health to ensure that as the services of the regional hospital boards are developed there shall be throughout full consultation with the local health authorities in order to ensure that what is being provided shall be put to the most effective use.—I am, etc.,

G. RAMAGE,

Secretary,

Association of County Medical Officers
of Health of England and Wales.
Stafford.

Cephaloridine

SIR,—Your comment on my letter (12 December, p. 1530) states that "about 80% of clinical isolates of *Proteus* are *P. mirabilis*." This is not everyone's experience.

For example, Millner¹ found 63% in 130 *Proteus* strains from urinary infections. In this hospital we found 49% of 147 strains from all sources,² whilst Dutton and Ralston³ found only about 35% of 57 strains from urinary infections to be *P. mirabilis*.—I am, etc.,

St. James' Hospital,
London S.W.12.

P. W. KIPPAX.

REFERENCES

- 1 Millner, P. F., *J. clin. Path.*, 1963, **16**, 39.
- 2 Kippax, P. W., *ibid.*, 1957, **10**, 211.
- 3 Dutton, A. A. C., and Ralston, M., *Lancet*, 1957, **1**, 115.

Plantar Warts and Athlete's Foot

SIR,—I find it difficult to accept Dr. A. A. Ross's objections (12 December, p. 1532) to my protest regarding barefoot gymnastics (14 November, p. 1268). Does Dr. Ross believe that an hour or so of barefooted exercise per week (during school terms only) is really likely to have a significant effect on flat-footedness or other foot deformities, particularly when the feet in question spend much of the rest of the week in hallux-valgus-producing shoes? Dr. Ross does not apparently regard a small increase in the number of plantar warts coming to his skin clinic with dismay. But for every extra case which he sees may there not be many treated at home under the guidance of general practitioners? In any case, whatever the theoretical advantages of barefoot gymnastics, the price is paid not in consultant's time, nor in the time of general practitioners, but in the hobbling discomfort of young children and in their fears of the even greater discomfort which may accompany treatment in the clinic. Incidentally, is hobbling a good exercise for the muscles and joints of the lower limbs?

With regard to athlete's foot, does Dr. Ross really advocate that children with the active condition should mix barefoot with uninfected children on the floor of the gymnasium? Perhaps his comments on this point were not intended to be other than superficial.—I am, etc.,

F. J. C. ROE.

Chester Beatty Research Institute,
London S.W.3.

Dangerous Anaesthetic Device

SIR,—Dr. Manley (12 December, p. 1532) is quite justified in vindicating his ventilator as being the only agent to interfere with the Bosun's audible warning system. In fact, any apparatus producing a sustained pressure rise of about 25 cm./Wg. (water-gauge) in the anaesthetic system distal to the flow meters may render the audible system inoperable. If this back pressure drops below the figure mentioned at any time then the warning device will sound and will not stop until a pressure of above 100 cm./Wg. is obtained.

Thus, ventilators which operate at low bag pressure (without pressure reduction to the patient circuit), such as the Barnet, East-Radcliffe, etc., and all closed-circuit ventilators are free from this hazard. If the flow control of the later Barnet models is set to give high restriction then back pressures of interfering magnitude may result.

The Manley and the Howells ventilators both maintain a back pressure in excess of 25 cm./Wg. (100–120 cm./Wg.) and therefore interfere with the Bosun's audible warning of oxygen loss.

Incidentally, these latter machines might be indicted for altering the Fluotec output of halothane by virtue of the same back pressures. However, tests performed at Cyprane Ltd. recently indicated that at normal source flow rates (4 litres and upwards) on the Howells ventilator the effect on the Fluotec output was negligible. It is probable that this holds true also for the Manley ventilator.—I am, etc.,

T. HILARY HOWELLS.

Department of Anaesthesia,
Royal Free Hospital,
London W.C.1.

Thyroid Carcinoma in the Newborn

SIR,—In a memorandum on a "Case of Thyroid Carcinoma in a Newborn Infant" (25 January 1964, p. 224) I have clearly been mistaken in ascribing the appearances to carcinoma. Dr. Winship, to whom the bulk of the material was sent, has very kindly drawn attention to a condition described as struma intralaryngotrachealis in Henke and Lubarsch¹ and illustrated by an identical lesion to that found in our case.

On behalf of Dr. Enid E. McRuer and myself I sincerely regret this misidentification of an unusual condition and hope this correction may be of assistance.—I am, etc.,

M. D. ROSS.

Harare Hospital,
Salisbury, S. Rhodesia.

REFERENCE

- 1 Henke and Lubarsch, *Handbucher speziellen pathologischen Anatomie u. Histologie*, Bd. 8, p. 331.