Current Practice

CHILD CARE IN GENERAL PRACTICE

"Normal Development"

R. S. ILLINGWORTH,* M.D., F.R.C.P., D.P.H., D.C.H.

All children are different. They differ in their intelligence, rate of development, their interests and aptitudes, and in their behaviour. In addition, the home environment of one child is very different from that of another and the environment has a profound effect on development. Various illnesses and mechanical or sensory defects greatly affect the course of development. In consequence, developmental assessment is fraught with difficulties. There is no place for spot diagnosis and no place for diagnosis based purely on clinical impressions. Each child has to be considered in detail, in order to avoid serious and unnecessary mistakes. Mistakes may be tragic in some circumstances—as in the assessment of suitability for adoption—a process undertaken far too lightly by many with no experience or knowledge of developmental assessment.

Patterns of Development

A child's intelligence has a profound effect on his development. *A mentally subnormal child is backward in all fields of development, except occasionally in sitting and walking, and still less frequently in sphincter control.* He is relatively more retarded in speech than in other fields, and relatively less retarded in motor development (sitting and walking). He shows much less interest in his surroundings than normal children. It follows that no child can be mentally subnormal merely because he is backward in individual fields of development. It also follows that the least important aspect of development for assessment purposes is the age of sitting and walking. One pays much more attention to his responsiveness, alertness, interest in his surroundings, and concentration, to the use of his hands, and later to his speech.

A mentally superior child may be advanced throughout, but more commonly there is either no recognized sign of superiority in infancy or else it requires an experienced eye to detect his unusually good alertness and social responsiveness, the advanced nature of his vocalizations, and later of his speech.

The rate of development is largely dependent on maturation of the nervous system, but it is also affected by environment. The mentally defective child is slow in maturation, and so is retarded in almost all aspects of development. He is commonly a sleepy baby—and is described by his mother as being "so good" and "not a bit of trouble." A child may be a slow starter—and so backward in all aspects of development. He eventually matures, and becomes average or superior. This pattern is rare.

Delayed maturation in individual fields of development is extremely common and often runs in the family. For instance, there may be a familial pattern of late walking, speech, or sphincter control.

A child's development, and later his intelligence quotient, is considerably affected by his environment. It is retarded

by emotional deprivation, so that a baby brought up in an institution is likely to be backward in all aspects of development. It is accelerated, up to a point, by a good, loving, and stimulating home. Lack of practice will delay development. For instance, if a mother deliberately keeps a child off his legs for fear he will develop knock-knee or bow legs, he will show defective weight-bearing. If she fails to give him the opportunity to use the pottie when he is beginning to ask for it, he will be delayed in acquiring sphincter control. If he is not given a chance to feed himself and dress himself, when developmentally ready for it, he will be retarded in these skills.

A variety of physical factors retard development. These include variations in muscle tone, and visual or auditory defects. Hypotonia retards sitting and walking, and if extreme may make both impossible: hypertonia, as in cerebral palsy, retards motor and manipulative development. Auditory defects retard speech, and if severe enough will make speech impossible without special training. Visual defects will impede manipulative development and to some extent walking.

Varieties of Patterns

One may summarize the different patterns of development as follows:

(I) *Average throughout in all fields.*

(II) *Average, becoming superior later.* This pattern may be due to failure to recognize the early signs of mental superiority, or to slow maturation, or to improvement of the environment.

(III) *Superior or advanced in certain fields.* Perhaps the commonest is advanced motor development. For instance, a child may sit without support by 5 months, instead of the average age of 7 months, or walk without support at 8 months, instead of the average age of 13 months. This pattern certainly does not indicate mental superiority. On the other hand, advanced speech always signifies mental superiority. (Delayed speech certainly does not *per se* denote mental inferiority.)

(IV) *Superior or advanced in all fields.* This signifies mental superiority.

(V) *Average or superior at first, deteriorating or slowing down in development.* This may be due to a worsening environment, with emotional deprivation. It may be due to a variety of illnesses, such as meningitis, encephalitis, and degenerative diseases of the nervous system. Mental deterioration, if not prevented by proper treatment, occurs in hypothyroidism, hypercalcaemia, phenylketonuria, leucosis, nephrogenic diabetes insipidus, the hypoparathyroidisms, galactosaemia, and possibly in certain carbohydrate enzyme defects and in other metabolic conditions. It results from severe hypoglycaemia (as from insulin overdosage), lead poisoning, the toxic effect of drugs (such as those used for epilepsy), or from head injury. It is a characteristic feature of mongolism that development slows down in later infancy. One may see

* Professor of Child Health, the University of Sheffield.

distressing deterioration in association with certain forms of epilepsy.

(VI) *Retarded in certain fields.* The commonest cause of delayed walking in a child with average intelligence is a familial pattern of late walking, or cerebral palsy. It may be due to hypotonia, or to lack of opportunity to practise. It is possibly caused sometimes by obesity. It is not due to congenital dislocation of the hip.

Delayed speech, in a child of average intelligence, may be a familial pattern of development: it may be due to an auditory defect, including high-tone deafness. Often there is no discoverable cause. In very rare cases it is due to psychotic illness, including autism. It is not due to tongue-tie, or to cleft palate, or to laziness—arising from "everything being done for him." It is common in twins for obscure reasons.

Delayed sphincter control, in a child of average intelligence, is commonly a familial pattern, depending on delayed maturation of the relevant part of the nervous system. It may also be due to a structural defect of the bladder neck, or in a girl to an ectopic ureter entering the vagina. In either case there would be constant dribbling incontinence. These are commonly psychological causes, dependent on faulty toilet training.

The most common error is over-enthusiastic training, leading to the child being conditioned against the pottie because of all the unpleasantness associated with it (in the way of smackings and enforced sitting on it), or to the usual response of negativism when over-anxiety is shown.

(VII) *Uniformly retarded throughout.* This means mental defectiveness.

(VIII) *Uniform retardation at first, followed by average or superior development.* This may be due to delayed maturation, which may be a familial pattern. It sometimes follows a really difficult birth, perhaps with cerebral haemorrhage. It may be the result of an initial bad environment, followed by improved circumstances—as when a child brought up in an institution is placed in a good foster home. It may result from the institution of appropriate treatment for a condition such as cretinism.

(IX) *Lulls in development.* One commonly finds that a child appears to make no progress in speech for several months, and then suddenly makes rapid strides forward. A child may be backward in speech at one time and advanced in speech a month later. In assessing such a child one pays much more attention to his understanding of words than to his ability to articulate them.

Diagnosis

Diagnoses are normally made on the basis of the history, the examination, special investigations where relevant, and the interpretation.

History

A careful history is essential in developmental assessment. One needs to know about hereditary diseases, and in particular about mental deficiency in either of the parents. One needs to know about conditions, prenatal or post-natal, which increase the risk of abnormality in the child. These conditions include rubella in early pregnancy, bleeding or toxæmia during pregnancy, extreme prematurity, a low birth weight in relation to the duration of gestation, severe anoxia at birth—especially in the absence of a good cause for it, such as prolapse of the cord—convulsions in the newborn period or later, or a history of pyogenic meningitis. Even if there is a history of any of these conditions, each child has to be considered on its merits, and it is very wrong to pay so much attention to these risk factors that one does not seriously assess the child himself as one sees him. It would be a grave error of judgment to regard a baby as unsuitable for adoption because of any of the above "risk" factors.

The history includes the management of the baby and the history of illnesses. This is necessary in order that one can determine whether there have been any factors which may have affected his development, such as illness or emotional deprivation.

Milestones of Development

The history includes the previous milestones of development. This is essential in order that one can determine the rate of development. The previous milestones in addition help one to compare the mother's version of the baby's development with one's own objective findings. Lateness in milestones, combined with the objective finding of backwardness, means much more than either of these alone.

The following are really essential questions about his milestones:

When did he begin to smile at you when you were talking to him (average 4 to 6 weeks)?

When did he begin to make noises as well as smile when you talked to him (usually one or two weeks after he began to smile)?

When did he begin to turn his head when he heard sounds (average 3 to 4 months)?

When did he begin to hold a rattle placed in the hand and play with it (average 3 to 4 months)?

When did he reach out for a toy and get it without it being placed in his hand (average 5 months)?

When did he begin to chew (average 6 months)?

When did he first sit for a few seconds on the floor without support and without rolling over (average 7 months)?

When did he begin to roll completely over from prone to supine (average 5 months) and from supine to prone (average 6 months)?

When did he begin to pull himself to the standing position at the side of furniture (average 8 months), to walk holding on to furniture (average 10 months), to creep on hands and knees (average 10 months), to walk without support at all (average 13 months)?

When did he begin to hold his arm out for a coat, or foot out for a shoe (average 10 months), wave bye-bye and play pat-a-cake (average 9 months), and begin to throw objects deliberately on to the floor (average 12 months)?

When did he begin to copy you doing things about the house, such as sweeping, brushing, washing-up (average 15 months)?

When did he begin to say single words *with meaning*, not just in imitation, and not just ma ma, dadada (average 1 year), and when did he begin to join two or three words together, not just in imitation (average 21 to 24 months)?

When did he begin to tell you that he has wet himself, or wants the pottie (average 15 to 18 months)? When was he reasonably reliable by day (average two years) and by night (average 3 years)?

When could he pick up a cup, drink from it, and put it down without help and without much spilling (average 15 months)?

How long does he play with one toy (this gives an idea of his powers of concentration)?

How much does he smile and laugh (this question is asked in the case of the baby from 6 weeks to 6 months; it gives an idea of his social development)?

It is absolutely essential in all cases to determine whether the baby was prematurely born, because if he was due allowance must be made. If he was born, say, 2 months prematurely, he missed two months' development *in utero*, and so 2 months must be added to all the milestones mentioned.

The following is a summary of other important milestones of development:

- 6 weeks:** Held up with hand under abdomen, momentarily holds head in same plane as rest of body. Prone—chin momentarily off couch.
- 3 months:** Prone—chin and front of chest off couch.
- 6 months:** Sits for seconds on floor with hands forward for support. Transfers object from one hand to the other. Prone—weight on hands with extended elbows. Supine—lifts head up spontaneously.
- 8 months:** Sits securely. Stands holding furniture.
- 10 months:** Goes for objects with index finger. Pickets up pellet between tip of thumb and tip of first finger.
- 13 months:** Feeds self with spoon.
- 15 months:** Tower of 2 one-inch cubes. Creeps up stairs.
- 18 months:** Up and down stairs, holding rail. Walks up stairs, one hand held.
- 2 years:** Tower of 6 or 7 one-inch cubes. Up and down stairs 2 feet per step. Runs. Turns door-knob. Puts on shoes, socks, pants. Points to 4 parts of body. Takes off shoes, socks. Dry by night. Imitates vertical stroke and circle. Turns pages of book singly.
- 2½ years:** Tower of 8 one-inch cubes. Knows full name, sex.
- 3 years:** Stands on one leg for seconds. Rides tricycle. Tower of 9 one-inch cubes. Names one colour. Copies circle. Imitates cross. Unbuttons buttons. Counts up to 10. Counts 3 numbers, 1 in 3 trials.
- 4 years:** Names 2 to 3 colours. Copies cross. Dresses and undresses fully, no help, except shoes-laces. Can button clothes. Repeats 3 figures, in each of 3 trials.
- 5 years:** Names 4 colours. Repeats 4 figures, in 2 out of 3 trials. Copies triangle. Can manage shoe-laces. Gives age.

Notes

- (i) The only equipment needed is 10 one-inch cubes.
- (ii) The term "copies" a cross means that the child is given a pre-drawn cross and asked to copy it. The term "imitates" means that the examiner draws it when the child is looking, and is asked to do the same.
- (iii) The figures are spoken slowly and clearly, and the child is asked to repeat them.

Examination

The examination must include a general physical examination, including examination of the central nervous system. It includes measurement of the maximum head circumference, examination of the hips for congenital subluxation, a rough test of hearing (after the age of 3 months), and testing of the urine for phenylpyruvic acid.

After the age of 5 months, one begins by placing a one-inch cube in front of the child and observing the hand movements. This test will show spasticity or ataxia or athetosis. When a baby is 10 months old or thereabouts one makes a very small pellet of paper and places it in front of him to test for the index-finger approach and for finger-thumb apposition.

In all children one examines the hips by flexing them to a right angle and abducting them. Abduction is limited in congenital subluxation or the spastic form of cerebral palsy.

In all children one tests the knee-jerks, beginning to tap over the front of the foot. A brisk response there suggests an exaggerated jerk. Experience alone will tell one whether it is within normal limits or not. One tests for ankle clonus. After a year one observes the child walking or trying to walk. After the third birthday one asks him to stand on one foot.

The maximum head circumference should be recorded for every child in every well-baby clinic, out-patient practice, or home. It gives invaluable information. The head size reflects the growth of the brain, so that the head is small if the brain is not growing properly. It is too large in the case of hydro-

cephalus. Serial readings against a chart (see Fig.) are more important than single readings in that they show the rate of increase.

Child with microcephaly. The middle line shows the 50th percentile—the average head circumference. Only 10% have a head as large as that shown by the upper line (the 90th percentile), and 90% have a head circumference larger than that shown by the lower line (the 10th percentile). The three lines show the rate of increase of head size. The crosses show the head circumference of a child with microcephaly, and indicate that the head is not growing as rapidly as it should do.

It must be borne in mind that a small child must be expected to have a smaller head than a big child, and vice versa, so that the head size must be related to the size of the baby as a whole. As babies take after their parents, an unusual head should make one look at the head of the mother and father.

After three months a rough hearing test, by the crinkling of tissue paper, should be part of the routine examination of all infants. In all babies phenylketonuria is eliminated by the Phenistix test.

Conclusion

There is no short cut to developmental diagnosis. Each child can only be assessed on the basis of the history, a full physical and developmental examination, and a careful appraisal and interpretation of one's findings. There is no place for spot diagnosis. Spot diagnoses in this subject do not impress parents, who feel that only a careful examination can lead to a proper opinion. One may be right in the case of a mongol, but a full detailed history and examination of a mongol will convince parents of the rightness of the verdict much more than a spot diagnosis will. One must refuse to allow the attractive appearance of a child to lead one to a diagnosis of mental superiority. One must be sure that one has seen the maximum performance of which the child is capable. It is easy to be misled by the failure of a child to co-operate.

Developmental diagnosis is not easy, and anyone who thinks that it is easy shows clearly that he does not understand it.

The limitations and difficulties must be clearly understood. I have described these, and the methods of testing, in detail elsewhere.¹ All one can hope to do is to say how far the child

has developed at his age. If due allowance is then made for factors which may have affected his development, or which may affect it in future, one can say much about his developmental potential. It is unlikely that one will be able to recognize mental superiority. One can certainly detect mental subnormality, and one can readily diagnose any but the milder forms of cerebral palsy. One may be able to say something

about what talents he has. One will never be able to say what he will do with them.

REFERENCE

- ¹ Illingworth, R. S., *Development of the Infant and Young Child, Normal and Abnormal*, 1963, 2nd ed. Livingstone, Edinburgh.

TO-DAY'S DRUGS

Prophylactic Iron and Folic Acid in Pregnancy

Iron

Normal Iron Balance

Normal healthy adult men and post-menopausal women do not become iron deficient when their diets contain 12 mg. or more of food iron per day. They absorb sufficient iron from this diet to meet the normal iron loss in urine, faeces, and from desquamating skin of 0.5 to 1 mg. each day. Thus from an average European diet containing 12–18 mg. of iron we absorb approximately 5–10% of the iron.

Menstruation imposes a further loss of about 20 mg. per month on the average (0.7 mg. each day), but losses vary from 2.3 to 79 mg.

Iron Requirements in Normal Pregnancy

The newborn infant contains about 280 mg. of iron (range 201–372 mg.); the placenta contains 75 mg. of iron (range 34–170 mg.); expansion of the mother's red-cell mass in pregnancy demands a further 200 mg. of iron; blood loss at delivery (primiparae, about 500 ml.; multiparae, about 300 ml.) results in a further loss of 130 to 200 mg. of iron; and lactation over six months may also result in the loss of 100–180 mg. of iron. Thus the net cost to the mother in terms of iron can vary from 550–950 mg. of iron. This will be offset to some extent by the amenorrhoea and the post-partum reduction of red-cell mass, so that there is a conservation of 380 mg. (340–580 mg.).

In terms of the amount of iron that must be absorbed each day, this is 0.6 mg. in the first trimester, rising to 2.8 mg. in the second and 3.5 mg. each day in the latter third of pregnancy. Perhaps it is as well to add that in pregnancy foetal requirements for iron are always met irrespective of the state of maternal iron stores.

Consequences of Increased Iron Requirement in Pregnancy

The pregnant subject attempts to meet the increased iron requirement by mobilizing her normal iron stores and by increasing the proportion of iron absorbed from food. Thus the proportion of a dose of ⁵⁹Fe as the ferrous salt that is absorbed during pregnancy increases from about 10% of the oral dose in early pregnancy to 30% at the 20th week and to over 40% after the 30th week of pregnancy, and this increased absorption is still present even if these subjects are given 1,000 mg. of parenteral iron in early pregnancy. However, observations with iron salts are not readily transferable to food iron, and the extent to which the pregnant subject is able to mobilize food iron is uncertain. What is certain is that a very high proportion of

pregnant subjects fail to meet the demands for iron from their usual dietary intake.

Under normal circumstances about 65% of the body iron is present in the red cells as haemoglobin (2,100–2,500 mg.) and 30% (800–1,500 mg.) is present as haemosiderin in the reticulo-endothelial cells of liver and marrow. This constitutes the iron stores. It is probable that in women iron stores are considerably less than this figure. Such iron stores are demonstrable by staining a film obtained from a sternal or iliac-crest marrow aspirate by Perls' prussian blue method, which stains the haemosiderin aggregates a deep greenish-blue colour. In the latter half of pregnancy demonstrable marrow iron stores are absent in about 80% of pregnant subjects.

Once the iron stores are mobilized the serum-iron level will fall. Many observers have shown that the average plasma iron is decreased in pregnancy and that the average transferrin level is increased, this being the usual response of the iron-carrying protein to iron deficiency.

Normal pregnancy is associated with a fall in the haemoglobin concentration, the maximum fall being reached at about 25–30 weeks of pregnancy. This is due to an increase in plasma volume which is greater in extent than the increase in the red-cell mass. Although there are many excellent studies demonstrating these volume changes it is probable that in most patients an inadequate supply of iron was a limiting factor in the expansion of the red-cell mass. In other words, steps were not taken to ensure adequate iron stores in early pregnancy. For this reason it is uncertain what fall in haemoglobin concentration should be regarded as physiological in pregnancy. Pending better information on this subject 12 g. per 100 ml. is a reasonable lower limit for the haemoglobin level at this period of pregnancy. After about the thirtieth week of pregnancy the haemoglobin concentration increases. Nevertheless, in the absence of additional sources of iron a high proportion of pregnant women will fail to regain their initial haemoglobin concentrations at the end of pregnancy. In the comprehensive study of Magee and Milligan¹ the mean haemoglobin concentration of early pregnancy had not been regained one year after delivery.

But the most decisive data indicating the need for additional iron in pregnancy comes from comparison of groups of pregnant subjects given supplemental iron with those not supplemented in this way. In such studies subjects not receiving additional iron approach the end of pregnancy with haemoglobin concentrations approximately 1 g. per 100 ml. lower than those taking an additional form of iron. Many patients receiving iron have higher haemoglobin concentrations at the end of pregnancy than in the beginning.

An old viewpoint which has recently been revived² is that iron had a pharmacological effect in raising haemoglobin levels other than its effect in meeting normal iron requirements. The work supporting this hypothesis³ has been criticized.⁴ Furthermore, in another study the administration of 1,000 mg. of parenteral iron to normal subjects was not followed by any