questioning. Dominian and Dobson (1969), in their study of patients from a coronary care unit, felt that oxygen masks were a source of distress to the patients, because of their association with serious illness as well as the discomfort, but this would not explain our finding that more patients were worried by oxygen therapy than mechanical ventilation. The thirstiness mentioned by a few of our patients was probably mainly due to restricted fluid intake allowance to the postoperative cardiac patients, but if inadequately humidified gases are administered this will also contribute.

Most of the patients were pleased to return to their own wards, as they took this as a sign of recovery. A few did not want to leave because they found it a very happy ward and one because she liked having a room to herself. Four patients thought the ward unpleasant but most found it pleasant, and several comments were added about the efficiency of the staff, the reassurance of never being left alone, and the pleasant atmosphere.

I am indebted to Dr. A. M. Hutton, director of the department of anaesthetics, for his help and encouragement in undertaking this survey.

REFERENCES

- Bevan, J. R. (1969). *Lancet*, **1**, 1310.
 Blachly, P. H., and Starr, A. (1964). *American Journal of Psychiatry*, **121**, 371.
 Both Ends of the Stethoscope (1969). *Lancet*, **2**, 262.
 Cole, L. B., Youngman, H. R., and Gandy, A. P. (1968). *Lancet*, **2**, 567.
 Dominian, J., and Dobson, M. (1969). *British Medical Journal*, **4**, 795.
 Druss, R. G., and Kornfeld, D. S. (1967). *Journal of the American Medical Association*, **201**, 291.
 Egerton, N., and Kay, J. H. (1964). *British Journal of Psychiatry*, **110**, 433.
 Fahy, A., Watson, B. G., and Marshall, M. (1969). *British Journal of Anaesthesia*, **41**, 439.
 Klein, R. F., Kliner, V. A., Zipes, D. P., Troyer, W. G., and Wallace, A. G. (1968). *Archives of Internal Medicine*, **122**, 104.
 Kornfeld, D. S. (1969a). *British Medical Journal*, **1**, 108.
 Kornfeld, D. S. (1969b). *Anesthesiology*, **31**, 166.
 Kornfeld, D. S., Zimberg, S., and Malm, J. R. (1965). *New England Journal of Medicine*, **273**, 287.
 Lazarus, H. R., and Hagens, J. H. (1968). *American Journal of Psychiatry*, **124**, 1190.
 Parker, D. L., and Hodge, J. R. (1967). *Journal of American Medical Association*, **201**, 702.
 Personal Paper (1969). *Lancet*, **2**, 1129.
 Sgroi, D., Holland, J., and Marwit, S. J. (1968). Psychological Reactions to Catastrophic Illness: A Comparison of Patients Treated in an Intensive Care Unit and a Medical Ward. Presented at Annual Meeting of Psychosomatic Society, Boston, Massachusetts, March 1968.
 Watts, G. T. (1969). *Queen's Medical Magazine*, **60**, No. 4, p. 8.

GENERAL PRACTICE OBSERVED

The Practice Nurse—A New Look

J. WESTON SMITH,* M.B., D.R.C.O.G., M.R.C.G.P. ; J. B. O'DONOVAN,* M.B., B.S., M.R.C.G.P.

British Medical Journal, 1970, **4**, 673-677

Summary: A nurse has been employed by three general practitioners in a semi-rural practice of 9,000 patients since April 1968. Of her 1,360 home visits, 838 were primary visits, of which only 87 required follow-up by the doctor and 484 were requested directly to the nurse. More advanced procedures carried out by the nurse included electrocardiography, pregnancy diagnosis, blood sampling, and cytology. This study suggests that the work of a practice nurse improves the medical service already provided and reduces the work-load of the doctors, particularly by primary screening by the nurse both at home and in the surgery.

Introduction

Over the past few years there has been a great increase in the numbers of nurses included in the general practice team, either privately employed or seconded by the local authority. The nurse has largely continued to fulfil her traditional roles in the community, though it is possible to extend her role in the practice team to follow-up visiting and primary assessment (Hasler *et al.*, 1968; Marsh, 1967; Royal College of General Practitioners, 1968a). In a survey in south-west England (Royal College of General Practitioners, 1968b) it was suggested that "a great deal of experimentation and a major change of attitude amongst doctors is necessary before full use of nursing skills becomes possible in family practice." There is little evidence of such a major change.

We show that the nurse has a definite part to play as the agent of first contact. We made a preliminary examination of this (Smith and Mottram, 1967), while the present study ex-

amines the situation in greater detail, concluding that the nurses' work in primary medical care is safe and is acceptable to patients.

Study

The practice, which is semi-rural and of about 90 square miles (233 sq. km.) in area, has about 9,000 patients and three partners—one senior (Dr. A, aged 60) and two juniors (Drs. B and C, aged 40-50). The practice nurse works about 30 hours weekly between 9 a.m. and 5 p.m. She is employed by the partners and uses a practice car for visiting. In 12 months from April 1968 we investigated the effect of employing a nurse for primary screening by visiting and revisiting selected patients; the range of work possible both in the home and in the surgery, provided that suitable equipment was made available; and the reaction of patients to the service.

Methods of Recording

All face-to-face contacts between the nurse and patients were recorded on forms according to a code, particularly those findings relating to home visits. Similar records of the doctors' contact with patients in the home and surgery were kept for 12 months from May 1968 (Table I). The number of consultations and visits is lower than the national norm, probably owing to two factors:

- (1) We also work in a local general-practitioner hospital with 68 beds and a busy casualty department; some of the work normally carried out in a practice—such as minor surgery, and treatment of minor trauma and septic conditions—is dealt with there.

* General Practitioner, Tamworth, Staffs.

- (2) The fact that a nurse is part of a practice team and is an acceptable alternative to the doctor on certain occasions probably reduces the doctors' work-load.

TABLE I.—Total Number of Doctor-Patient Consultations and Visits and Distribution of Patients by Age and Sex

Doctor	No. of Consultations	No. of Visits	Consultations				Ratio of Males to Females
			With Patients Aged Under 10 Years		With Patients Aged Over 60		
			No.	%	No.	%	
A* ..	7,572	1,400	365	6.5	1,865	33.1	2:1
B ..	6,677	251	1,272	19.2	295	4.4	2:3.5
C ..	7,666	791	1,272	16.5	890	11.5	2:3.5

*Figures for eight months only (October 1968—May 1969).

Nurse's Duties

To analyse the nurse's duties, a record card was kept for each patient, and all the necessary nursing services recorded on it. In addition to name, age, marital status, and a rough diagnosis, we recorded the source of the demand for the nurse's services at home and the mode of action she adopted in each case.

Patient Contacts

In comparison with figures given by Pinsent (1968) and the Royal College of General Practitioners (1968a) employing a nurse for our type of work appeared to shift the demand and actual work away from the older age groups towards the younger (Table II). This may be due partly to district nurses

TABLE II.—Total Number Nurse-Patient Contacts and Distribution by Age and Sex

Age Group (Years)	Surgery	Home	Males		Females	
			No.	%	No.	%
0-4	988	432	762	45.6	658	16.4
5-14	341	315	269	16.1	387	9.7
15-44	2,415	228	344	20.6	2,299*	57.8
45-65	446	92	185	11.0	352	8.8
>65	106	293	112	6.7	288	7.2
Total	4,296	1,360	1,672	100	3,984	100

*1,408 antenatal visits.

retaining their traditional role of caring for the elderly sick; an examination of the work done by nurses in other surveys seems to substantiate this.

Home Visiting

Origin of Request

Significantly 55% of the demands for the nurse's services for patients under the age of 14 were made directly to her. On the other hand, the older practitioner made much less use of the nurse for domiciliary services—about 1.6% of the total, except in the older age groups (Table III). Of the 3,807 visits

TABLE III.—Origin of Request for Nursing Service at Home

	Age Group (Years)					Total
	0-4	5-14	15-44	45-64	≥65	
First visit request by:						
Dr. A	7	3	1	1	10	22
Dr. B	44	40	52	9	12	157
Dr. C	59	40	52	12	12	175
Revisit for:						
Dr. A	6	1	3	2	5	17
Dr. B	17	9	9	5	9	49
Dr. C	20	8	35	35	34	132
Routine revisit	19	17	13	24	198	271
First request for nurse	238	187	44	2	13	484
Independent request at home	10	4	7	2	1	24
Not known	12	5	12			

in the practice the nurse carried out 1,360, of which 838 were first visits. This meant a great saving of time, since one visit takes 16-20 minutes, of which only a fraction is spent with the patient. Since the nurse has been employed requests for her services have steadily increased, so that visiting is now an established part of her work.

The most common conditions requiring a visit were minor disorders of the upper respiratory tract and common infectious diseases (Table IV). Some of these 356 visits represent a new service rather than a substitute for the doctor.

TABLE IV.—Analysis of Primary Requests for Nurse*

Age Groups (Years)	0-4		5-14		15-44 >44			Total
	M.	F.	M.	F.	M.	F.	All	
Sex								
Number of Patients	100	82	65	71	6	32	15	371
Communicable disease	37	47	24	36	0	4	0	148
E.N.T. and respiratory	50	49	46	49	0	25	0	219
Gastro-intestinal disorders	20	12	4	7	0	3	3	49
Other disorders	17	6	11	10	6	6	12	68
Total	124	114	85	102	6	38	15	484

*Four of the requests were not known.

Mode of Action

In considering the action taken by the nurse on seeing the patient at home we kept to the criteria laid down in January 1967.

- (1) In the nurse's opinion it was unnecessary for the doctor to visit or for the patient to visit the doctor.
- (2) A doctor's visit to the house was unnecessary, but the patient was advised to visit the doctor.
- (3) Commonsense measures were suggested by the nurse, who, in addition, would ask the doctor to prescribe a simple placebo.
- (4) The nurse would make a provisional diagnosis if it was thought to be an epidemic condition and report it to the doctor, who would visit the patient when he was in the neighbourhood.
- (5) If pyrexia with other disquieting symptoms were present the nurse would advise the household that the doctor would be calling later on.
- (6) The doctor was contacted urgently.

During the time the scheme has operated we have modified it slightly so that, during an epidemic instead of the doctor visiting when in the neighbourhood, the family were instructed to contact the practice again immediately if they had cause for concern and were given information about the expected course of the condition (Table V). In only 87 of the 838 primary visits was a follow-up visit by the doctor indicated; the need to visit the same day was for very young children, whereas elderly people needed more "when convenient" visits.

TABLE V.—Mode of Action by Nurse at Home

	Age Group (Years)					Total
	0-4	5-14	15-44	45-64	≥65	
(1) No action	6	1	0	1	14	22
(2) Patient to visit doctor later	8	15	23	3	7	56
(3) General advice	188	171	86	21	28	494
(4) Medicine to be prescribed by doctor	170	109	49	13	54	395
(5) Doctor to visit when in neighbourhood	0	0	2	5	28	35
(6) Doctor to visit that day	13	9	6	1	8	37
(7) Doctor contacted urgently	2	6	7	0	0	15
(8) Other services	7	11	19	1	33	71
(9) Nurse to call again	4	6	6	7	9	32
Services, including E.C.G., blood sample, immunizations, dressings, etc.	23		30	38	112	203

Of the 484 requests for primary visits made directly to the nurse 25 patients subsequently had to see the doctor. This

contrasts with the 354 primary visits given to the nurse by the doctors, which necessitated 63 (18%) personal visits by a doctor (Table III).

Surgery Duties

Formerly, when patients arrived without an appointment it was always a matter of anxiety whether they needed to be seen immediately. On finding the doctors fully booked, patients now ask to see the nurse, who can then assess the urgency of the request. We had not expected this demand, so no figures have been recorded during the study period.

During the study period the amount of work covered by the nurse proved greater than previously envisaged. At first her function was to assess the degree of illness and help the doctor to decide the urgency and place of consultation. In times of epidemic, however, she did act as diagnostician and was able to advise, especially when to contact the doctor.

Advanced Procedures

Since the number of advanced procedures carried out by the nurse had increased, we analysed a four-week period during August and November 1969 and January and March 1970. (Tables VI and VII). The results suggested that the demand for this type of service greatly increases as its availability becomes known. We believe some of the procedures are better carried out by a nurse, while others, being technical skills—such as, electrocardiography, Pregnosticon, routine blood

TABLE VI.—*Mode of Action of Nurse in Surgery*

	Age Group (Years)					Total
	0-4	5-14	15-44	45-64	≥65	
Immunization and vaccination	865	143	92	19	27	1,146
Weight-reducing advice	7	46	455*	222*	21*	751
Desensitization	2	1	1			4
Skin testing, E.C.G., etc.	1	0	14	9	10	34
Injections and blood sample	12	34	240*	85	53	424
Ear syringe	2	3	59	20	9	93
History, temperature, blood pressure, and swab	15	11	30	53	10	119
Urine analysis	3	6	32	1	2	44
Dressings	13	18	60	18	69	178
Antenatal	0	10	1,398	0	0	1,408

*High preponderance of females.

Antenatal-urine analysis, blood pressure, and weighing only.

TABLE VII.—*Further Analysis of More Advanced Procedures*

	Survey Year	Four Sample Weeks 1969-70
E.C.G.	50*	8
Pregnancy diagnosis	26	12
Blood sample	17†	25
Cytology	37	16

*High preponderance of males. †High preponderance of females.

pressure readings—mean that one person doing them all the time becomes adept and the equipment is kept in good order. Some of these skills were acquired after we started the scheme.

The number of routine dressings and attention to minor trauma is probably lower than in most practices, because of our link with the local casualty department. Similarly, we are careful to restrict the nurse's activities in the more traditional fields associated with district nursing.

Patients' Reactions

The Royal College of General Practitioners (1968a, p. 21) states "One of the conclusions of this study may well be that

the feelings and attitudes of the doctors and nurses and of the patients if they can be assessed seem likely to be much more valuable in evaluating the success of the experiment than the measures of work load and the time saved."

We hoped that the introduction of a practice nurse would provide a better service for our patients and give the doctors more time for consultations requiring their greater skills and other aspects of their work, including hospital duties. Since patients, however, might assume the nurse was a "substitute for the doctor" we sent out a preliminary questionnaire to patients who had been seen by the nurse. Of these only 0.5% were dissatisfied (70% replied, though return postage was not prepaid). A further questionnaire was sent to 150 patients, who were selected at random from three groups, but not matched for age, sex, etc.: patients who had been visited by the nurse at home; patients who saw the nurse at the surgery; and patients who had never seen the nurse. A prepaid envelope was also enclosed, and 108 replies were received (see Questionnaire).

Copy of Questionnaire

We should like you to indicate who you would prefer to do the following things relating to your medical care			
	Doctor	Nurse	Either
Change a dressing			
Syringe your ears			
Give an injection			
Examine your throat			
Take your blood pressure			
Give diet instruction			
Decide what medicine you need			
Explain what is wrong with you			
Vaccinate your baby			
Take your medical history			
Discuss your menstrual irregularities			
Decide whether x-ray examination is needed			
Decide if blood examination is necessary			
Discuss family difficulties			
Tell your hidden fears to			
Decide how ill your child is			
Decide whether it is safe to take your child out			

The answers to the questionnaires were divided into three groups (Figs. 1, 2, and 3): (1) no change in patients' attitudes (in other words, these patients continued to prefer the doctor to decide the issue); (2) increased acceptance of the nurse (these were regarded as proper duties for the nurse to perform); (3) shift to accepting the nurse once the patients had met her and understood what she was trying to do.

Fig. 1.—Doctor preference.

FIG. 2.—Increase in nurse acceptance.

FIG. 3.—Shift to nurse acceptance.

Discussion

In a particular practice with particular personnel we feel it is possible to provide an acceptable and useful nursing service. Nevertheless, two questions arise: is this possible in other practices—in other words, is it of wider application? Secondly is it desirable from both the patients' and the professions' (medical and nursing) points of view? If the answers are yes, then the problem is how to put such a service into action.

Personal selection and deployment of the nurse are essential. Despite the absence of formal training for this particular role she has acquired the necessary skills and attitudes and become a well-integrated member of the practice team.

Patients are now aware of the service she provides and use it with confidence. Moreover, we think that a new type of practice nurse, suitably trained, should replace practice nurses, district nurses, and health visitors as we know them.

Furthermore, our findings show that there is a place for primary screening by a nurse; this is contrary to the opinions of some writers (Marsh, 1969), but, provided the service is introduced properly, patients soon feel that it is an extra service rather than a substitute for doctor care. Williams (1967) suggested that this had introduced a new concept regarding the National Health Service; it may well be what the Service needs.

Lay people seem to appreciate our concept of the practice nurse, while the medical and nursing professions tend to be hostile. The question of legal responsibility has yet to be resolved, but, provided the nurse is able to undertake the tasks given to her and every action is fully documented and agreed, all reasonable steps have been taken to guarantee a proper standard of care for the patient.

It is very difficult to assess accurately the effect of this service on the doctors' work-load. The Royal College of General Practitioners' (1968a) survey estimated the saving of time to be about 4–8%, but other writers have found it almost impossible to assess. We certainly do less visiting, but the number of patients seen does not appear to decrease, and consultation times are longer. The quality of the work done by doctors and the effect on their lives are noticeably different. Furthermore, there is less frustration when a nurse is used intelligently in the practice team, probably because the doctor feels that his time is not being misused by doing work that another trained person can do. On the other hand, using the more advanced skills she has been taught in her nursing school and learning new ones, is an impetus and encouragement to the nurse.

Official Recognition

In view of the interest shown and the number of doctors who are starting similar schemes, official recognition must be given soon by both professions and the Ministry. Certain practices should be recognized for training purposes and, by their organization and methods, show that they are capable of teaching suitable candidates. The service must not be used for cheap medicine or as a convenient way of providing more free time for the medical profession.

About two and a half hours of work are spent each week per thousand patients using a nurse in the manner we describe, but the demand is increasing and we are placing another nurse in the team. There must surely be reserves of State-registered nurses whose skills are unused owing to family commitments and who would be unable to fit in with the 24-hour service expected by other sections of the Health Service. The hours of work are suitable (about 9 a.m. to 5 p.m.) for the married woman with a family who wants to come back to nursing.

In discussing who should be agent of first contact, Anderson (1969) suggested that with multiple access a paramedical worker should act as the agent of primary medical care parallel with the doctor. Some patients may elect to consult the general practitioner, others to consult the nurse or the social worker in the first instance. At the moment the medical and nursing professions are still uncertain of their role in the provision of community services. Possibly a team such as we have described may help to solve many problems of the National Health Service.

We appreciate the interest shown by members of the medical and nursing divisions of the Ministry of Health and Social Security. We should also like to thank the directors and staff of the Royal College of General Practitioners Research Unit in Birmingham for their advice and assistance with data-processing. We would like to thank Sister E. M. Mottram, without whose ready and patient help this study would not have been possible. We acknowledge our

debt of gratitude to Beryl Weston Smith, whose assistance in voluntarily recording the figures, designing the graphs, and giving helpful criticism and advice in setting out the paper contributed to the success of the project.

REFERENCES

- Anderson, J. A. D. (1967). *Lancet*, 2, 679.
 Hasler, J. C., et al. (1968). *British Medical Journal*, 3, 734.
 Marsh, G. N. (1967). *British Medical Journal*, 1, 489.
 Marsh, G. N. (1969). *British Medical Journal*, 4, 42.

- Pinsent, R. J. F. H. (1968). *Journal of the Royal College of General Practitioners*, 15, 402.
 Royal College of General Practitioners (1968a). *The Practice Nurse*, London, Royal College of General Practitioners.
 Royal College of General Practitioners (1968b). *General Practice in South West England*, London, Royal College of General Practitioners.
 Smith, J. W., and Mottram, E. M. (1967). *British Medical Journal*, 4, 672.
 Smith, J. W., and O'Donovan, J. B. (1970). *Journal of the Royal College of General Practitioners*. In press.
 Williams, D. L. (1967). *British Medical Journal*, 4, 803.

NEW APPLIANCES

Fine Continuous-feed Aneurysm Needle

Mr. BRIAN L. DOWLING, surgical tutor, Nuffield Department of Surgery, Radcliffe Infirmary, Oxford, writes: The instrument described here is based on the continuous-feed aneurysm needle of Tibbs (1952). In order to facilitate the ligation of small vessels the instrument is constructed in the form of a traditional aneurysm needle (Fig. 1). The "needle" component of the device is at right-angles to its main axis and permits easy use in a deep wound. The needle is hollow (Fig. 2), allowing free flow of the suture material without "bow-stringing."

This material can be compressed by the thumb of the right hand against the grip at the top of the shaft, allowing ligation by the one-hand technique using the left hand. The spool is contained within the handle (Fig. 3) and is large enough to hold suture material for even a long operation. The material used has generally been 00 silk, but loading could be done with any suitable material.

The instrument has been in use for 18 months and has proved invaluable. It has been particularly useful in the ligation of small mesenteric vessels in the preparation of ileostomies, Roux loops, and interposed jejunal loops. It has also proved of value in biliary and gastric surgery, and has been used with success in ligating the small vessels encountered during partial thyroidectomy. A continuous-feed aneurysm needle of this type avoids the problems of using artery forceps on small delicate vessels and provides a rapid method of repeating ligation.

The prototype instrument was made by Mr. Don Hopkins in the workshops of the Nuffield Department of Surgery. Arrangements are being made for its manufacture by Messrs. Charles Thackray of Leeds.

REFERENCE

- Tibbs, D. J. (1952). *British Medical Journal*, 2, 1201.

FIG. 1.—Complete aneurysm needle threaded for use. FIG. 2.—Tip of the instrument, showing the silk passing through the hollow curved needle and eye. FIG. 3.—Handle, unscrewed, showing the spool on its guide.