Published in final edited form as:

Invest Ophthalmol Vis Sci. 2006 June ; 47(6): 2336–2340.

A Prospective Assessment of the Y402H Variant in Complement Factor H, Genetic Variants in C-reactive Protein, and Risk of Age-Related Macular Degeneration in Men

Debra A. Schaumberg, ScD, OD, MPH,

Division of Preventive Medicine, Children's Hospital; Schepens Eye Research Institute, Department of Ophthalmology, Harvard Medical School, Boston

William G. Christen, ScD,

Division of Preventive Medicine, Department of Ophthalmology, Harvard Medical School, Boston

Piotr Kozlowski, PhD,

Division of Hematology, Department of Ophthalmology, Harvard Medical School, Boston

David T. Miller, MD, PhD,

Division of Hematology, Brigham and Women's Hospital; Division of Genetics, Department of Ophthalmology, Harvard Medical School, Boston

Paul M. Ridker, MD, MPH, and

Division of Preventive Medicine, Center for Cardiovascular Disease Prevention, Department of Ophthalmology, Harvard Medical School, Boston

Robert YL Zee, PhD

Division of Preventive Medicine, Center for Cardiovascular Disease Prevention, Department of Ophthalmology, Harvard Medical School, Boston

Abstract

Background: Two biologically related factors, complement factor H (CFH) and C-reactive protein (CRP) have been associated with age-related macular degeneration (AMD). The Y402H variant of CFH is located within the binding site of CFH for CRP. Although plasma CRP levels have been related to AMD, and plasma CRP levels are partly determined by genetic variation, there is no information on whether genetic variants in *CRP* are associated with AMD.

Methods: We performed a prospective analysis among 111 men who eventually developed AMD and 401 men who remained free of AMD, all participants in the Physicians' Health Study. We determined genotypes for the common T>C single nucleotide polymorphism (SNP) in exon 9 of *CFH* (rs1061170; protein Y402H) as well as seven previously described *CRP* SNPs (rs3093059, rs2794521, rs3091244, rs1417938, rs1800947, rs1130864, rs1205). We used logistic regression analysis to evaluate individual SNPs as well as six *CRP* haplotypes for association with AMD.

Results: The high-risk C allele of *CFH* was present in 45% of cases and 34% of controls. We observed an odds ratio for AMD of 1.46 (95% CI=1.05-2.04) for TC heterozygotes and 2.13 (95% CI=1.10-4.16) for CC homozygotes assuming a multiplicative (log-additive) model, and calculated an attributable fraction of 25% (95% CI=1% to 44%). For *CRP*, single-marker or haplotype-based analysis failed to reveal any significant associations with risk of AMD.

Conclusion: These prospective data confirm an association between the Y402H variant of *CFH* and risk of AMD. In contrast, although biologically plausible, genetic variation in *CRP* does not appear to be associated with risk of AMD. Further prospective study of larger numbers of subjects is needed to substantiate available information on the genetic epidemiology of AMD.

Introduction

A body of recent work suggests that aging may be intimately related to chronic low-grade inflammation, resulting from the body's response to accumulated stress from myriad stressors such as UV radiation, oxidation, and infection. This chronic stress leads to the development, at varying rates among individuals, of a subclinical chronic inflammatory response (1). The level of this inflammatory response predicts the development of several age-related degenerative diseases (2), including age-related macular degeneration (AMD) (3), the leading cause of blindness and visual impairment among aging populations.

Consistent with a pathogenic role for inflammation, recent work has identified a strong association between a common variant of the gene for complement factor H (*CFH*) (dbSNP: rs1061170, sequence: T1277C, protein: Y402H) and the risk of age-related macular degeneration (AMD) (4-10). This association is plausibly causative as the *CFH* gene maps to a location on chromosome 1q termed the *ARMD1* locus that was linked with AMD in several studies, relevant biological mechanisms can be theorized, and its significance has now been consistently replicated in a number of studies. However, the magnitude of prior estimates, with relative risks in the range of 2.5 to 7.4, must be interpreted in light of the study design and selected nature of cases as well as controls, including, for example, clinic-based ascertainment of prevalent cases in most instances as well as the restriction in some studies that controls had no family history of AMD (6), all of which might bias risk estimates upward (11). Moreover, although *CFH* (or a tightly linked gene) appears to play a role in AMD, the high prevalence of the risk allele in the population suggests a stochastic effect or the existence of interacting risk factors.

In addition, plasma levels of C-reactive protein (CRP), an acute phase reactant and systemic marker of subclinical inflammation, have been previously found to be associated with risk of age-related macular degeneration (AMD) (12,13). *CFH* interacts with high affinity to CRP to reduce deposition of the terminal attack complex of complement (C5b-9), and the presumed causal Y402H variant of *CFH* is predicted to lead to changes in the activities of the binding sites on *CFH* for CRP that could alter the ability of *CFH* to suppress complement-mediated damage (14). Recent work has shown that variation in plasma CRP is at least partly determined by genetic variation in the gene for CRP (15,16). However, there is no information on whether genetic variants in *CRP* are associated with AMD.

We therefore investigated the association between the Y402H variant in *CFH* and seven common variants in *CRP* and AMD in a prospective study of a subset of subjects from the Physicians' Health Study (PHS), a cohort of 22,071 initially healthy male physicians.

Materials and Methods

Study Population

We conducted a prospective study among a subset of participants in the PHS cohort, a randomized, double-masked, placebo-controlled trial of aspirin and beta-carotene for the primary prevention of cardiovascular disease and cancer among 22,071 US male physicians that began in 1982 and has been conducted in accordance with the Declaration of Helsinki. At baseline, 14,916 PHS participants provided a blood sample. Among these subjects, 1712 were selected for participation in a nested case-control study of the genetic epidemiology of

cardiovascular disease. Each cardiovascular disease case was matched to one control chosen randomly from participating physicians who reported no cardiovascular disease at the time the case reported his event, and met matching criteria of age, smoking habit, and time since randomization. For the present study, we excluded 88 non-Caucasian men, 69 men for whom relevant genotype data were not available, 17 men who reported a diagnosis of AMD before baseline, 27 men with a diagnosis of deep vein thrombosis. Among the remaining men, we identified 111 Caucasian participants with a confirmed diagnosis of incident AMD. For the control group, we excluded 999 men who were ≤ 65 years old leaving 401 Caucasian men available to serve as controls. The availability of *CFH* and *CRP* genotype data among this subset of PHS participants provided a cost-effective means to investigate whether variants of these genes are associated with AMD.

Ascertainment of AMD Cases

Procedures for documentation of incident cases of AMD have been described previously (17). Beginning with the 7-year follow-up questionnaire we asked PHS participants about the diagnosis of AMD, and updated this information on all subsequent annual follow-up questionnaires. We requested the month and year of diagnosis and the name and address of the diagnosing eye doctor, and signed permission to review medical records. Upon receipt of permission, we sent a letter to the participant's eye doctor containing a brief questionnaire to obtain information on the date of diagnosis, the best-corrected visual acuity at the time of diagnosis, and the chorio-retinal lesions that were observed at diagnosis (drusen; RPE changes including atrophy, hypertrophy and RPE detachment; geographic atrophy; subretinal neovascular membrane; disciform scar). For the present study, we included all confirmed cases of AMD whether or not visual acuity was affected.

Validation of AMD Case Ascertainment

To validate our methodology for AMD case ascertainment, we performed a validation study in the Nurses' Health Study (NHS), in which we use parallel methodology for confirmation of AMD. Briefly, we requested 30-degree color stereo photographs from cases reported on the 1990, 1992, and 1994 NHS questionnaires and subsequently confirmed by independent review of medical record information to meet our case definition. Two experienced retinal specialists with specific expertise in research on macular degeneration reviewed the slides using a magnified (4x) stereo viewer to evaluate a circle with a radius of two disc diameters from the center of the fovea. We have reviewed slides from 180 of a possible 210 eligible cases. Among these, we received 143 slide sets that were of sufficient quality to grade, 132 (92%) of which were judged to have definite AMD, and 5 (3%) to have probable AMD. Thus, 95% of medical record-confirmed cases of AMD were considered definite or probable AMD upon independent review of fundus photographs.

SNP Genotyping

CFH genotypes were determined using a fluorescence-based detection method (Applied Biosystems Assay-by-Design). Seven *CRP* SNPs were genotyped by either Sequenom matrix-assisted laser desorption and ionization-time of flight (MALDI-TOF) mass spectrometry or ABI PRISM TaqMan assays according to standard protocols as previously described (15). To confirm genotype assignments, we performed genotyping in duplicates blinded as to case-control status, and two independent observers carried out scoring. Disagreements ($<1\%$ of all scoring) were resolved by a joint reading, and if necessary, repeat genotyping. The *CRP* SNPs included: A T>C substitution at -757 (rs3093059), at C>T substitution at -717 (rs2794521), a triallelic C>T/A substitution at -286 (rs3091244), a T>A substitution at 349 (rs1417938), a G>C substitution at 1059 (rs1800947), a C>T substitution at 1444 (rs1130864), and a G>A substitution at 1846 (rs1205).

Statistical Analysis

Genotype and allele frequencies between cases and controls were compared using χ^2 -analysis. Tests for Hardy-Weinberg equilibrium (HWE) were carried out by χ^2 -analysis. Pairwise linkage disequilibrium (LD) was examined as described by Devlin and Risch (18). We used logistic regression models to estimate the odds ratios (OR) and corresponding 95% confidence intervals (CI) for the effect of genotypes on risk of AMD adjusted for other risk factors. Haplotype frequencies were estimated from genotype data using PHASE v2.1 (19,20), and the haplotype distribution between cases and controls was compared by likelihood ratio test. In addition, we examined the relationship between haplotypes and AMD outcome by logistic regression analysis using a baseline-parameterization approach (21), adjusting for age, smoking, body mass index, diabetes, prior myocardial infarction or stroke, hypertension and randomized treatment group. A two-tailed P-value of ≤ 0.05 was considered to represent a statistically significant result. All analyses were carried using SAS v9.1/Genetics. For analysis of single *CRP* SNPs, the study had 80% power to detect odds ratios ranging from 1.8 for the most prevalent SNP to 2.2 for the least prevalent SNP (22). Finally, we calculated the attributable fraction in the population and its 95% CI as a measure of the proportion of AMD associated with the *CFH* polymorphism (23).

Results

Among the 111 participants with AMD, 72 had visual acuity of 20/30 or worse due to AMD, including 21 cases with the neovascular form of the disease, and 51 cases with dry AMD. Control subjects were by design significantly older than the AMD cases (Table 1). Apart from controls also being more likely to have hypertension, the distribution of other factors such as smoking status, body mass index, history of hyperlipidemia, diabetes mellitus, use of aspirin, and family history of premature coronary artery disease were not different between AMD cases and controls. There was no evidence for departures from HWE for any of the SNPs in either cases or controls.

The high-risk C allele of *CFH* was present in 45% of cases and 34% of controls ($P=0.005$), and TC and CC genotypes were more common among cases as presented in Table 2 (P for trend=0.008). In logistic regression models for all AMD cases and controls, we observed significant association of the *CFH* polymorphism with AMD (P for trend=0.02), with an odds ratio of 1.46 (CI=1.05-2.04) for TC heterozygotes and 2.13 (CI=1.10-4.16) for CC homozygotes, assuming an additive model (such that the effect of the number of risk alleles is additive on the log-odds scale) adjusted for age, smoking, body mass index, history of hypertension, presence or absence of diabetes, randomized aspirin and beta-carotene treatment assignments, and cardiovascular disease status. In a similar model limited to cases who had visual acuity of 20/30 or worse due to AMD, the OR (CI) estimates were 1.65 (1.12-2.42) for TC heterozygotes and 2.72 (1.25-5.86) for CC homozygotes. Based on these data, the population attributable risk for *CFH* Y402H was estimated to be 25% (95% CI= 1% to 44%).

The estimated ORs were of similar magnitude among neovascular cases, though confidence intervals were wide due to the small number of cases (data not shown). In additional analyses in which we included men ≤ 65 years old in the control group estimates were also not different apart from narrower confidence intervals due to the increased sample size (data not shown).

In marker-by-marker chi-square analyses of the seven *CRP* SNPs (Table 3), none of the alleles or genotypes was significantly associated with AMD. Similarly, we did not observe any significant associations between the seven *CRP* SNPs and AMD in logistic regression models adjusted for age, smoking, and other factors, and assuming an additive model (Table 4). The *CRP* variants tested were in LD amongst each other as previously reported (15). Haplotype frequencies $\geq 1\%$ based on the seven genotyped *CRP* SNPs were not significantly different

among cases and controls, and haplotype-based logistic regression analysis did not uncover any significant associations with risk of AMD, although confidence intervals were wide (data not shown).

Discussion

Data derived from epidemiological, genetic, laboratory, and pathological studies support the theory that inflammation has a role in the initiation and progression of AMD (3-6,12). In this prospective study of a subset of AMD cases and controls from the PHS, we have observed a significant association of the Y402H variant of CFH and risk of AMD, confirming the results of prior studies. In contrast, we did not observe any significant association between common genetic variants in *CRP* and risk of AMD, though power for detection of modest associations was limited.

Participants in the present study were originally selected for a study of the genetic epidemiology of cardiovascular disease. However, any bias should be minimal given the lack of association between the Y402H variant and cardiovascular disease in this population (24), and our control for these conditions in regression models. Assessment of AMD relied on self-reports confirmed by review of medical record information (17,25, 26), so some cases of AMD may have been missed or misdiagnosed. However, in a cohort study, including in the case of nested case-control sampling from a cohort, a disease definition that has low sensitivity but high specificity does not bias the observed relative risk because there is a proportional reduction of cases in each exposure group so long as case ascertainment is unrelated to the exposure of interest (here the *CFH* and *CRP* genotypes) (27). Although we do not know the sensitivity of our case definition, our validation indicates high specificity (28). Nonetheless, because subjects were not examined, it remains possible that associations were underestimated. The likelihood of underestimation may be increased given the lower prevalence of CFH Y402H variants observed among the cases in the present study as compared to others (4-8). Given the relatively small number of cases, this observation may be due to chance, or possibly to the inclusion of a case group with milder forms of AMD.

Chronic inflammation could influence the risk of AMD through various pathways including endothelial dysfunction in choroidal vessels, development of basal deposit and drusen, and degeneration of Bruch's membrane. All of these potential points of attack would result in alterations in the transport and exchange of nutrients and waste materials between the RPE and choroid, and could theoretically cause the characteristic changes observed in AMD. Inflammatory stimuli are also known to increase the production of reactive oxygen intermediates, which are thought to play a key role in the pathogenesis of AMD (29,30). Furthermore, inflammation can reduce the bioavailability of antioxidants, setting the stage for a vicious cycle of altered redox status and increased oxidative stress (29).

The present findings confirm a number of prior reports that a common variant in *CFH* is strongly associated with risk of AMD (4-8). *CFH* plays an intimate role in the regulation of alternative pathway of complement activation, and alterations in the ability of *CFH* to suppress complement-mediated damage among people with the Y402H variant of *CFH* could increase the level of subclinical inflammation and spur the progression of AMD (3,31,32).

CFH interacts biologically with *CRP*, and *CFH* Y402H is located within the binding site on *CFH* for *CRP* (14). *CRP* is a major acute-phase reactant produced by the liver. Plasma levels of *CRP* may remain slightly elevated in states of chronic low-level inflammation that occur in a subset of otherwise healthy individuals during aging. Thus, mildly elevated levels of *CRP* appear to be a useful marker of a heightened state of inflammation or nonspecific immune reactivity. Higher *CRP* levels have previously been found to be associated with the presence and progression of advanced AMD (12,13); although not all investigators have observed a

relationship (33-35). It is plausible that CRP has some direct pathophysiological role in AMD, perhaps mediated through its capacity to induce complement activation via the alternative pathway and contribute to tissue damage through a number of complement-mediated mechanisms (36). However, the biological functions of CRP have yet to be fully elucidated, and it remains possible that plasma CRP levels are a risk marker, perhaps of a heightened state of immune reactivity or inflammatory activity, with no direct biological role in AMD pathogenesis. The triallelic SNP located at -286 from the start of transcription has been recently associated with higher plasma CRP levels among carriers of the rarer T and A alleles at this locus (16,37). However, in the present study we did not observe carriers of the rarer alleles to be more likely to develop AMD. It is possible that the association of variation within *CRP* with AMD may be influenced by variation in *CFH* or other factors, a hypothesis that the present study was not adequately powered to investigate.

These prospective data confirm the association between *CFH* Y402H and AMD. Incident cases are preferred over prevalent cases because polymorphic genetic systems are often related to a number of conditions, which could at least in theory alter the clinical course or survival of affected individuals and thus result in biased estimates of association. At the same time, prior reports suggest that *CFH* Y402H is more strongly associated with advanced disease (9), and consequently the magnitude of the association observed in any particular study is likely to be influenced by the distribution of case severity. In that regard, the present study included a majority of cases with early AMD, which may account for the more modest effect observed. In addition, the relatively small number of cases increases the chance that associations were underestimated. In contrast to *CFH*, we found no evidence that common genetic variants in *CRP* are associated with risk of AMD. Further prospective study of larger groups of subjects is needed to fully understand the interrelationships of these and other genes as well as behavioral risk factors in the development of this important cause of blindness and visual impairment.

References

1. Franceschi C, et al. Inflamm-aging. An evolutionary perspective on immunosenescence. *Ann N Y Acad Sci* 2000;908:244–54. [PubMed: 10911963]
2. Ballou SP, Kushner I. Chronic inflammation in older people: recognition, consequences, and potential intervention. *Clin Geriatr Med* 1997;13(4):653–69. [PubMed: 9354747]
3. Hageman GS, et al. An integrated hypothesis that considers drusen as biomarkers of immune-mediated processes at the RPE-Bruch's membrane interface in aging and age-related macular degeneration. *Prog Retin Eye Res* 2001;20(6):705–32. [PubMed: 11587915]
4. Klein RJ, et al. Complement factor H polymorphism in age-related macular degeneration. *Science* 2005;308(5720):385–9. [PubMed: 15761122]
5. Haines JL, et al. Complement factor H variant increases the risk of age-related macular degeneration. *Science* 2005;308(5720):419–21. [PubMed: 15761120]
6. Edwards AO, et al. Complement factor H polymorphism and age-related macular degeneration. *Science* 2005;308(5720):421–4. [PubMed: 15761121]
7. Zarepari S, et al. Strong association of the Y402H variant in complement factor H at 1q32 with susceptibility to age-related macular degeneration. *Am J Hum Genet* 2005;77(1):149–53. [PubMed: 15895326]
8. Hageman GS, et al. A common haplotype in the complement regulatory gene factor H (HF1/CFH) predisposes individuals to age-related macular degeneration. *Proc Natl Acad Sci U S A* 2005;102(20):7227–32. [PubMed: 15870199]
9. Rivera A, et al. Hypothetical LOC387715 is a second major susceptibility gene for age-related macular degeneration contributing independently from complement factor H to disease risk. *Hum Mol Genet*. 2005

10. Conley YP, et al. Candidate gene analysis suggests a role for fatty acid biosynthesis and regulation of the complement system in the etiology of age-related maculopathy. *Hum Mol Genet* 2005;14(14): 1991–2002. [PubMed: 15930014]
11. Lohmueller KE, et al. Meta-analysis of genetic association studies supports a contribution of common variants to susceptibility to common disease. *Nat Genet* 2003;33(2):177–82. [PubMed: 12524541]
12. Seddon JM, et al. Progression of age-related macular degeneration: prospective assessment of C-reactive protein, interleukin 6, and other cardiovascular biomarkers. *Arch Ophthalmol* 2005;123(6): 774–82. [PubMed: 15955978]
13. Seddon JM, et al. Association between C-reactive protein and age-related macular degeneration. *JAMA* 2004;291(6):704–10. [PubMed: 14871913]
14. Giannakis E, et al. A common site within factor H SCR 7 responsible for binding heparin, C-reactive protein and streptococcal M protein. *Eur J Immunol* 2003;33(4):962–9. [PubMed: 12672062]
15. Miller DT, et al. Association of common CRP gene variants with CRP levels and cardiovascular events. *Ann Hum Genet* 2005;69(0):623–638. [PubMed: 16266402]
16. Carlson CS, et al. Polymorphisms within the C-reactive protein (CRP) promoter region are associated with plasma CRP levels. *Am J Hum Genet* 2005;77(1):64–77. [PubMed: 15897982]
17. Christen WG, et al. A prospective study of cigarette smoking and risk of age-related macular degeneration in men. *JAMA* 1996;276(14):1147–51. [PubMed: 8827967]
18. Devlin B, Risch N. A comparison of linkage disequilibrium measures for fine-scale mapping. *Genomics* 1995;29(2):311–22. [PubMed: 8666377]
19. Stephens M, Smith NJ, Donnelly P. A new statistical method for haplotype reconstruction from population data. *Am J Hum Genet* 2001;68(4):978–89. [PubMed: 11254454]
20. Stephens M, Donnelly P. A comparison of bayesian methods for haplotype reconstruction from population genotype data. *Am J Hum Genet* 2003;73(5):1162–9. [PubMed: 14574645]
21. Wallenstein S, Hodge SE, Weston A. Logistic regression model for analyzing extended haplotype data. *Genet Epidemiol* 1998;15(2):173–81. [PubMed: 9554554]
22. Schlesselman, JJ. Case-Control Studies: Design, Conduct, Analysis. Oxford University Press; New York: 1982.
23. Walter SD. The estimation and interpretation of attributable risk in health research. *Biometrics* 1976;32(4):829–49. [PubMed: 1009228]
24. Zee RY, Diehl KA, Ridker PM. Complement factor H Y402H gene polymorphism, C-reactive protein, and risk of incident myocardial infarction, ischaemic stroke, and venous thromboembolism: A nested case-control study. *Atherosclerosis*. Oct 13;2005 Epub ahead of print
25. Christen WG, et al. Prospective cohort study of antioxidant vitamin supplement use and the risk of age-related maculopathy. *Am J Epidemiol* 1999;149(5):476–84. [PubMed: 10067908]
26. Schaumberg DA, et al. Body mass index and the incidence of visually significant age-related maculopathy in men. *Arch Ophthalmol* 2001;119(9):1259–65. [PubMed: 11545630]
27. Rothman, KJ.; Greenland, S. Modern epidemiology. Philadelphia; Lipincott-Raven: 1998. p. 133-34.
28. Seddon JM, et al. A prospective study of cigarette smoking and age-related macular degeneration in women. *JAMA* 1996;276(14):1141–6. [PubMed: 8827966]
29. Lavrovsky Y, et al. Role of redox-regulated transcription factors in inflammation, aging and age-related diseases. *Exp Gerontol* 2000;35(5):521–32. [PubMed: 10978675]
30. Winkler BS, et al. Oxidative damage and age-related macular degeneration. *Mol Vis* 1999;5:32. [PubMed: 10562656]
31. Johnson LV, et al. A potential role for immune complex pathogenesis in drusen formation. *Exp Eye Res* 2000;70(4):441–9. [PubMed: 10865992]
32. Mullins RF, et al. Drusen associated with aging and age-related macular degeneration contain proteins common to extracellular deposits associated with atherosclerosis, elastosis, amyloidosis, and dense deposit disease. *FASEB J* 2000;14(7):835–46. [PubMed: 10783137]
33. McGwin G, et al. The relation between C reactive protein and age related macular degeneration in the Cardiovascular Health Study. *Br J Ophthalmol* 2005;89(9):1166–70. [PubMed: 16113374]
34. Klein R, et al. Systemic markers of inflammation, endothelial dysfunction, and age-related maculopathy. *Am J Ophthalmol* 2005;140(1):35–44. [PubMed: 15939388]

35. Klein R, et al. Early age-related maculopathy in the cardiovascular health study. *Ophthalmology* 2003;110(1):25–33. [PubMed: 12511342]
36. Du Clos TW. Function of C-reactive protein. *Ann Med* 2000;32(4):274–8. [PubMed: 10852144]
37. Kovacs A, et al. A novel common single nucleotide polymorphism in the promoter region of the C-reactive protein gene associated with the plasma concentration of C-reactive protein. *Atherosclerosis* 2005;178(1):193–8. [PubMed: 15585218]

Table 1

Baseline characteristics of study participants.

	Controls (N=401)	AMD Cases (N=111)	P-value
Age (years)	70.1±0.2	65.4±0.7	<0.0001
Smoking Status (%)			0.32
Never	41.2	43.6	
Past	49.2	42.7	
Current	9.5	13.6	
Body Mass Index (kg/m ²)	24.8±0.1	24.9±0.3	0.71
Blood Pressure (mmHg)			
Systolic	134.7±0.7	132.4±1.4	0.13
Diastolic	81.1±0.4	80.4±0.8	0.45
Hyperlipidemia (%)	16.1	14.0	0.60
Hypertension (%)	50.8	35.5	0.004
Diabetes (%)	6.8	5.4	0.61
Aspirin use (%)	44.9	40.5	0.41
Family History of Premature CAD (%)	5.4	8.2	0.28

Mean±SE

Table 2
Complement factor H genotype and allele distribution.

rs1061170 (Y402H)	Controls (N=401)	AMD Cases (N=111)	P-value
Genotype (%)			
<i>TT</i>	44.89	33.33	
<i>TC</i>	41.40	44.14	
<i>CC</i>	13.72	22.52	0.008*
T	0.66	0.55	
C	0.34	0.45	0.005

P-value for the 2 degree of freedom exact test of homogeneity of genotype distribution among cases and controls is 0.03.

* P-value for exact test for trend across genotypes.

Table 3

C-reactive protein genotype and allele distribution.

Genotype (%) or Allele	Controls N=401	AMD Cases N=111	P-value*
rs3093059 (-757T>C)			
TT	86.90	90.11	
TC	12.80	9.89	
CC	0.30	0.00	0.68
T	0.93	0.95	
C	0.07	0.05	0.39
rs2794521 (-717C>T)			
CC	58.55	51.61	
CT	33.91	43.01	
TT	7.54	5.38	0.27
C	0.76	0.73	
T	0.24	0.27	0.50
rs3091244 (-286C>T>A)			
CC	36.97	44.44	
CT	37.88	35.56	
TT	8.79	5.56	
CA	11.82	8.89	
TA	4.24	4.44	
AA	0.30	1.11	0.55
C	0.60	0.65	
T	0.33	0.29	
A	0.07	0.06	0.52
rs1417938 (349T>A)			
TT	49.21	50.94	
TA	40.84	38.68	
AA	9.95	10.38	0.92
T	0.70	0.70	
A	0.30	0.30	0.86
rs1800947(1059G>C)			
GG	86.78	86.49	
GC	12.72	13.51	
CC	0.50	0.00	0.92
G	0.93	0.93	
C	0.07	0.07	0.96
rs1130864(1444C>T)			
CC	46.24	48.39	
CT	43.64	40.86	
TT	10.12	10.75	0.90
C	0.68	0.69	
T	0.32	0.31	0.84
rs1205(1846G>A)			
GG	42.77	42.86	
GA	44.28	43.96	
AA	12.95	13.19	1.00
G	0.65	0.65	
A	0.35	0.35	0.99

All genotype distributions in Hardy-Weinberg equilibrium.

* P-value for exact test of genotype distribution or chi-square test of allele distribution.

Table 4

Single-marker logistic regression analyses for the variants evaluated and risk of AMD.

<i>Additive mode</i>	OR*	95%CI*	P-value
CFH			
rs1061170 (Y402H)	1.46	1.05-2.04	0.02
CRP			
rs3093059(-757T>C)	0.66	0.29-1.54	0.34
rs2794521 (-717C>T)	1.03	0.68-1.56	0.88
rs3091244 (-286C>T>A)	0.86	0.67-1.10	0.23
rs1417938 (349T>A)	0.90	0.62-1.29	0.55
rs1800947 (1059G>C)	1.14	0.59-2.21	0.70
rs1130864 (1444C>T)	0.90	0.61-1.34	0.61
rs1205 (1846G>A)	1.15	0.78-1.68	0.49

* Adjusted for randomized treatment group, age, BMI, history of hypertension, presence or absence of diabetes, smoking status, and CVD status. Results are based on an additive model where genotypes were coded 0, 1, 2, and thus estimates presented are for subjects heterozygous for the risk allele at each locus. The OR for subjects homozygous for the risk allele is equal to the square of the OR for the heterozygotes.