Published in final edited form as:

Tetrahedron Lett. 2007 March 5; 48(10): 1827–1830.

Toward the Total Synthesis of Scleritodermin A: Preparation of the C₁-N₁₅ Fragment

Diver Sellanes, Eduardo Manta, and Gloria Serra*

Cátedra de Química Farmacéutica, Departamento de Química Orgánica, Facultad de Química, UDELAR, Av. General Flores 2124, C. C. 1157, Montevideo, Uruguay.

Abstract

The synthesis of the C₁-N₁₅ fragment of the marine natural product Scleritodermin A has been accomplished through a short and stereocontrolled sequence. Highlights of this route include the synthesis of the novel ACT fragment and the formation of the α -keto amide linkage by the use of a highly activated α , β -ketonitrile.

Keywords

Scleritodermin A; thiazole; α -ketoamide; Wittig

Marine sponges of the order Lithistida are excellent sources of bioactive metabolites.¹ They produce cyclic peptides with non-proteinogenic amino acids and polyketide moieties, such as Cyclotheonamides,² Oriamide,³ and Keramamides.⁴ Recently, another macrocyclic compound, Scleritodermin A (**1**), was isolated by Schmidt and co-workers from the sponge *Scleritoderma nodosum*.⁵ It shows significant cytotoxic activity against a panel of human tumor cells lines (IC₅₀ < 2 μ M). The structure of Scleritodermin A incorporates a novel conjugated thiazole moiety 2-(1-amino-2-*p*-hydroxyphenylethane)-4-(4-carboxy-2,4-dimethyl-2*Z*,4*E*-propadiene)-thiazole (**ACT**), L-proline, L-serine and the unusual amino acids keto-*allo*-isoleucine and *O*-methyl-*N*-sulfo-D-serine.

It has been suggested that the α -keto amide function of the Cyclotheonamides is involved in the deactivation of a protease.^{6,7} The same function, is presented in immunosuppressants such as rapamycin and FK-506.⁸

In spite of such an interesting biological activity, the synthetic challenge defined by the assemblage of the thiazole ring, the polyketide chain and the α -keto amide function in the macrocycle, has not been described in the literature up to date.

Our interest on the synthetic studies of marine natural products⁹ stimulated us to embark upon a total synthesis of this compound and its analogs. We describe, herein, the synthesis of the **ACT** fragment and its coupling to the α , β -ketonitrile derived from L-*allo*-Ile to obtain the key intermediate C₁-N₁₅ fragment.

Our retrosynthetic analysis of Scleritodermin A is shown in Scheme 1. Disconnection at the ester group of the macrolactone and cleavage of the amide bond between keto-*allo*-Ile and L-

*Corresponding author. Tel: + 598 2 9244856; fax: +598 2 924 19 06, email: gserra@fq.edu.uy.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Proline give the C₁-N₁₅ fragment (**2**). This fragment contains a conjugated thiazole and the α -ketoamide moiety, also found in other marine natural products like Oriamide and Keramamides. The disconnection of the amide bond in fragment **2** produces the **ACT** derivative, which could be obtained from thiazole **6**. We propose to synthesize this compound (**6**) by cyclodehydration of a β -hydroxy thioamide derived from L-serine and L-tyrosine.

For synthesizing the **ACT** derivative we began by protecting the amino group of the *O*-benzyl-L-tyrosine with Boc. Then, a coupling reaction with L-serine methyl ester provided product **3** in 96 % yield,¹⁰ Scheme 2. When the corresponding β -hydroxy thioamide **5** was obtained by using the Lawesson's reagent (after protection of the alcohol group), the reaction proceeded in poor yield. In contrast, **5** was obtained in high yield using Wipf's procedure by cyclodehydration of the β -hydroxy amide with DAST,¹¹ giving the oxazoline **4** (87 %) and then thiolysis using H₂S,¹² (92 %). For the synthesis of the thiazole rings our group had reported the use of Deoxo-Fluor and *in situ* oxidation,^{9c} but no reports were found in the literature using DAST and BrCCl₃/DBU. To produce the desired thiazole **6** from the thioamide **5**, we investigated both procedures. The better yield was obtained using DAST/BrCCl₃/DBU (80%). The issue of potential racemization of the tyrosine stereogenic center under the dehydration-dehydrogenation conditions was addressed by chiral HPLC analysis of the thiazole derived from dipeptide **5**.¹³

For synthesizing the **ACT** derivative, the ester group of **6** had to be converted to the corresponding aldehyde. Since the reduction of **6** using DIBALH rendered the aldehyde **7** in poor yield, we used instead LiAlH₄ and then activated MnO₂ (80% yield two steps, Scheme 3).

The Wittig reaction between the aldehyde **7** and 2-(triphenylphosphoranylidene)-propionaldehyde rendered exclusively the *E* alkene (**8**) in 92% yield.

For the last coupling reaction of the **ACT** derivative synthesis (**9**), we investigated different conditions. When the ylide ethyl 2-(triphenylphosphoranylidene)propionate was refluxed in benzene, the *E* isomer was obtained exclusively. Then, the reaction was done in MeOH at 0 ° C,¹⁴ but again only the *E* isomer was obtained. Finally, we used the modification of Still-Gennari for the HWE reaction,¹⁵ and the **9** fragment was obtained in *Z*: *E*, 80: 20 using K₂CO₃ or 97: 3 using KN(TMS)₂ as base. The configuration of the two double bonds was confirmed by NOE correlation between: 1) H₇ of the thiazole ring and the methyl group attached to C₄ and 2) H₃ and the methyl group attached to C₂ (see Scheme 1). Finally, the amine **10** was obtained by cleaving the Boc group of **9**.

Our synthesis of the α -ketoamide **2** (C₁-N₁₅ fragment) utilized the reaction of the nucleophile **10** and a highly electrophilic α , β -ketonitrile derived from L-*allo*-Isoleucine. This procedure was described by Wasserman *et al.*¹⁶ for the synthesis of Cyclotheonamide E₂ and E₃. The synthesis of Cyclotheonamides A and B have been reported by Schreiber,¹⁷ Wipf,¹⁸ Shioiri,¹⁹ Maryanoff⁶ and Ottenheijm.²⁰ All of them obtained the α -ketoamide function by oxidation of α -hydroxy precursors in the final steps of the procedures.

Protection of the amine group of L-*allo*-Isoleucine with Boc and further reaction of its carboxylic acid with the (triphenylphosphoranylidene)acetonitrile ylide using the coupling reagent EDCI, allowed us to obtain the cyano keto phosphorane **11** (Scheme 4). Ozonolysis of **11** generated the α , β -ketonitrile, not isolable. Finally, **10** was added to obtain the desired C₁-N₁₅ fragment (**2**) in 49% yield. The structure of this product was confirmed by the presence of a signal at 194.5 ppm (¹³C-NMR) assignable to the C₁₃ (keto function).²¹

In conclusion, a key fragment of Scleritodermin A was prepared in good overall yield from readily available starting materials. Further progress toward the total synthesis of this natural product will be reported in the due course.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

This work was supported by a grant from NIH/FIRCA (Fogarty International Research Collaboration Award). The authors acknowledge a doctoral fellowship from PEDECIBA (Programa de Desarrollo de Ciencias Básicas) (Diver Sellanes).

References and Notes:

- Bewley CA, Faulkner DJ. *Angew Chem Int Ed* 2162;1998:37.
- a) Nakao Y, Oku N, Matsunaga S, Fusetani N. *J Nat Prod* 1998;61:667. [PubMed: 9599274] b) Fusetani N, Matsunaga S, Matsumoto H, Takebayashi H. *J Am Chem Soc* 1990;112:7053.
- Chill L, Schleyer M, Kashman J. *Tetrahedron* 16147;1997:53.
- a) Itagaki F, Shigemori H, Ishibashi M, Nakamura T, Sasaki T, Kobayashi J. *J Org Chem* 1992;57:5540. b) Kobayashi J, Itagaki F, Shigemori H, Takao T, Shimonishi Y. *Tetrahedron* 1995;51:2525. c) Uemoto H, Yahiro Y, Shigemori H, Tsuda M, Takao T, Shimonishi Y, Kobayashi J. *Tetrahedron* 1998;54:6719.
- Schmidt EW, Ravento-Suarez C, Bifano M, Menendez AT, Fairchild CR, Faulkner DJ. *J Nat Prod* 475;2004:67.
- Maryanoff BE, Greco MN, Zhang H-Ch, Andrade-Gordon P, Kauffman JA, Nicolaou KC, Liu A, Brungs PH. *J Am Chem Soc* 1225;1995:117.
- Lewis SD, Ng AS, Baldwin JJ, Fusetani N, Naylor AM, Sahfer JA. *Thromb Res* 173;1993:70.
- Schreiber SL, Crabtree GR. *Immunol Today* 1992;13:136. [PubMed: 1374612]
- a) Serra G, González D, Manta E. *Heterocycles* 1995;41:2701. b) Serra G, Mahler G, Manta E. *Heterocycles* 1998;48:2035. c) Mahler SG, Serra GL, Antonow D, Manta E. *Tetrahedron Lett* 2001;42:8143. d) Scarone L, Sellanes D, Manta E, Wipf P, Serra G. *Heterocycles* 2004;63:773. e) Mahler G, Serra G, Dematteis S, Saldaña J, Domínguez L, Manta E. *Bioorg Med Chem Lett* 2006;16:1309. [PubMed: 16384701] f) Sellanes D, Scarone L, Mahler G, Manta E, Baz A, Dematteis S, Saldaña J, Domínguez L, Serra G. *Lett Drug Design Disc* 2006;3:625.
- Plant A, Stieber F, Scherkenbeck J, Loesel P, Dyker H. *Org Lett* 3427;2001:3.
- Phillips AJ, Uto Y, Wipf P, Reno MJ, Williams DR. *Org Lett* 1165;2000:2.
- Wipf P, Miller CP, Venkatraman S, Fritch PC. *Tetrahedron Lett* 6395;1995:36.
- Chiralcel OD column; $\lambda = 228$ nm; 10% 2-propanol:hexane; 1.0 mL/min; $t_R = 14, 8$ min.
- Valverde S, Lomas MM, Herradon B, Ochoa SG. *Tetrahedron* 1895;1987:43.
- Still W, Gennari C. *Tetrahedron Lett* 4405;1983:24.
- Wasserman HH, Zhang R. *Tetrahedron Lett* 3743;2002:43.
- Hagihara M, Schreiber SL. *J Am Chem Soc* 6570;1992:114.
- Wipf P, Kim H. *J Org Chem* 5592;1993:58.
- Deng J, Hamada Y, Shioiri T, Matsunaga S, Fusetani N. *Angew Chem Int Ed Engl* 1729;1994:33.
- Bastiaans HMM, Van der Baan J, Ottenheijm HCJ. *J Org Chem* 3880;1997:62.
- Synthesis of the C₁-N₁₅-Fragment (2):** Compound **9** (128 mg, 0.23 mmol) was dissolved in CH₂Cl₂ (2 mL) and TFA (1 mL) was added at 0°C. The reaction mixture was stirred until TLC indicated that all starting material had been consumed. The solvent was evaporated and the residue was dissolved with saturated aqueous sodium bicarbonate. The mixture was extracted with EtOAc, and the combined organic layers were dried with MgSO₄, filtered, and concentrated to afford the nucleophile **10** as a yellow oil (103 mg). **10** was used without further purification. A solution of **11** (126 mg, 0.24 mmol) in CH₂Cl₂ (5 mL) was ozonized at -78°C until the color of the solution remained blue (or yellow-blue). After the solution was purged with N₂ to remove the excess O₃, the nucleophile

10 (103 mg, 0.22 mmol) was introduced at -78°C . The reaction was stirred at -78°C for 2 h and then quenched with HCl 1M and extracted with AcOEt. The combined organic layers were dried with MgSO_4 , filtered and concentrated in vacuo. Purification by flash chromatography (SiO_2 , EtOAc/n-hexane, 1:2) gave the desired product **2** (Z/E mixture) as a yellow oil (76 mg, 49 %). **2**: $R_f = 0.45$ (EtOAc/n-hexane, 1:2); **IR** (film) 2926, 1716, 1699, 1508, 1248, 1016, 748; **$^1\text{H-NMR}$** (CDCl_3 , 400 MHz) δ 0.68 (d, 3 H_{35} , $J = 6.9$ Hz), 1.00 (t, 3 H_{34} , $J = 6.9$ Hz), 1.31 (m, 5 $\text{H}_{33, 27}$), 1.44 (s, 9 H_{31}), 1.96 (m, 1 H_{32}), 2.06 (s, 3 H_{24}), 2.20 (s, 3 H_{21}), 3.29 (m, 2 H_{10}), 4.23 (q, 2 H_{26} , $J = 7.2$ Hz), 5.03 (m, 3 $\text{H}_{28, 15}$), 5.20 (m, 1 H_9), 5.50 (m, 1 H_5), 6.31 (s, 1 H_{22}), 6.50 (s, 1 H_1), 6.88 (d, 2 H_{13} , $J = 8.7$ Hz), 7.03 (m, 3 $\text{H}_{12, 3}$), 7.42 (m, 5 $\text{H}_{17, 18, 19}$), 7.67 (m, 1 H_6); **$^{13}\text{C-NMR}$** (CDCl_3 , 100 MHz) δ 12.0 (C_{34}), 14.1 (C_{35}), 14.5 (C_{27}), 17.8 (C_{21}), 22.0 (C_{24}), 28.6 (C_{31}), 30.0 (C_{33}), 36.7 (C_{32}), 41.4 (C_{10}), 52.5 (C_5), 58.8 (C_9), 61.1 (C_{26}), 70.4 (C_{15}), 79.9 (C_{30}), 115.3 (C_{13}), 117.5 (C_3), 124.2 (C_1), 127.8 (C_{18}), 128.3 (C_{19}), 128.9 (C_{17}), 129.3 (C_{11}), 129.6 (C_{23}), 130.8 (C_{12}), 136.4 (C_{20}), 137.5 (C_{16}), 137.9 (C_{22}), 153.6 (C_2), 152.4 (C_{29}), 158.4 (C_{14}), 167.4 (C_4), 169.2 (C_{25}), 170.2 (C_7), 194.5 (C_8); **HRMS** m/z calcd for $\text{C}_{39}\text{H}_{49}\text{N}_3\text{O}_7\text{S}$ (M^+) 703.3291, found 703.3247.

Scheme 1.
Retrosynthetic analysis of Scleritodermin A (1)

Scheme 2.
Synthesis of thiazole **6**

Scheme 3.
Synthesis of ACT derivative

Scheme 4.
Synthesis of the $\text{C}_1\text{-N}_{15}$ fragment