animal feed to prevent crossover into further species.

While the incubation period for all the slow spongiform viral encephalopathies is long, making an epidemiological study of the extent of bovine spongiform encephalopathy time consuming and expensive, use could be made of an antiserum developed to the so called "prion protein" (PrP 27-30) which has been used successfully to identify these abnormal extracellular filaments which are found in scrapie and Creutzfeldt-Jakob disease.⁵ The appearance of bovine spongiform encephalopathy has given an urgent impetus to the intellectual challenge of finding the infectious agent. Current evidence favours the "virino" (RNA containing subviral particle) theory.⁶

ROY L BISHOP

Glasgow G21 4PG

- 1 Brown P, Rohwer RG, Gajdusek DC. Sodium hydroxide decontamination of Creutzfeldt Jakob disease virus. *N Engl J Med* 1984;310:727.
- 2 Gajdusek DC. Unconventional viruses and the origin and disappearance of kuru. *Science* 1977;197:943-60.
- 3 Kimberlin RH, Walker CA. Pathogenesis of mouse scrapie: patterns of agent replication in different parts of the CNS following intraperitoneal infection. *J R Soc Med* 1982;75: 618-24.
- 4 Clarke MC, Kimberlin RH. Pathogenesis of mouse scrapie: distribution of the agent in the pulp and stroma of infected spleens. *Vet Microbiol* 1984;9:215-25.
- 5 DeArmond SJ, McKinley MP, Barry BA, Braunsfeld MB, McColloch JR, Prusiner SB. Identification of prion amyloid filaments in scrapie infected brain. *Cell* 1985;41:221-35.
- 6 Kimberlin RH. Scrapie agent: prions or virinos? *Nature* 1982; 297:107-8.

that the same practices should be revisited, but this was deemed to be unnecessary.

Whether these differences of view were put before the members of the committee at its meeting on 25 February is not known. Certainly a written response by the region's general practice sub-committee to the 1987 visit sent to the officers of the committee was not made available to the committee members. The chairman's letter of 3 March informing the region of the decision to withdraw training recognition did not give the specific grounds for coming to that decision. Several letters have been sent to the committee requesting details, but the region is still awaiting a written response. That the reported state of the medical records was central to the decision was brought to the attention of the region only orally when the decision had already been made.

As a former joint honorary secretary and chairman of the committee I am dismayed and shocked by its action. That a national body recognised by the secretary of state for the purposes of certification under the vocational training regulations should apparently ignore the region's representations in accepting uncritically the 1987 visitors' report, using controversial evidence to damn the whole region and failing to make the reasons clear in writing, is a matter of grave concern.

S E JOSSE

North Middlesex Hospital,
London N18 1QX

Intravenous volume replacement

Mr G Ramsay rightly gives prominence to the treatment of hypovolaemic shock, a life threatening condition encountered by a wide range of doctors in many specialties. Yet all too often this potentially remediable condition is treated in an ad hoc, empirical way. As Mr Ramsay showed much of this confusion stems from the colloid versus crystalloid controversy.

The crux of this problem seems to be whether clinical reliance is placed on the physiology of the mechanisms protecting against oedema within the pulmonary circulation.¹ Proponents of crystalloid infusion argue that although such infusions will reduce the colloid osmotic pressure the fall in colloid osmotic pressure will be reflected in the pulmonary interstitium owing to the comparative permeability of the pulmonary capillary membrane to protein. This effect coupled with the efficient lymphatic drainage of the pulmonary interstitium prevents an increase in pulmonary hydrostatic pressure and thus attenuates any tendency towards pulmonary oedema.

Other studies relating the comparative importance of left atrial pressure and the colloid osmotic pressure to the development of pulmonary oedema show an increased rate of formation of oedema at pulmonary capillary wedge pressures exceeding 23 mm Hg; this critical pressure point is reduced by a fall in colloid osmotic pressure.² Allowing for individual variation, the critical pressure for the formation of oedema when the colloid osmotic pressure is allowed to fall may well lie within the range of wedge pressures used as end points in resuscitation.³ This uncertainty may be further compounded if the left atrial pressure is inferred from a central venous pressure reading.

The concept of early and late shock is generally regarded as valid, which would suggest that speed is essential in resuscitation to prevent the build up and release of toxic intermediates from the hypoxic microcirculation. Whether peripheral oedema resulting from crystalloid infusion matters is still controversial, but evidence shows that compression of the capillary beds by oedema may retard the restoration of tissue oxygenation.⁴

Mr Ramsay mentioned the adverse properties

of dextrans but not their potential advantages. Although the use of dextran 70 may be associated with a higher incidence of mild or moderate reactions, the incidence of life threatening allergic reactions is no higher for dextran 70 than for gelatin preparations.⁵

Shock is recognised as a hypercoagulable state,⁶ and activation of cascade systems may have an important role in the pathogenesis of associated conditions such as the adult respiratory distress syndrome, disseminated intravascular coagulation, and multiple organ failure. Many studies support the effect of dextrans in attenuating this increase in coagulant activity, and such considerations may arguably make dextran 70 the preferred agent for use in the initial stages of resuscitation.^{3,7}

KENNETH J POWER

Department of Anaesthesia,
Royal United Hospital, Bath

- 1 Zarins CK, Rice CL, Peters RM, Virgilio RW. Lymph and pulmonary response to isobaric reduction in plasma oncotic pressure in baboons. *Circ Res* 1978;43:925-30.
- 2 Guyton AC, Lindsey AW. Effect of left atrial pressure and decreased plasma protein concentration on the development of pulmonary oedema. *Circ Res* 1959;7:649-57.
- 3 Haljamäe H. Rationale for the use of colloids in the treatment of shock and hypovolaemia. *Acta Anaesthesiol Scand* 1985;29: 48-54.
- 4 Shoemaker WC, Schluchter M, Hopkins JA, Appel PL, Schwartz S. Fluid therapy in emergency resuscitation: clinical evaluation of colloid and crystalloid regimens. *Crit Care Med* 1981;9:367-8.
- 5 Ring J, Messmer K. Incidence and severity of anaphylactoid reactions to colloid volume substitutes. *Lancet* 1977;i:466.
- 6 Barrett J, Sheaff C, Smith S, Jonasson O. Correlation of spontaneous micro-aggregate formation with the severity of trauma in man. *J Trauma* 1983;23:389-92.
- 7 Modig J. Effectiveness of dextran 70 versus Ringer's acetate in traumatic shock and adult respiratory distress syndrome. *Crit Care Med* 1986;14:454-7.

Low birth weight and hypertension

In their study of 77 men aged 28 Professor Gerhard Gennser and colleagues found a relation between increased diastolic blood pressure and low birth weight for gestational age (28 May, p 1498). In a prospective study of 3240 children born in Britain and followed up to the age of 36 Wadsworth *et al* found an inverse relation between birth weight and systolic blood pressure in both men and women.¹

In collaboration with Dr J Golding we analysed the blood pressures of a national sample of 9921 10 year old children born in Britain who were followed up from birth. Again blood pressure was inversely related to birth weight. Systolic pressure increased by 0.43 mm Hg for each kilogram increase in weight at 10 years but fell by 0.80 mm Hg for each kilogram increase in birth weight; 95% confidence intervals for these figures were 0.39 to 0.46 and 0.39 to 1.22.

We showed that 75 years ago geographical differences in mortality from stroke and ischaemic heart disease in Britain were closely related to differences in maternal mortality from causes other than puerperal fever.² Higher maternal mortality, mainly from toxæmia and puerperal haemorrhage, occurred among women with poor physique and health. On the basis of the known "tracking" of blood pressure through childhood into adult life and the inverse relation between birth weight and blood pressure we postulated that hypertension was a link between an adverse intrauterine environment and the risk of cardiovascular disease in adult life. Hypertension could be established in the womb through mechanisms such as those suggested by Professor Gennser and others or through accelerated growth of low birthweight babies during early infancy as suggested by Ounsted *et al*.³

We are carrying out further studies of the relation between intrauterine growth and blood

Vocational training in general practice

I appreciate that Dr Dorothy Ward, the immediate past chairman of the Joint Committee on Post-graduate Training for General Practice, had to condense into a few words the reasons why the committee adopted its policy on the withdrawal of training approval in the North East Thames region (28 May, p 1534). The action was no doubt meant to uphold standards and demonstrate that the committee meant business. The implication is not certain as the committee does not have the legal power to refuse certificates of prescribed or equivalent experience after individual applications made under the terms of the vocational training regulations. The approval of hospital posts for general practice training and trainers in general practice is a regional responsibility, and a region's judgment cannot be altered by the committee.

There are several inaccuracies in Dr Ward's letter. In the 1985 visit to North East Thames one training practice out of the 12 visited had records that did not demonstrate the region's minimum criteria of acceptability. At this time the region's policy was that, although the records should be in chronological order, it accepted as a minimum that 75% of the records in a teaching practice had to be in order. In one practice the trainer's records were acceptable but those of his partners were not. After this visit the general practitioner subcommittee revised its policy to comply with that of the committee. After the 1987 visit I personally revisited four of the five training practices that had been inspected. In my opinion the records met the region's and committee's criteria. The fifth trainer had already been highly commended by the visitors. Two of the trainers caused concern to the visitors for reasons that had nothing to do with the state of their records. In fact the course organiser had suggested that they should be inspected as anxieties had been expressed about them which it was thought could be explored with the visitors. In subsequent meetings with and correspondence to the officers of the committee the differences of view between the visitors and the region concerning the medical records were made clear. I suggested