acknowledgment, let alone explanation, of the philosophical debate about this crucial point.

As Dr Higgs says, the principle of double effect deserves a mention in this report. Once it is accepted that there is a distinction in moral terms between the active and passive taking of life one can produce an elaborate and well reasoned justification of our present easy conscience with withdrawing treatment while holding out against active intervention to terminate life. If this is not accepted one has to consider either that active euthanasia, like the withdrawal of treatment, can be morally acceptable or that withdrawal of treatment, like active intervention to terminate life, is morally wrong. Both are attitudes that doctors will find difficult to integrate with current practice.

It is a pity that the report did not take the opportunity to explore the important philosophical questions behind this distinction. The distinction may be a valid one in moral terms, but it would have been a better report had it contained the arguments behind this. As it is one could be left with the suspicion that this is a particular can of worms that the working party was not keen to open up lest it should lead to some uneasy answers.

J P WIGHT

Northern General Hospital,
Sheffield S5 7AU

Organs for transplantation

Professor Geoffrey D Chisholm (21 May, p 1419) discusses the "required request" and "routine inquiry" methods of obtaining organs for transplantation.

Reservations about "beating heart donors" originate from the 1979 memorandum to the 1976 report of the conference of medical royal colleges. The report recommended that absence of response to tests for brain stem death may indicate a hopeless prognosis and is a helpful guide to discontinuing artificial ventilation. The memorandum, however, incorrectly equated that state with death itself on the basis that "by then all functions of the brain have totally and irreversibly ceased." There is ample evidence that this latter deduction was, and remains, untrue.

Relatives who refuse permission for organs to be removed because the patient "has already been through enough" may have greater insight into the situation than the doctors wanting to harvest the organs. The explanation of Professor Chisholm's observation that the results of kidney transplantation from cadavers have come closer to the results with live donations is that the cadavers have come closer to being live donors. We must distinguish clearly between those donors who will be dead after life support is stopped (from whom kidneys and eyes may usefully be taken) and those who are nearly dead (from whom hearts, lungs, livers, and pancreas can be obtained while they remain ventilated, perfused, and reactive). Relatives should be given the opportunity to give or withhold consent for either circumstance.

Some anaesthetists recognise the need to anaesthetise such "cadavers" for surgery and to paralysed them, so the option of anaesthesia for the donor should also be offered specifically to donors or relatives of donors. Prior consent from donors is by a donor card, and this should be rephrased to allow consent to be given for removal of organs either before or after withdrawal of life support. Such consent should be countersigned by the doctor who has explained the procedure.

Professor Chisholm is wrong when he writes that "a patient not on a respirator can never be a donor." It is not necessary to be a "beating heart" donor for one's corneas and kidneys to be of use. People who have deleted all except eyes and kidneys from the organs listed on their donor cards

in the hope that this will ensure that they are not used as "beating heart" donors have no such security at present. This wish should be included explicitly on the donor card.

DAVID J HILL

Addenbrooke's Hospital,
Cambridge CB2 2QQ

Medical education

Dr H J N Andreyev (7 May, p 1326) should have been taught basic scientific principles while he was at secondary school. I was fortunate in being taught under H E Armstrong's "heuristic" method. Instead of being told to "verify the gas laws" I was given apparatus and told how to measure the effect of change of temperature on the pressure of a gas confined to a constant volume and of changes of pressure on its volume when kept at a constant temperature. In such ways I learnt to do reproducible experiments, to distinguish a dependent from an independent variable, and to consider the variability of observation and the dangers of coming to a conclusion on inadequate data.

When I came to the wards and clinics I had to amass my own data in a world in which the data were changing constantly. The rules which seemed to apply to one patient might not apply to another, or even to the same patient at a different time. I had to amass empirical experience. Tenderness at McBurney's point usually suggests appendicitis—but once when it appeared without any of the other signs of appendicitis it turned out to be caused by a hydronephrosis which had ruptured while the patient was at rest in bed. I know of no general scientific principles which will allow you to make these distinctions. If there were any, life would be much easier.

Doctors are technicians; peculiar ones, I agree, and with a remarkable field of work. My contemporaries were fortunate, for a registrar's round rarely had more than five or six students around the bed. Grand rounds were show occasions at which only the clinical clerk had much of a look in. Practical teaching is usually better given by juniors than by consultants, unless the consultant has a special delight in teaching or acting a part.

GEORGE DISCOMBE

Haywards Heath,
West Sussex RH16 1HH

Obstructive uropathy

Dr Kathleen Lyons and others (28 May, p 1517) state that dilatation is the hallmark of obstruction of the urinary tract. While this is widely accepted, it is not true.

Ormand in his report of retroperitoneal fibrosis comments on the almost normal appearance of the renal pelvis in a man with acute anuria,¹ and this has been noted by other authors.² Similar appearances have been noted in obstruction by carcinoma.³ Lalli refers to the common appearance of undilated pelvicaliceal systems in retroperitoneal fibrosis.⁴ Davies suggested that the pelvicaliceal system in acute obstruction was not dilated,⁵ and this has been confirmed by recent ultrasound studies.^{6,7} A range of appearances has been described in obstruction caused by various diseases.⁸ Appreciable dilatation of the pelvicaliceal system and ureters may be present (especially in pregnancy) with no obstruction.⁹

Ultrasound is not good at detecting dilatation of the pelvicaliceal system and is subject to false positive and false negative errors.^{10,11} Enthusiasm for a new technique which is painless, easy, and relatively fast have made ultrasound investigations popular. Failure to follow a sequential series of investigations has been a factor.¹²

In all medical practice it is important to understand the disease. Ultrasound is an imaging technique which shows anatomy and not physiology. It was predictable that it would be less than perfect in investigating many cases of obstruction. The duty of the radiologist is to understand the disease process and choose the most suitable imaging technique or sequence to elucidate the problem. It is too easy for ultrasound studies to become routine examinations which do not contribute to the diagnosis.¹³

PETER DAVIES

Department of Radiology,
City Hospital,
Nottingham NG10 4FG

- 1 Ormand JK. Bilateral ureteric obstruction due to envelopment and compression by an inflammatory retroperitoneal process. *J Urol* 1948;59:1072-9.
- 2 Saxton HM, Kilpatrick FR, Kinder CM, Lessof MN, McHardy-Young S, Wardle DFH. Retroperitoneal fibrosis. *Q J Med* 1969;38:159-81.
- 3 Elliott S, Davies P. The urographic appearances of ureteric obstruction secondary to bladder carcinoma. *Clin Radiol* 1986;37:495-8.
- 4 Lalli AF. Retroperitoneal fibrosis and inapparent obstructive uropathy. *Radiology* 1977;122:339-42.
- 5 Davies P. Urographic appearances in renal failure due to intraluminal ureteric obstruction. *J R Soc Med* 1982;75:768-71.
- 6 Hill MC, Rich JL, Mordait JG, Finder CA. Sonography versus excretion urography in acute flank pain. *American Journal of Roentgenology* 1985;144:1235-8.
- 7 Laing FC, Jeffrey BR, Wing VW. Ultrasound versus excretion urography in acute flank pain. *Radiology* 1985;154:613-6.
- 8 Minford J, Davies P. The urographic appearances in acute and chronic retroperitoneal fibrosis. *Clin Radiol* 1984;38:51-7.
- 9 Davies P. Ultrasound does not diagnose ureteric obstruction. *J R Soc Med* (in press).
- 10 Talner L, Scheible W, Ellenbogen PM, Berk CH, Gosink BB. How accurate is ultrasonography in detecting hydronephrosis in azotemic patients? *J Urol Radiol* 1981;3:1-6.
- 11 Amis ES, Cronan JJ, Pfister RC, Yoder JC. Ultrasonic inaccuracies in diagnosing renal obstruction. *Urology* 1982;19:101-5.
- 12 Anonymous. Diagnosing obstruction in renal failure. *Lancet* 1984;ii:848-9.
- 13 Mindel S. The full potential of ultrasound. *Br Med J* 1988;296:929.

Drug and alcohol misuse in pregnancy

Dr Nabeel D Sulaiman and others (28 May, p 1500) found that 2% of a sample of 901 women in early pregnancy consumed 100 g or more absolute alcohol a week. The psychiatric department at Guy's Hospital recently set up a service within the antenatal clinic for pregnant drug addicts. To gauge some idea of the size of the problem we decided to carry out a random screening. On their initial visit to the antenatal clinic 218 pregnant women were asked to give urine samples. The purpose of the test was explained and verbal permission sought from each patient. Only two women refused; one of them later consented but no drugs or alcohol were found in her urine sample.

Of the remaining 216 women, four had positive results for alcohol, two for opiates, and one for benzodiazepine. Apart from one woman with epilepsy who was taking prescribed phenobarbitone, no other sedative or stimulant drug was detected in the urine samples. All four women who showed alcohol in their urine were over 30. The urine test was sensitive to a blood alcohol concentration of 2.2 mmol/l (10 mg/100 ml). Thus women who consumed alcohol before or during lunch or had drunk heavily the night before were detected. These patients may correspond to Dr Sulaiman and others' category of women who consume 10 or more standard drinks (100 g) a week. If so our findings are nearly identical. Only one of the four women agreed to be interviewed by a psychiatrist, and she claimed abstinence despite her results.

Both women with opiates detected in their urine were under 30. They both accepted treatment and were using less than 20 mg heroin a day. One smoked and the other injected heroin intra-

venously. Both successfully completed gradual withdrawal with prescribed methadone during their second trimester. The one patient who had positive results for benzodiazepines was 19.

At the outset we were concerned about the ethics of screening patients for illicit drugs. Dr J E M Gregg and others (14 May, p 1400) wrote, "Routine screening of urine [in pregnant women] for drugs would be impracticable and possibly unethical." We asked permission from each patient before screening and were pleasantly surprised to find almost 100% compliance. Although 4% of our sample had positive results for drugs or alcohol, this does not mean that they were all addicted or regularly misusing these substances. This figure is lower than expected, given the amount of alcohol¹ and drug abuse^{2,3} in the general population. Some women may stop taking drugs of their own accord when they realise that they are pregnant. Indeed, the two opiate addicts who accepted treatment had already started to reduce their consumption before they were seen. Another explanation may be that opiate addicts seek antenatal care elsewhere knowing that this hospital does not have an established tradition of treating opiate dependence. It would be interesting to hear the experience of other hospitals and the results of similar screening carried out elsewhere.

MERVYN LONDON
MAURICE LIPSEDEGE

Department of Psychiatry,
United Medical and Dental Schools of
Guy's and St Thomas's Hospitals,
Guy's Hospital,
London SE1 9RT

1 Dunbar GC, Morgan DDV. The changing pattern of alcohol consumption in England and Wales 1978-85. *Br Med J* 1987;295:807-10.

2 Richards T. Drug addicts and the GP. *Br Med J* 1988;296:1082.

3 Royal College of Psychiatrists. *Drug scenes: a report on drugs and drug dependence*. London: Gaskell, 1987.

Gastric lavage

Dr V Danel and others (28 May, p 1507) add weight to the argument to stop the routine use of gastric lavage in our accident and emergency departments.

They state that a similar study has not yet been completed. Such a study was, however, reported by Tenenbein *et al.*¹ Ten volunteers ingested 5 g of ampicillin. All subjects acted as controls and on separate occasions had various treatments one hour after ingestion. Serial serum ampicillin concentrations were used to estimate the amount of drug absorbed. The reductions of absorption compared with controls were 32% for gastric lavage, 38% for emesis induced by ipecacuanha ($p < 0.01$), and 57% for activated charcoal (50 g) with magnesium sulphate (30 g) ($p < 0.01$).

The debate between the choice of ipecacuanha or charcoal will continue, but surely the case against gastric lavage is beyond doubt. I suspect that if this procedure had to be performed by doctors instead of nurses the controversy would have been settled already.

RAY MCGLOONE

Wakefield WF1 3RS

1 Tenenbein M, Cohen S, Sitar DS. Efficacy of ipecac-induced emesis, orogastric lavage, and activated charcoal for acute drug overdose. *Am Emerg Med* 1987;16:838-41.

Respiratory illness and home environment of ethnic groups

Dr R J W Melia and colleagues (21 May, p 1438) classified their Asian patients by the language spoken at home. They also coined the graceful word "Urdus" to mean Urdu speakers.

Speaking Urdu does not relate to ethnicity. Urdu, which is a variant of Hindi, is the national language of Pakistan, but there are fewer Urdu speakers there (estimated as eight million) than in India (estimated as 35 million).¹ As speaking Urdu does not relate to origin, culture, race, or lifestyle what is the rationale for its use to classify ethnicity? Gujarati speakers probably originate from the state of Gujarat and Punjabi speakers probably originate from the state of Punjab so using language to classify ethnicity remains a crude but justifiable device. Validation of such classification is, however, desirable.

The term "Asian" is widely used by the public, including people from India and Pakistan, and mass media to describe people whose roots seem to lie in the Indian subcontinent. It is a convenient term to describe a disparate group of people who are no less heterogeneous than Europeans. Scientists should beware of using "Asian" in this inaccurate way. They may mislead the international research community and cause affront, or at least bemusement, to other Asians who are excluded by the narrow use of this word.

Writers of scientific papers who use the term "Asian" should define its meaning. Those who subdivide their Asian groups should discuss the validity of this action. My approach is to italicise "Asian" to remind readers that the term is being used in a restricted way. The invention of a suitable acronym is overdue.

R S BHOPAL

Unit of Community Medicine,
University of Newcastle upon Tyne,
Newcastle upon Tyne NE2 4HH

1 Anonymous. *The world in figures*. London: Hodder and Stoughton, 1987.

AIDS: a bill of rights for the surgical team

Professor H A F Dudley and Mr A Sim (21 May, p 1449) discuss the rights of the surgical team treating a patient who is positive for human immunodeficiency virus (HIV). Now we have had the first medical casualty in the United Kingdom apparently related to occupational exposure to HIV all surgeons must realise that surgical practice in certain areas will have to change if projections of future incidence are borne out.¹

Previous reports have been reassuring, suggesting that the risk is negligible if simple precautions are taken.^{2,3} We note, however, that 1% of HIV contaminated needlestick injuries or mucosal exposures lead to seroconversion.⁴ Such injuries are common, and their frequency and the incidence of glove failure or penetration have been reported.^{5,7} No such data exist for endoscopic urological surgery.

In a recent survey of urological surgery at this hospital (427 consecutive operations, 123 open and 304 endoscopic) we were surprised to find that contamination of mucous membranes or prolonged exposure of apparently normal skin occurred during 99 (33%) endoscopic procedures. Our study confirmed the frequency of needlestick injuries (13 instances) and glove perforations (42) in open surgery. During endoscopic surgery, however, the surgeon's face was often contaminated, particularly the eyes (46 instances). The risk from such exposure is not known, but seroconversion in three laboratory workers with no direct evidence of penetrating skin injury makes it imperative to guard against this risk.⁸ Because endoscopic surgery reduces the opportunity for contamination through penetrating injuries does not mean that the risks should be underestimated. The frequency of contamination may even be greater.

While a high index of suspicion and a carefully

taken sexual, drug, and medical history in young single men will indicate some in high risk groups it is unrealistic to expect to predict all, and more general measures may be necessary. The surgeon's face must be protected, and the previously easy-going acceptance of prolonged exposure of the surgeon's skin to potentially infected body fluids must change. This will require the reorganising of surgical work to reflect the greater time required for safe surgery. Redesign of most endoscopic equipment, particularly urological instruments, should allow a completely closed system of drainage of potentially infective effluent or irrigant. The wider use of a video camera would keep the surgeon's face at a reasonable distance from the source of likely contamination. The cost implications of such changes are likely to be large.

T A MCNICHOLAS

D J JONES

G N A SIBLEY

St Peter's Hospital,
London WC2

1 Tindall G, Lansdowne R. AIDS: public panic. *J R Soc Med* 1987;80:780-2.

2 Geddes AM. Risk of AIDS to health workers. *Br Med J* 1986;292:711-2.

3 Adler MW. *ABC of AIDS*. London: British Medical Journal, 1987.

4 Adler MW. Patient safety and doctors with HIV infection. *Br Med J* 1987;295:1297-8.

5 Lafferty K, Wyatt AP. After safe sex, safe surgery? *Br Med J* 1987;295:392.

6 Hussain SA, Latif ABA, Choudhary AAAA. Risk to surgeons: a survey of accidental injuries during operations. *Br J Surg* 1988;75:314-6.

7 Brough SJ, Hunt TM, Barrie WW. Surgical glove perforations. *Br J Surg* 1988;75:317.

8 Anonymous. Hospital workers have AIDS virus. *Nature* 1987;327:261.

Risk factors for ischaemic brain infarction

Dr E Ernst (21 May, p 1469) suggested that abnormal flow properties of blood could predispose to strokes. Impaired white cell rheology has assumed increasing importance as a determinant of cerebral blood flow, especially in acute ischaemia, and may contribute to ischaemic brain infarction.¹

During an acute infection the number of white cells increases and fibrinogen concentrations are raised, this being the expression of the acute phase proteins.² Blood flow in the nutritive capillaries may be impaired as a result. The potential flow resistance can be studied by filtering blood or blood cell suspensions through polycarbonate filters with pores of similar diameter to that of the nutritive capillaries.³ The preliminary results of our studies using a modified version of the Lennie filtration method⁴ on white cell rheology show that acute bacterial infections cause a decrease in whole blood filterability by a rise in plasma viscosity and monocyte rigidity.

Impaired white cell rheology (particularly in the monocyte subpopulation), whatever the original cause, may play an important part in the development of ischaemic stroke.

GIOVANNI CIUFFETTI

MICHELE MERCURI

MARIA TERESA RIZZO

RITA LOMBARDINI

LAURA SANTAMBROGIO

Universita Degli Studi,
06100 Perugia,
Italy

1 Lowe GDO. Blood rheology in arterial disease. *Clin Sci* 1986;314:137-46.

2 Harkness J. The viscosity of human blood plasma: its measurement in health and disease. *Biorheology* 1971;8:171-93.

3 Lowe GDO. Blood rheology in general medicine and surgery. *Clin Haematol* 1987;1:827-61.

4 Lennie SE, Lowe GDO, Barbenel JC, Forbes CD, Foulds WS. Filterability of white blood cell subpopulations, separated by an improved method. *Clinical Hemorheology* 1987;7:811-6.