criteria be waived because of the urgency of the situation is shared by the gay community. We should not, however, abandon all scientific caution for a treatment that may be worse than the disease. The best way to do this is to encourage enrolment into the trial and to hope that results will become apparent quickly so that the trial need not run its planned course and that publication and dissemination of the results will be rapid so that the usual long delays can be avoided.

ALEX MILLS

Lesbian and Gay Medical Association,
London WC1N 3XX

- 1 McKerrrow G. 2000 guinea pigs for £3 million AZT trial. *Capital Gay* 1988 Oct 21:1,9.
- 2 Mills AF. We need this trial of AZT. *Capital Gay* 1988 Oct 28:2.
- 3 Watney S. They cannot justify putting 2000 people through 3 years of stress. *Capital Gay* 1988 Nov 4:2.
- 4 Davies P. Dr Gelmon meets her critics. *Capital Gay* 1988: Nov 4:11.
- 5 Mann C. Don't accept this "doctor knows best" attitude. *Capital Gay* Nov 11:2.
- 6 Mills AF. One way to fight HIV is to support important research. *Capital Gay* 1988 Nov 18:2.

Echocardiography

Dr J B Chambers and colleagues (29 October, p 1071) provided an excellent review of the state of cardiac ultrasound imaging and Doppler investigation, but their last line—sadly, the one that many administrators may study ("the bottom line")—does no service to their subject. The statement about colour flow Doppler that "there would have to be a high clinical load to justify its purchase" is extremely misleading.

Clearly there must always be adequate clinical usage of any method of investigation to justify its purchase. But ultrasound is safe, painless, and speedy. The capital equipment cost is low compared with other imaging techniques (radiographic, isotopic). Colour Doppler imaging is a truly magical advance compared with earlier techniques. In the review I counted nine separate statements in which the authors emphasised the advantages of colour Doppler over pulsed or continuous wave. For example, "Colour flow Doppler mapping is speedy and simple to use"; "colour flow Doppler may quickly show abnormalities of flow even without echocardiographic abnormalities"; and "Colour flow Doppler makes contrast echocardiography for detecting shunts virtually redundant."

The working group of the British Cardiac Society recommended provision for 250 echocardiograms per year per 100 000 population.¹ This might mean an average district general hospital performing 500 studies. Even in 1985, however, the year that provided these figures, several district hospitals were performing twice this number of investigations. These presumably were the hospitals in which there was a consultant with the training to run an echocardiography laboratory as the technique in its present form (two dimensional imaging) is only a decade old. An increase in demand may be expected. There would seem to be a reasonable clinical load to justify modern equipment.

But there is a more pressing need for modern and not outdated equipment: regional centres have never been so hard pressed. It is a waste of regional time, facilities, and transport costs to send a patient to the regional centre without a precise diagnosis. One should argue that never has there been a greater need for diagnostic precision in the periphery than now. Incorrect assessment or diagnosis at a district general hospital may enhance the pride of the staff at the regional centre ("we got it right again—anyone should have known it was an arteriovenous fistula of the sinus of Betelgeuse") but it represents a waste of money and is unnecessary.

Drs Chambers and Monaghan are two of this country's most skilful echocardiographers and

usually echocardiography's most persuasive advocates. In the text of their review they emphasise the value of colour Doppler imaging. May we please put it on the bottom line?

ANDREW McLEOD

Poole General Hospital,
Poole,
Dorset BH15 2JB

- 1 Working group of the British Cardiac Society. Report. Cardiology in the district hospital. *Br Heart J* 1987;58:537-46.

Effectiveness of geriatric rehabilitative care

Dr David C Kennie and others (29 October, p 1083) published the first prospective study of collaborative care in the aftercare of patients with hip fractures. It may be, however, that more caution is needed in interpreting the data presented.

Age and cognitive state are major determinants of morbidity and mortality after hip fracture.¹⁻⁴ It is therefore crucially important to isolate the effects of these two factors from the influence of the treatment regimen. The treatment group were younger (median age was 5 years less) and had better cognitive state (scores on the Pfeiffer short portable mental state questionnaire were higher) than the controls.

The probability derived from a χ^2 statistic may not be a valid means of dismissing selection bias in the treatment group. Furthermore, the subdivision of mental test results into four groups may not be justified by Pfeiffer's original work on the test.⁵ He combined the moderate and severely impaired groups and found a 92% agreement with a clinical diagnosis of "organic brain syndrome." He also emphasised the need to take educational attainment into account in scoring performance.

Although the effects of age are tabulated in control and treatment groups separately, this procedure may unmask the effects of the higher cognitive state in the treatment group. Tabulation of cognitive state in control and treatment groups may be equally misleading.

Ideally, the design should incorporate age and cognitive state matching or stratification at entry to the study to avoid these difficulties in interpretation.

Finally, the authors give no information on mortality in the treatment and control groups. This information is vital in interpreting median length of stay in hospital, and any difference in mortality between the groups is likely to influence outcome.

NIGEL SMITH

Department of Health Care of the Elderly,
University Hospital,
Nottingham NG7 2UH

- 1 Evans GJ, Prudham D, Wandless I. A prospective study of fracture of the proximal femur: factors predisposing to survival. *Age Ageing* 1979;8:246-50.
- 2 Cobey JC, Cobey JH, Conant L, et al. Indicators of recovery from fractures of the hip. *Clin Orthop* 1976;117:258-62.
- 3 Kreutzfeld TJ, Haim M, Bach E. Hip fracture among the elderly in a mixed urban and rural population. *Age Ageing* 1984;13:111-9.
- 4 Katz S, Heiple KG, Downs TD, Ford AB, Scott CP. Long-term course of patients with fracture of the hip. *Surg Gynecol Obstet* 1967;124:1219-30.
- 5 Pfeiffer E. A short portable mental state questionnaire for the assessment of organic brain deficit in elderly patients. *J Am Geriatr Soc* 1975;23:433-41.

AUTHORS' REPLY.—Age and cognition clearly affected the study outcomes but were correlated variables. Taken together, they did not fully explain the differences between the two groups, nor did mortality between trial entry and orthopaedic inpatient discharge (five in treatment and four in control group).

Stratification would have required large

numbers of patients to ensure that all the known influences on the trial outcomes were equally distributed between the two trial groups. The complexities of this design were inconsistent with the practical and clinical realities of establishing this trial in a district hospital.

Any interpretation of mental state is difficult in the absence of data before the fracture. Cognition at trial entry was influenced by postoperative confusion, and cognition later might have been influenced by group treatment effects. Dr Smith's concerns about the mental test scores, however, are not entirely justified. Pfeiffer's short portable mental state questionnaire is part of the larger OARS (Older American research and service center instrument) questionnaire for assessing the elderly. It has been used for more than a decade in over 150 evaluative studies.¹ The influence of educational attainment on test scores has also been shown to be minimal.² Finally, Hodkinson's mental test score³ was applied concurrently with the short portable mental state questionnaire and, although they were not published, gave closely correlated results.

DAVID C KENNIE

JOHN REID

IAN R RICHARDSON

CHRISTINE KELT

Department of Geriatric Medicine,
Royal Infirmary,
Stirling FK8 2AU

- 1 George L, Fillenbaum GG. OARS methodology, a decade of experience in geriatric assessment. *J Am Geriatr Soc* 1985;33:607-15.
- 2 Erkinjuntti T, Sulkava R, Wikström J, Autio L. Short portable mental status questionnaire as a screening test for dementia and delirium among the elderly. *J Am Geriatr Soc* 1987;35:412-6.
- 3 Hodkinson HM. Evaluation of a mental test score for assessment of mental impairment in the elderly. *Age Ageing* 1972;1:233-8.

Slowing the speedy

I cannot accept Dr Robert West's assertion that the most effective way to reduce fatalities is vigorous enforcement, or even reduction, of the current 70 miles/h (113 km/h) speed limit (29 October, p 1069).

No doubt one reason that this speed limit is ignored by as many as 40% of vehicles¹ is that it is patently safe to travel at higher speeds on motorways in good conditions. If it was safe for cars to travel 70 miles/h in 1965 then in today's cars, with their improved braking, road holding, and power, it should be safe to travel at a higher speed. The Association of Chief Police Officers' main priority should be to prevent driving liable to cause accidents, and its recent statement on speed limits seems to be addressed to this priority. The association apparently recognises that it is relatively safe to travel at speeds of up to 84 miles/h (135 km/h).

While it may be true that high speed motorway driving often leads to dangerous high speed urban driving, rigorous enforcement of motorway speed limits seems an odd and inefficient approach to the problem. Motorways are considerably safer to travel on than other roads. In 1986, 4.5% of deaths from road accidents in Britain occurred on motorways, despite their carrying 14.2% of traffic.³ Higher speed limits do not appear to affect this phenomenon. For example, in West Germany, with no mandatory maximum speed limit, 8.5% of road traffic accident deaths occur on motorways carrying 26.8% of traffic.

A more pragmatic solution is needed. In the short term it would be appropriate to raise the motorway speed limit to 84 miles/h (135 km/h) while vigorously enforcing urban speed limits near motorway exits. To reduce motorway accidents due to impaired driving conditions it would be helpful to ensure that motorists obey the speed limits on overhead motorway signals. A more long