TABLE 11—Out of hours consultations per doctor and rates (per 1000 patients per year) for 1988

	Plymouth		London	
	No of consultations	Rate	No of consultations	Rate
Emergency surgery	122	10.17	62	4.43
Visits	364	30.33	236	16.86
Night visits	59	4.92	34	2.43
Total	511	42.58	298	21.29

index is based on eight doctors sharing the on call duties in London (that is, it excludes trainees) and five doctors in Plymouth (except the emergency surgery, which is taken by six doctors).

The figures are self explanatory, but some comment is worth while. In 1988 out of hours consultations represented 6% of all our consultations, compared with 4% for the London practice. Perhaps the most relevant statistic is the out of hours risk to the doctor on-call; our rate is roughly double that in London.

It should be emphasised that we do not work in a readily identifiable deprived inner city area such as Tower Hamlets. I estimate a Jarman deprivation index of 10–20 for our worst area, compared with the London authors' index of 50–60 in some areas. I suspect that our figures are representative of many parts of the country that are equally alleged to be not "deprived." Hence it is clear that the out of hours demand is what society wants and expects, a point that Mr Clarke seems to have ignored.

With pecuniary disincentives for appointing additional partners, an ever increasing daytime workload, and a ridiculous out of hours burden how are we to find the time for all the extra commitments of the new contract?

E L BROWN

St Neot's Surgery,
Plymouth PL2 3BJ

- Livingstone AE, Jewell JA, Robson J. Twenty four hour care in inner cities: two years' out of hours workload in east London general practice. *Br Med J* 1989;299:368-70. (5 August.)
- Jarman B. Identification of underprivileged areas. *Br Med J* 1983;286:1705-9.

Treating patients with asthma who are dependent on systemic steroids

SIR,—Drs R J Shiner and D M Geddes have stated that the important anxiety concerning the long term use of systemic steroids is that patients commonly develop osteoporosis.¹ They have outlined the methods that can be used to reduce the dose of steroids to avoid this complication.

An alternative would be to prevent and even correct corticosteroid induced osteoporosis by using oestrogens. Prolonged corticosteroid treatment is associated with decreased plasma oestradiol and androstenedione concentrations.² It also reduces testosterone and androstenedione concentrations in men.³ The loss of skin collagen and skin thickness after corticosteroid treatment is also a feature of the oestrogen deficiency state that follows the menopause.⁴ All of these changes can be corrected by oestrogen treatment in postmenopausal women, and we have reported an increase in bone density after percutaneous oestrogen or testosterone treatment in postmenopausal women receiving systemic corticosteroids.⁵

We have 11 postmenopausal patients receiving steroids long term; they had a mean increase in spinal bone density of 8.4% and an increase in the proximal femur bone density of 7.4% after one year of having percutaneous implants of 50 mg oestradiol and 100 mg testosterone. A further six patients receiving corticosteroids long term and

also hormone implants for a mean of 8.4 years have been shown to have normal bone density in cross sectional studies. The improvement in bone density after one year clearly continues with duration of hormone treatment.

We believe that hormone replacement treatment with or without testosterone is an essential adjunct to corticosteroid treatment in both premenopausal and postmenopausal women. It is logical, but unproved, that testosterone may be necessary to reduce many of the skeletal and psychosexual complications of long term corticosteroid treatment in men. Research on such neglected male patients is urgently required, but in the mean time it is difficult to understand why oestrogens are not used to prevent steroid induced osteoporosis in women.

JOHN STUDD
MICHAEL SAVVAS

King's College Hospital,
London SE5 9RS

MARGARET JOHNSON

Royal Free Hospital,
London

- Shiner RJ, Geddes DM. Treating patients with asthma who are dependent on systemic steroids. *Br Med J* 1989;299:216-7. (22 July.)
- Crilly RG, Marshall DH, Nordin BEC. Metabolic effects of corticosteroid therapy in post-menopausal women. *J Steroid Biochem* 1979;11:429-33.
- MacAdams MR, White RH, Chipps BE. Reduction of serum testosterone levels during chronic glucocorticoid therapy. *Ann Intern Med* 1986;104:648-51.
- Brincat M, Wong Ten Yuen A, Studd JWW, Montgomery J, Magos AL, Savvas M. Response of skin thickness and metacarpal index to estradiol therapy in post-menopausal women. *Obstet Gynecol* 1987;70:538.
- Studd JWW, Savvas M, Johnson M. Corticosteroid induced osteoporosis and hormone implants. *Lancet* 1989;i:563.

Acute right ventricular infarction

SIR,—Although we agree that haemodynamic monitoring with pulmonary artery catheters is an important part of the management of right ventricular infarction, as stated by Dr J L Caplin,¹ we do not agree with Drs N Naqvi and S Mushahwar that the management of patients with cardiogenic shock after acute myocardial infarction is impossible without a pulmonary artery catheter and radial artery line.² We also do not believe that pulmonary artery pressure monitoring can be discussed in the context of hypotension complicating acute myocardial infarction without mentioning the benefits of thrombolysis and possible complications of invasive pressure monitoring in this group of patients.

The second international study of infarct survival indicated that thrombolysis was most beneficial in patients at greatest risk of death, such as those with anterior infarcts and those with systolic blood pressures below 100 mm Hg.³ This type of myocardial infarct is often complicated by left ventricular failure, which should be obvious with the aid of clinical examination and a chest radiograph showing pulmonary congestion. In this group of patients the first priority should undoubtedly be the early administration of a thrombolytic agent. The second international study showed a 53% reduction in vascular mortality at five weeks when streptokinase and aspirin were administered within four hours of the onset of symptoms compared with a 32% reduction when they were administered between five and 12 hours.

We believe that the benefits of invasive pressure monitoring in anterior myocardial infarction with hypotension are outweighed by the proved benefits of early thrombolysis, the potential delay in administration of streptokinase while pulmonary artery pressure monitoring is started, and the risk of haemorrhage from the puncture site after thrombolysis. For the same reason we believe

that a radial artery line is also contraindicated. Hypotension and heart failure in this group of patients should be treated by diuretics and inotropic support.

With inferior myocardial infarction complicated by hypotension, especially with a raised jugular venous pressure and clear lung fields, we agree that pulmonary artery pressure monitoring contributes important information towards the management of the patient, which should generally consist of optimising filling pressures by the administration of intravenous fluids, inotropic support, and occasionally diuretics. Even under these conditions, however, the benefits of thrombolysis at an early stage outweigh those gained by pressure monitoring.

Our current clinical practice is to ensure the early administration of a thrombolytic agent in all patients without a contraindication who have a clinical myocardial infarction. In patients with anterior myocardial infarction complicated by hypotension and radiographic evidence of pulmonary oedema we try to avoid invasive haemodynamic monitoring within the first 24 hours. In myocardial infarction complicated by hypotension with clear lung fields, especially clinical right ventricular infarction, we insert pulmonary artery catheters at an early stage. After thrombolysis these catheters are inserted through the femoral or basilic veins by an experienced operator of at least registrar grade. Catheter insertion at these sites allows local pressure haemostasis in the event of haemorrhage.

We believe that Drs Caplin and Naqvi and Mushahwar have overstated the role of pulmonary artery pressure monitoring in hypotension complicating acute myocardial infarction without mentioning the benefits of thrombolysis and discussing the risks and benefits of inserting a pulmonary artery catheter after thrombolysis.

M R THOMAS
D SMYTH

Department of Cardiology,
King's College Hospital,
London SE5 9RS

- Caplin JL. Acute right ventricular infarction. *Br Med J* 1989;299:69-70. (8 July.)
- Naqvi N, Mushahwar S. Acute myocardial infarction. *Br Med J* 1989;299:459-60. (12 August.)
- ISIS-2 Collaborative Group. Randomised trial of intravenous streptokinase, oral aspirin, both, or neither among 17 187 cases of suspected acute myocardial infarction: ISIS-2. *Lancet* 1988;ii:349-60.

General practitioners' referrals

SIR,—Ms Angela Coulter and colleagues have studied cross boundary referrals to outpatient clinics and found that general practitioners referred to a large number of locations.¹ They speculated about the reasons for referrals to other health authorities and pointed to the need to establish what existing patterns of referral are before arranging hospital service contracts as a result of the white paper.

Cross boundary flow is an existing financial problem for some Scottish health boards, with large payments being made from one health board to another on the basis of admissions under Scottish health authorities revenue equalisation (SHARE).² Before evaluating the costs and benefits of providing new services within a health authority compared with contracting services from other health authorities we need to know about the reasons for admission to hospitals elsewhere.

We have studied all patients resident in Ayrshire and Arran Health Board who were admitted to hospitals in Greater Glasgow Health Board for orthopaedic and ear, nose, and throat surgery during 1986. Patients were identified by reference to the computer tapes of discharge data (SMR1) held by Greater Glasgow Health Board. The