Current Practice

GROWING POINTS

Autoimmune Disease—II*

G. L. ASHERSON,† D.M., M.R.C.P.

The first part of this article described the way in which autoimmune responses—namely, autoantibodies, delayed hypersensitivity, and sometimes tissue damage—could be produced in animals. These autoimmune responses occur when (a) the animal is exposed to antigens similar to but not identical with self; (b) the animal is exposed to its own body constituents, together with an adjuvant such as Freund's complete adjuvant; and (c) spontaneously in New Zealand black mice. The tissue damage seen is regarded as autoimmune in nature because

(1) It is provoked by the injection of the appropriate organ, preferably in Freund's complete adjuvant, which increases and modifies immune responses. Well-characterized antigens such as thyroglobulin are able to cause autoimmune disease.

(2) It can be prevented by treatment with cortisone and folic-acid antagonists under conditions which limit immune responses, and by antigen under conditions which limit delayed hypersensitivity.

(3) The histology of the affected organ, showing infiltration with lymphocytes and histiocytes, is characteristic of delayed hypersensitivity.

(4) In the case of allergic encephalomyelitis the lesions can be transferred to normal recipients by spleen cells.

(5) Circulating autoantibodies can be demonstrated.

It is possible to cause autoantibody formation in animals, however, without producing tissue damage. For instance, rats injected with carbon tetrachloride produce autoantibodies against their own livers which are not thought to damage the liver.

It is easy to recognize disease produced by the injection of antigen as autoimmune. In man, allergic encephalomyelitis sometimes follows the injection of rabbit spinal cord used in the treatment of suspected rabies. The post-cardiotomy and post-myocardial-infarction syndromes sometimes follow damage to the heart, and the autoimmune response is probably responsible for the clinical picture. Ligation of the vas deferens by making available or altering testicular antigen causes the formation of autoantibody to sperm. These are probably responsible for infertility in some patients.

In most cases, however, the agent causing an autoimmune response is unknown, and this makes it difficult to determine whether the autoimmune response causes tissue damage or whether tissue damage gives rise to an autoimmune response. There are three possibilities:

(A) The initiating agent causes an immune response which is responsible for the main features of the disease. An example is human encephalomyelitis produced by rabbit spinal cord. If the tissue damage caused by the autoimmune response leads to further damaging autoimmune responses the disease will be self-perpetuating.

(B) The initiating agent causes tissue damage leading to an autoimmune response which is responsible for the main features of the disease. The post-cardiotomy syndrome is a possible example. Here again the autoimmune response may be self-perpetuating.

(C) The initiating agent causes the tissue damage and is responsible directly for the clinical and pathological changes of the disease.

The autoimmune response is a result of the tissue damage, or is a direct effect of the initiating agent but is unimportant in pathogenesis. The formation of autoantibodies to pancreas seen in obstruction of the pancreatic duct is a possible example.

Because it is often uncertain whether an autoimmune process is the result or cause of disease, the term putative autoimmune disease is used in this article. There are certain common features of putative autoimmune diseases. These are:

(i) Raised serum gammaglobulins, which offer indirect evidence of immunological activity because antibodies are gammaglobulins.

(ii) Direct evidence of an autoimmune response, as shown by auto-antibody and delayed hypersensitivity.

(iii) Tissue lesions characterized by infiltration with lymphocytes, histiocytes, and plasma cells.

(iv) Genetic incidence of the disease, and clinical and serological association with other autoimmune diseases, in either the patient or his family.

(v) Association with enlargement of the thymus or thymic tumour or both.

(vi) Ameliorating effect of corticosteroids and cytotoxic drugs.

(vii) Production of similar disease of an autoimmune aetiology in animals.

The occurrence of these features in a disease provides evidence that it resembles other putative autoimmune diseases.

Diseases in which Autoimmune Reactions may be Provoked by Micro-organisms

Rheumatic Fever.—This disease is always associated with group A beta-haemolytic streptococcal infection. These organisms have an antigen similar to human heart. Autoantibodies which react with streptococcal extracts, and can be absorbed by the human heart, may be found by gel-precipitation techniques in acute rheumatic fever. They also occur in acute glomerulonephritis and less frequently in uncomplicated streptococcal pharyngitis. The autoantibodies probably arise because of exposure to streptococcal antigen resembling human heart. The cause of tissue damage is less clear. Patients with rheumatic heart disease who undergo cardiac surgery have an increased incidence of post-cardiotomy syndrome, compared with patients with congenital heart disease. This syndrome is characterized by precordial pain, fever, and pericardial effusion. There are autoantibodies against heart, and the response to steroids is good. The raised incidence in subjects with past rheumatic fever may be related to the fact that they have already produced an autoimmune response to heart tissue.

Acute Glomerulonephritis.—This is associated with certain strains of group A beta-haemolytic streptococci. The type-12 nephritogenic strain has antigens similar to those of renal glomeruli. Acute glomerulonephritis is one of the few diseases (systemic lupus erythematosus is another) in which the level of serum complement is low. A low serum-complement level provides some evidence of complement-fixing antigen-antibody reactions *in vivo*. Serum complement is often raised when there is tissue damage, and not all immune reactions fix complement. Hence the finding of normal serum comple-

* The first part of this article appeared in last week's issue (p. 417).

† Reader in Immunology, Department of Bacteriology, the London Hospital Medical College, London E.1.

ment is not evidence against the presence of immunological disease.

Renal Disease.—Renal damage is a common cause of death in systemic lupus erythematosus, and it is often suggested that various forms of glomerulonephritis are immunological in nature. There are several different methods of producing immunological renal disease:

(a) Masugi nephritis, caused by the injection of antikidney serum. For a long time this was the chief evidence that renal disease might be immunological.

(b) Immune response to foreign antigens adhering to the glomeruli. Certain bacterial antigens adhere to the glomeruli, and immune responses against them may cause renal damage.

(c) Injection of kidney tissue in Freund's complete adjuvant. This may be an example of an autoimmune response against the kidney.

(d) Circulating antigen-antibody complexes. When antibody is mixed with an excess of antigen soluble antigen-antibody complexes are formed. These complexes are often able to bind (fix) complement and to cause renal damage. Their presence in the circulation is usually associated with a low serum-complement level, and they may be responsible for the renal damage in systemic lupus erythematosus.

The demonstration of gammaglobulin bound to the tissues by immunofluorescence is often taken as evidence of local combination of antibody and antigen. The presumption is strengthened by the demonstration of complement in the lesion by immunofluorescence. Nevertheless, bound gammaglobulin is seen in the glomeruli in certain experimental diseases of non-immunological nature, and this evidence requires careful assessment in renal disease.

Ulcerative Colitis.—In about 10% of these patients autoantibodies to colon can be demonstrated by immunofluorescence. Autoantibodies can also be demonstrated by haemagglutination, using red cells coated with polysaccharide derived from colonic tissue. These autoantibodies react with an antigen which occurs in sterile human and rat colon, and in certain strains of *Esch. coli*. The suggestion is that these autoantibodies arise because the human body is stimulated by bacterial antigens similar to but not identical with its own tissue components. But it is not certain that autoantibody production is an important part of the disease, and it is possible that immune reactions to bacterial residues in the tissues has a more central role. In some cases food allergy, particularly to milk, is important; and in controlled trials withdrawal of milk has been shown to diminish the frequency of relapse.

Coeliac Disease.—Antibodies to gluten show a high frequency and titre in patients with coeliac disease. There is no evidence that this is an autoimmune disease, but it may be an immune reaction to foreign antigens. Autoantibodies to gut occur in pernicious anaemia and ulcerative colitis, and in both of these the incidence of carcinoma of the affected organ is raised. In coeliac disease the incidence of lymphosarcoma of the small intestine is raised (but this is still low).

Glandular Fever.—Autoantibodies against formed elements of the blood are sometimes found, and they may be responsible for haemolytic anaemia and thrombocytopenia. The virus has not been isolated, and we do not know whether it shares antigens with human tissue.

Primary Atypical Pneumonia.—This is due to *Mycoplasma pneumoniae*. Cold agglutinins (which are autoantibodies directed against the I antigen of human erythrocytes) are often found, and they may present clinically as Raynaud's phenomenon or haemolytic anaemia and jaundice on exposure to cold. A raised titre of antibodies to *M. pneumoniae* is sometimes found in the Stevens-Johnson syndrome.

Hashimoto's Thyroiditis Group of Diseases

This group of diseases includes Hashimoto's thyroiditis, pernicious anaemia, and non-tuberculous Addison's disease of

the adrenal. Certain forms of hypoparathyroidism and diabetes may ultimately be shown to belong to this group. The diseases have several common features:

(I) Autoantibodies occur which react with the affected organ, and can be demonstrated by complement fixation and immunofluorescence. Autoantibody against cell secretions—thyroglobulin and intrinsic factor—occur in Hashimoto's thyroiditis and pernicious anaemia.

(II) Patients with one disease commonly have autoantibodies characteristic of the other diseases of the group. These autoantibodies are called organ-specific, because they react with antigen from one organ only. In contrast, the autoantibodies in systemic lupus erythematosus react with widely distributed nuclear and cytoplasmic antigens. These non-organ-specific antibodies are rare in the Hashimoto's thyroiditis group of diseases.

(III) Patients sometimes have more than one disease in the group.

(IV) The patients give a family history of their own disease and of other diseases in the group.

These diseases are clinically important, because they may present insidiously and simulate psychological disorders.

Nearly all patients with Hashimoto's thyroiditis have autoantibodies against thyroid by at least one test, and the diagnosis is usually wrong if the tanned-cell-haemagglutination test for thyroglobulin, the complement-fixation reaction for antibody against cytoplasmic antigen, the immunofluorescent test for antibody against thyroglobulin, the second colloid antigen (CA2), and the cytoplasmic antigen are all negative. Low titres of antibody against thyroglobulin by the tanned-cell test are not uncommon, being shown in about 15% of women over the age of 50. Like many other autoantibodies which occur in a few apparently normal people, the incidence is much higher in women than in men. The complement-fixation reaction is rarely positive in the absence of disease. The immunofluorescent and complement-fixation tests for cytoplasmic antigen run in parallel. The immunofluorescent test is a little more sensitive (90% of patients with Hashimoto's thyroiditis give positive results), but there is a higher incidence of borderline positive results (0–15%) in apparently healthy subjects.

In the differential diagnosis of carcinoma of the thyroid and Hashimoto's thyroiditis the absence of autoantibodies is strong evidence against Hashimoto's thyroiditis. The converse is not true, and the presence of autoantibodies does not exclude carcinoma.

Tissue damage in these diseases is probably due to autoimmune processes. This is suggested by the histology of the affected organ, which is infiltrated with lymphocytes and histiocytes, and the finding of autoantibodies. Cytotoxic antibodies to thyroid cells in tissue culture are almost invariably present in active Hashimoto's thyroiditis. The process which initiates autoantibody formation is unknown. The classical view was that the thyroid antigens were secluded and therefore unable to induce tolerance, but recently thyroglobulin has been demonstrated in the circulation. Perhaps its levels are too low or too variable for the maintenance of tolerance.

Graves's Disease.—Patients with Graves's disease have a thyroid-stimulating factor which behaves differently from thyrotrophic hormone, and is known as the long-acting thyroid stimulator or L.A.T.S. Recently it has been shown that L.A.T.S. is a gammaglobulin, which presumably acts by combining with the thyroid gland and stimulating the action of thyrotrophic hormone. The implication is that the hypersecretion in Graves's disease is due to an autoantibody. The thymus is usually enlarged in Graves's disease, and contains germinal centres, which are morphological evidence of antibody production. They are not found in the normal thymus. Similar changes are seen in myasthenia gravis and systemic lupus erythematosus, and in a few cases of Hashimoto's thyroiditis that have been examined. The incidence of autoantibodies to thyroid, as detected by conventional tests, is raised in Graves's disease. There is, however, a particularly high incidence of autoantibodies to thyroglobulin in exophthalmic goitre.

Pernicious Anaemia.—Autoantibodies against intrinsic factor are a distinctive feature of this disease, and they do not occur in patients with uncomplicated gastritis. They are detected by their effect on the electrophoretic mobility of radioactive-B₁₂ intrinsic-factor complexes. Autoantibodies against the secretory cells of the stomach can be detected by complement-fixation and immunofluorescence in 50 to 90% of patients with pernicious anaemia. As in ulcerative colitis, the incidence of carcinoma of the affected organ is raised. The incidence of antibodies to stomach is increased in chronic gastritis (diagnosed by biopsy) and idiopathic iron-deficiency anaemia. Antibodies against stomach are not reported in infantile pernicious anaemia. These patients have normal acid secretion, and infantile pernicious anaemia is a distinct disease from the adult form.

Myasthenia Gravis and the Role of the Thymus

Autoantibodies against muscle have been demonstrated in myasthenia gravis. The techniques used for their detection include complement fixation, immunofluorescence, and tanned-cell haemagglutination. Autoantibodies are also produced against an antigen in the thymus, and occasionally against nuclei. Myasthenia gravis is sometimes associated with other putative autoimmune diseases, such as Graves's disease. Lymphorrhages (accumulations of lymphocytes) similar to those seen in thyrotoxicosis, Hashimoto's thyroiditis, and Addison's disease are found in the muscles. In this disease the relationship between the autoantibodies and the characteristic neuromuscular block is unknown.

The thymus is often enlarged, and contains germinal centres which provide morphological evidence of antibody production. In some cases there is a thymic tumour, and these patients characteristically have high titres of autoantibodies and respond poorly to thymectomy. Thymic tumours also associated with aplastic anaemia, dermatomyositis, polymyositis, granulomatous myocarditis, hypergammaglobulinaemic purpura, and hypogammaglobulinaemia. The question arises whether autoimmunity is important in any of these diseases.

The role of the thymus in disease is still obscure. In early post-natal life cells leave the thymus to populate the lymphoid system, and the thymus produces a hormone which favours the development of immunologically competent cells. The thymus is required in adult life for the rapid restoration of immunological function after damage to the lymphoid system by x-rays and other agents. The presence of germinal centres in the thymus in myasthenia gravis presumably reflects antibody production which may be directed against autoantigens.

Diabetes Mellitus

The possibility of autoimmune mechanisms in diabetes mellitus is under discussion. The most suggestive evidence is the association of Hashimoto's thyroiditis, Addison's disease, and sometimes hypoparathyroidism with diabetes (Schmidt's syndrome). Antibodies against insulin are commonly found in treated diabetics. They may be directed against human insulin, and arise as a result of stimulation with foreign insulin. They are sometimes responsible for insulin resistance.

Systemic Lupus Erythematosus and Related Diseases

Systemic lupus erythematosus is characterized clinically by multisystem involvement. The increase in gammaglobulin suggests increased antibody production. The wide range of autoantibodies characteristically react with widely distributed antigen, and these are described as non-organ-specific; the incidence of organ-specific autoantibodies of the type seen in the Hashimoto thyroiditis group of diseases is little altered.

The antibodies against nuclei (antinuclear factor) are usually detected by immunofluorescence. This test is nearly always positive in systemic lupus erythematosus, and diagnosis of the disease is nearly always wrong when the antinuclear-factor test is negative. The antinuclear-factor test, unlike the L.E.-cell test, is usually positive even when the disease is clinically inactive. The L.E.-cell test also detects antibodies against nuclei. It is positive much less frequently than is the antinuclear factor test, but there are very occasional patients with positive L.E. and negative antinuclear-factor tests.

Antibodies against cytoplasm are also found, and are detected by complement fixation—the autoimmune complement-fixation reaction. There is sometimes a biologically false-positive test for syphilis. Rheumatoid factor (which behaves like an antibody to altered gammaglobulin) occurs in about a third of patients. Autoantibodies against formed blood elements and coagulation factors may cause haemolytic anaemia, thrombocytopenia and bleeding disorders, and some cases of apparent idiopathic thrombocytopenia and thrombotic thrombocytopenic purpura develop features of systemic lupus erythematosus.

Patients with systemic lupus erythematosus have an unusual tendency to give drug and blood-transfusion reactions, and relapses may be provoked by drug reactions. In children the anticonvulsant drugs may precipitate an illness resembling systemic lupus erythematosus; the prognosis of this drug-induced disease is apparently better than that of the "idiopathic" disease. There is evidence that when patients with systemic lupus erythematosus are injected with antigens which do not cause autoantibody formation in normal subjects they produce a range of autoantibodies.

The histology of the thymus in systemic lupus erythematosus is of considerable interest. Lymphoid infiltrates and germinal centres are found, but few observations are available, because corticosteroids and terminal illness cause involution of the thymus.

The incidence of hypergammaglobulinaemia, antinuclear factor, and diseases of the systemic lupus erythematosus group is slightly raised in the relatives of patients with systemic lupus erythematosus, and in a few families the incidence of related disease is very high. The familial association with hypogammaglobulinaemia is an intriguing finding.

The aetiology of systemic lupus erythematosus is unknown, although New Zealand black mice provide a good natural model of the disease. Because of the many different autoantibodies found, this disease can be described as a disorder or breakdown of immune tolerance, but it is not clear whether this is due to exposure to altered antigens or to a defect in the antibody-producing system.

Rheumatoid Arthritis.—Rheumatoid factor, which behaves like an antibody to altered gammaglobulin (and some of the allotypes of gammaglobulin), is found in most patients with chronic rheumatoid arthritis showing joint erosions on radiological examination and nodules. It is often absent in the first year of the disease, and it is found in about 5% of normal subjects. Rheumatoid factor can be produced in animals by immunizing with antigen-antibody complexes; by infection, which probably acts by producing antibody-antigen complexes *in vivo*; and by immunization with altered gammaglobulin. Rheumatoid factor is often found in subacute bacterial endocarditis, where it has been suggested as a diagnostic aid, and in a range of diseases—including infectious hepatitis, cirrhosis of the liver, sarcoid, scleroderma, and pulmonary fibrosis. In most diseases, other than rheumatoid arthritis, the latex-fixation test (based on human gammaglobulin) is more often positive than the Rose-Waaler reaction (based on rabbit gammaglobulin).

Antinuclear factor and positive L.E. tests are found in some cases of rheumatoid arthritis. Systemic lupus erythematosus rarely develops in these patients, but they tend to have somewhat more severe disease with a raised incidence of arthritis.

Though the arteritis involves digital and visceral vessels, and the peripheral neuropathy is attributed to it, renal damage is rare. The affected joints in rheumatoid arthritis show infiltration with lymphocytes, histiocytes, and sometimes plasma-cells, and the formation of lymphoid follicles. Pannus is formed and there is erosion of cartilage. These changes can all be produced by implanting particulate (insoluble) antigens into the joints of animals; streptolysin S of the streptococcus has a similar effect. These findings raise the question of whether the human disease is due to the presence of antigenic material in the joints, and there is current interest in the possible role of mycoplasma and other micro-organisms.

Sjögren's Disease presents as failure of secretion by the lacrimal and salivary glands, and it is often associated with rheumatoid arthritis. It is of considerable immunological interest, because organ-specific antibodies against intralobular salivary-gland ducts can be demonstrated by immunofluorescence, while non-organ-specific antibodies, which react with nuclear and cytoplasmic cell components, often occur. In some patients the clinical picture merges with that of systemic lupus erythematosus.

Dermatomyositis.—Some types of dermatomyositis may belong to the systemic lupus erythematosus group of diseases. Antinuclear factor is sometimes found, and the disease may develop features of systemic lupus erythematosus. In adults there is a frequent association with carcinoma.

Miscellaneous Conditions

Liver Disease.—Patients with infectious hepatitis often have antibodies against altered gammaglobulin (positive latex test) and complement-fixing antibodies against cytoplasm. This is an example of autoantibody formation caused by infection.

Cirrhosis of the liver and infiltration of the liver with lymphocytes is sometimes associated with minor manifestations of systemic lupus erythematosus, such as rashes and haemolytic anaemia. Many of the patients have positive L.E.-cell tests, and antinuclear factor, antibody against smooth muscle, and complement-fixing antibodies against cytoplasm are found. This disease, which is called lupoid hepatitis, responds to some extent to cortisone.

Primary biliary cirrhosis presents as jaundice simulating that of extrahepatic biliary obstruction. The vast majority of these patients have positive autoimmune complement-fixation reactions, often to very high titres. The antibody reacts with a mitochondrial antigen: this is useful in differential diagnosis, because antibodies against mitochondria are rare in extrahepatic biliary obstruction.

Multiple Sclerosis.—The ease with which encephalomyelitis can be produced in animals by immunization with brain in Freund's complete adjuvant has led to the suggestion that multiple sclerosis may be an autoimmune disease. The gammaglobulin in the cerebrospinal fluid, which is responsible for the positive colloidal-gold reaction, is raised. Recently auto-antibodies have been demonstrated which damage myelinated cells in tissue culture; but similar antibodies occur in motor-neurone disease, and these particular autoantibodies may be the result rather than the cause of tissue damage.

Carcinomatous Neuropathy.—Carcinoma, especially oat-cell carcinoma of the bronchus, is often associated with neuropathy which is not due to local metastases. Antibodies against the nerve bodies of posterior-root ganglia and other cells are found in sensory carcinomatous neuropathy. Tumours sometimes secrete material which is not produced by the normal cells from which they are derived. The question arises whether the tumours associated with certain forms of carcinomatous neuropathy have an antigen similar to brain tissue and so cause autoimmune disease.

Peripheral Neuritis.—Autoantibodies against nervous tissues are found in about a quarter of patients with the Guillain-Barré syndrome of peripheral neuritis associated with high protein content but low cell count in the cerebrospinal fluid.

Pemphigus is characterized by Nikolsky's sign—that is, the stratum corneum can be easily separated from the underlying epidermis—which is the result of damage to the prickle-cell layer. Autoantibodies against these latter cells can be demonstrated by immunofluorescence.

Aphthous Ulcer.—Autoantibodies against buccal mucosa have been demonstrated by complement fixation and immunofluorescence.

Conclusion

Immunologists at the turn of the century were aware of the damaging effect of immune reactions, and they considered the possibility of autoimmune disease. Interest in autoimmune disease revived with the identification of autoimmune haemolytic anaemia and the detection of autoantibodies adhering to red cells with the help of antihuman gammaglobulin serum (Coombs test). Burnet's concept of immune tolerance indicated a special mechanism whereby animals do not produce antibodies against their own tissue components, and this implied that altered physiology might lead to autoimmune reactions and disease. Two sets of observations were historically important. Firstly, the discovery of the L.E. test and the demonstration of antibodies against nuclei in systemic lupus erythematosus; this provided an example of a multisystem disease associated with multiple autoantibody production. Secondly, the production of various autoimmune diseases, including encephalomyelitis and thyroiditis, by immunization with the corresponding organ in Freund's complete adjuvant, and the demonstration of autoantibodies in Hashimoto's disease.

Early interest was focused on the following factors:

(1) Were the antibodies found in putative human autoimmune disease true antibodies? In all cases investigated autoantibodies have been shown to have the physicochemical properties of antibodies, and they are found by methods used to detect antibodies. Rheumatoid factor was originally considered to be an exception because it is a macroglobulin; but it is now known that many antibodies, especially those produced shortly after immunization or in response to particulate antigen, are macroglobulins.

(2) Were the antibodies indeed autoantibodies and not isoantibodies reacting with antigens produced in some but not all humans, and not in the patient from whom the antibody was taken? With the exception of some of the rheumatoid factors, the initial view that an antibody was an autoantibody has proved to be correct.

(3) Is autoantibody or autoreactive delayed hypersensitivity responsible for tissue damage? Methods are not yet available to answer this question. It seems likely that in some situations both circulating antibody and cellular factors are important.

(4) Are autoimmune responses the cause or merely the result of tissue damage?

Despite the inability to answer the last two questions, and the unsatisfactory situation in which putative autoimmune disease is virtually synonymous with the discovery of an autoimmune system, current research interest is now centring on the mechanism of production of autoimmune responses.

Three of the lines of approach are (a) the role of antigens similar but not identical with self; (b) the role of micro-organisms, including mycoplasma and viruses; and (c) the role of the hydrolytic enzymes in the intracellular organelles called lysosomes, which are able to degrade normal tissue components.

REFERENCES

- Asherson, G. L., *Abstr. Wld Med.*, 1965.
Glynn, L. E., and Holbrow, E. J., *Autoimmune Disease*, 1965. Oxford.
Humphrey, J. H., and White, R. G., *Immunology for Students of Medicine*, 2nd edition, 1964. Oxford.