Correspondence

Letters to the Editor should not exceed 500 words.

Aspirin and Children

SIR,—There seems to be increasing concern, amounting almost to panic, about the use of aspirin in therapeutics. If one remembers that any useful drug is potentially toxic, it is still true to say that aspirin is one of the safest and most useful drugs we have today.

In the paper by Dr. J. O. Craig and others (26 March, p. 757) salicylates are said to be of real value in children only in rheumatic fever and rheumatoid arthritis. Alternatives are then suggested to their use as antipyretics and as sedatives. Leaving aside the question whether the alternatives are as good (and I do not think they are), no mention is made of salicylates as analgesics—a very common use in children. To have a drug which is analgesic, antipyretic, and sedative in one is an advantage too great to throw away.

The way to prevent salicylate poisoning in children is by educating the parents about the correct dosage, making sure that it is clearly and prominently marked on the aspirin containers, and educating medical students in the dosage and signs of over-dosage of salicylates in children. There will always remain a few cases of salicylate poisoning in children, some of which may prove fatal, but when these are compared with the immense benefit which salicylates give to many thousands of children in Great Britain alone

annually their retention and widespread use is fully justified.—I am, etc.,

University College of Swansea, T. C. DANN.
Swansea.

SIR,—Drs. J. O. Craig, Isobel C. Ferguson, and J. Syme suggest that aspirin should be withheld from young children (26 March, p. 757). They say that tepid sponging is more effective in reducing temperature, and that chloral is a safe and effective sedative. No doubt this is true, but they do not suggest what analgesic should be given in its place. They do not recommend paracetamol.

It appears that they think young children do not feel pain in circumstances which do not warrant the use of a powerful (and potentially dangerous) analgesic, or, if they do, that they should be allowed to suffer. Perhaps they would care to nurse a child with otitis media with tepid sponging and chloral while awaiting its response to an antibiotic. It's a pity they don't have to try.

Dose for dose aspirin must be one of the safest and most effective drugs we use, and I shall continue to allow my young patients the relief from pain and discomfort which it can afford.—I am, etc.,

Enfield, Essex.

J. F. WOOD.

Popliteal Aneurysms

SIR,—The leading article on popliteal aneurysms (12 March, p. 625) is important, excellent, and timely, for these aneurysms are common.

Atherosclerotic popliteal aneurysms in the 45 to 60 age group seem to follow two patterns: single short or saccular in the normotensive, bilateral long or fusiform in the hypertensive.

Complete thrombosis of the aneurysm causes gangrene if the thrombus extends over a long enough area to block the several branches of the popliteal artery; thrombosis in the single short saccular aneurysm may not produce more than severe claudication, while thrombosis in the long fusiform aneurysm will almost certainly cause gangrene.

The importance of pressure behind the knee in precipitating thrombosis is well illustrated by two amputees; one where thrombosis of his popliteal aneurysm developed after he had had to sit for four hours snowbound in his car, and in the other where minor embolic episodes caused transient paraesthesiae and coldness of his left foot while driving his car in the week preceding complete thrombosis of the left of his

bilateral popliteal aneurysms. The long length seat cushioned under the knee would be best avoided by the atherosclerotic and at worst should be occupied for a short time only.

Operative treatment for a popliteal aneurysm is mandatory, except where other aneurysms are evident in addition to bilateral popliteal aneurysms, for in such a patient with aneurysms of several different arteries the atherosclerosis is so generalized that death is likely to precede gangrene of a foot. Excision of the saccular aneurysm and end-to-end graft can safely be done, but it is equally satisfactorily treated by proximal and distal ligation and a bypass graft, which is the procedure carried out for the fusiform aneurysm where an extensive dissection of this aneurysm would almost certainly cause venous thrombosis. The main reason, probably, that the synthetic graft has replaced the freeze-dried homograft is its ready availability. The freeze-dried homograft can be very satisfactory, for such a graft used to replace a saccular aneurysm in a man aged 57 remained patent for nine years and when it thrombosed the claudication allowed him function adequate for his retirement. It may

be that thrombosis develops earlier in a small-bore synthetic graft than in a homograft or an autogenous vein graft, but the dissection necessary to obtain a length of vein adequate for a bypass graft is extensive and unreasonable if it is carried out on a leg afflicted with a fusiform aneurysm. A graft which thromboses after 10 days preserves the limb but does not, of course, prevent claudication. When a popliteal aneurysm is revealed, and particularly if there are repeated small embolic episodes, proximal ligation of the aneurysm alone would be better than no operation; it would prevent major gangrene. Sympathectomy can be worth while when only minor nutritional changes persist in the toes some weeks after thrombosis of the popliteal aneurysm.—I am, etc.,

Leeds 2.

H. S. SHUCKSMITH.

SIR,—I was interested to see that both Mr. H. H. G. Eastcott and Professor A. J. H. Rains (26 March, p. 799) agree that ligation of the superficial femoral artery, as practised by John Hunter, still has a place in the treatment of popliteal aneurysm. I certainly agree, but would add that it is good policy to perform a lumbar sympathectomy immediately prior to ligating the vessel. This not only improves the collateral circulation but, as Sir James Learmonth first taught me, not infrequently causes the aneurysm to thrombose before the ligation is applied.

As regards resection and replacement, most students (postgraduate as well as undergraduate) appear to regard this as a fairly new procedure. It is worth pointing out, therefore, that Lexer of Freiburg resected an axillary aneurysm and replaced it by a vein graft (albeit unsuccessful) in 1907, and by 1913 successful operations of this kind had been reported by Lexer and also by that remarkable British surgeon, Hogarth Pringle. A specimen showing a patent vein graft which was obtained from one of Hogarth Pringle's patients at necropsy 3½ years after resection and replacement of a popliteal aneurysm exists in the museum of the Royal College of Surgeons of Edinburgh, and a photograph of it is reproduced in my book.¹—I am, etc.,

MICHAEL WOODRUFF.

University of Edinburgh
Medical School,
Edinburgh 8.

REFERENCES

- Woodruff, M. F. A., *The Transplantation of Tissues and Organs*, 1960. Springfield, Ill.

SIR,—I was most interested in your leading article on popliteal aneurysms (12 March, p. 625).

My experience of aneurysms of the popliteal artery due to atherosclerosis over the last 35 years—and I have seen a great many—differs in many important aspects from that expressed in your leading article.

There are two distinct types of popliteal aneurysm due to atheroma. The first type, which accounts for at least 90% of these aneurysms, is saccular and arises from a