attacks, occurring between 8.30 and 9 a.m., were noted to follow runs of coupled ectopic beats, and recurrent ventricular fibrillation was shown by E.C.G. (Fig. 1). Each attack either terminated spontaneously or was terminated by external cardiac massage, but the basic dysrhythmia of coupled ventricular ectopic beats persisted.

FIG. 2.—Electrocardiogram taken before and after intravenous administration of propranolol.

Propranolol, 5 mg., was then injected slowly intravenously. By the end of the injection the E.C.G. showed no ectopic beats (Fig. 2). Whereas before the injection the patient had been suffering syncopal attacks every three to four minutes, after the propranolol one hour passed before she had a further attack. The attacks then recurred and became more frequent and prolonged. She was therefore given a further intravenous injection of 5 mg. propranolol at 12.30 p.m. and suffered no further syncopal attacks for three hours. Coupled ventricular ectopic beats then recurred, and she had one short attack of ventricular fibrillation. Propranolol, 20 mg. by mouth, was started and given six-hourly. She had no further attacks, and her atrial fibrillation was converted to sinus rhythm with one direct-current shock one week later. She was discharged home on anti-coagulants alone.

COMMENT

Until the advent of direct-current shock (Lown *et al.*, 1962 ; Oram *et al.*, 1963), quinidine was the only satisfactory way of reverting atrial fibrillation to sinus rhythm. A test dose to detect hypersensitivity and anticoagulation to prevent embolism are probably necessary precautions. Digitalis has been recommended to avoid paradoxical tachycardia (Sokolow, 1951), though some authorities think that dysrhythmias occur more commonly when the two drugs are combined, and therefore advise that digitalis should not be used (Gold, 1950).

In spite of careful observation of the patient and his E.C.G. occasional ventricular dysrhythmias can occur in an almost totally unpredictable manner. These would normally be expected to occur during, or within six to eight hours of the end of, the course of quinidine (Goodman and Gilman, 1955). However, toxic effects may appear much later (Slonim, 1961), and in our patient the maximal toxic effect was not reached until 14 hours after the last dose. This should act as a warning to prolong observation of patients having quinidine therapy until well after the last dose has been given.

Sudden collapse during quinidine therapy is now thought to be due usually to ventricular fibrillation. In a review of the literature Thomson (1956) found an overall incidence of sudden death of 3.3%, most of these deaths being presumably due to

a dysrhythmia. Rokseth and Storstein (1963) report a series of 274 patients treated without fatality, though 12 of their patients suffered syncopal attacks. Proved ventricular fibrillations as a toxic effect of quinidine therapy has been reported on numerous occasions (Davis and Sprague, 1928 ; Sokolow, 1951 ; Selzer and Wray, 1964 ; Davies, Leak, and Oram, 1965). Rainier-Pope *et al.* (1962) make the point that in their two patients, as in ours, the attacks were heralded by frequent ventricular ectopic beats.

Quinidine toxicity has been treated in a number of different ways with varying degrees of success, the aim of treatment being to tide the patient over until the quinidine has been metabolized. Molar sodium lactate (Bellet, 1963), sympathomimetics (Rokseth and Storstein, 1963 ; Linenthal and Zoll, 1963), calcium sodium versenate (Luchi *et al.*, 1963), and external massage and direct-current shock (Rainier-Pope *et al.*, 1962) are among the remedies that have been used either clinically or experimentally. In our patient the E.C.G. abnormality was closely similar to that occurring in digitalis intoxication, and may well have been due to a synergistic effect of the two drugs, in spite of the small dose of digoxin used. It was felt, therefore, that the dysrhythmia might respond to treatment with one of the adrenergic beta-receptor-blocking drugs, which have been used with good effect in digitalis intoxication (Stock and Dale, 1963 ; Vaughan Williams and Sekiya, 1963). In our patient the results of this therapy appeared to indicate that propranolol has a specific effect in the suppression of this dysrhythmia. It provides further evidence of the usefulness of beta-adrenergic blockade in the treatment of drug-induced ventricular dysrhythmias.

I wish to thank Dr. W. van't Hoff for allowing me to publish details of this case, and Dr. J. P. P. Stock for his help in the preparation of the paper.

A. SEATON, B.A., M.B., M.R.C.P.,
Medical Registrar, North Staffordshire Royal Infirmary,
Stoke-on-Trent ; present address, Royal Southern
Hospital, Liverpool.

REFERENCES

- Bellet, S. (1963). *Clinical Disorders of the Heart Beat*, 2nd ed. Kimpton, London.
- Davies, P., Leak, D., and Oram, S. (1965). *Brit. med. J.*, 2, 517.
- Davis, D., and Sprague, H. B. (1928). *Amer. Heart J.*, 4, 559.
- Gold, H. (1950). *Quinidine in Disorders of the Heart*. Hoeber, New York.
- Goodman, L. S., and Gilman, A. (1955). *Pharmacological Basis of Therapeutics*, 2nd ed. Macmillan, New York.
- Linenthal, A. J., and Zoll, P. M. (1963). *Circulation*, 27, 5.
- Lown, B., Amarasingham, R., and Neuman, J. (1962). *J. Amer. med. Ass.*, 182, 548.
- Luchi, R. J., Helwig, J., and Conn, H. L. (1963). *Amer. Heart J.*, 65, 340.
- Oram, S., Davies, J. P. H., Weinbren, I., Taggart, P., and Kitchen, L. D. (1963). *Lancet*, 2, 159.
- Rainier-Pope, C. R., Schrire, V., Beck, W., and Barnard, C. N. (1962). *Amer. Heart J.*, 63, 582.
- Rokseth, R., and Storstein, O. (1963). *Arch. intern. Med.*, 111, 184.
- Selzer, A., and Wray, H. W. (1964). *Circulation*, 30, 17.
- Slonim, R. (1961). *J. Amer. med. Ass.*, 177, 791.
- Sokolow, M. (1951). *Amer. Heart J.*, 42, 771.
- Stock, J. P. P., and Dale, N. (1963). *Brit. med. J.*, 2, 1230.
- Thomson, G. W. (1956). *Circulation*, 14, 757.
- Vaughan Williams, E. M., and Sekiya, A. (1963). *Lancet*, 1, 420.

Unilateral Renal Vein Thrombosis with Nephrotic Syndrome

Brit. med. J., 1966, 1, 1523-1524

Bilateral renal vein thrombosis, though rare, has been described as a cause of the nephrotic syndrome. However, only a few case reports of unilateral renal vein thrombosis associated with the nephrotic syndrome have been published (Ashner, 1927 ; Pollak *et al.*, 1956 ; Morris *et al.*, 1963).

This report describes such a case. The diagnosis of renal vein thrombosis was suspected from the renal histopathology and radioisotope renography and was confirmed by inferior vena cavography.

CASE REPORT

A 19-year-old youth who worked in a grain shop was referred to us for puffiness of the face and swelling of the feet of five months' duration. The onset was insidious and was not preceded by pain in the loins. There was no history of respiratory symptoms suggestive of pulmonary embolism.

On examination he had a moderate degree of anasarca, mainly in the face and the legs. There was no evidence of collateral circulation suggestive of inferior vena caval obstruction. Blood-pressure was 110/76 mm. Hg and the fundus oculi were normal. There were no other abnormal features.

Examination of the urine revealed a specific gravity of 1018, a quantitative proteinuria of 6 g./day, and, on microscopic examination, a moderate number of granular and hyaline casts and occasional pus cells. Urine cultures, repeated six times, were sterile. Routine blood counts were normal, and E.S.R. (Westergren) was 42 mm. at the end of one hour. Total proteins were 4.12 g./100 ml., and paper electrophoresis of the serum revealed that albumin was 1.082, α_1 -globulin 0.249, α_2 -globulin 0.874, β -globulin 0.998, and γ -globulin 0.915. Blood cholesterol was 286 mg./100 ml., blood urea nitrogen 11 mg./100 ml., and blood creatinine 1.2 mg./100 ml. The glucose-tolerance curve was within normal limits. The Congo-red test was negative. L.E. cells were not detected. Serum sodium was 132 mEq/l. and potassium 4.2 mEq/l. Bleeding, clotting, and prothrombin times were normal. Fluoroscopy of the chest did not reveal any abnormality. Intravenous pyelogram showed good excretion of the dye on both sides.

Renal biopsy, initially carried out on the left side, showed minimal thickening of the basement membrane of the glomerular tufts, severe degenerative and necrotic changes in the tubules, and oedema and infiltration with chronic inflammatory cells of the interstitium. Renal biopsy on the right side showed normal findings. The presence of interstitial oedema and marked tubular damage which was out of proportion to the glomerular involvement led us to explore the probability of renal vein thrombosis as the cause of the nephrotic syndrome.

Renogram studies were done after an injection of 25 μ c of 131 I-labelled Hippuran (sodium iodohippurate). Two scintillation probes were placed over the kidneys, which were localized by means of the I.V.P. films, and simultaneous tracings of the pattern of excretion of Hippuran were obtained on the recorder. The renogram showed a lower peak of the vascular phase on the left side, thus supporting our belief that there was left renal vein thrombosis. The renogram curve of the right kidney appeared normal.

Inferior vena cavogram carried out through the retrograde saphenous vein confirmed the diagnosis of left renal vein thrombosis by revealing a block at the origin of the left renal vein.

The patient was given prednisolone in a dosage of 40 mg. daily for 10 days: the oedema subsided, but the proteinuria, though it lessened, persisted. Anticoagulants were not administered.

COMMENT

Renal vein thrombosis with or without thrombosis of the inferior vena cava is being increasingly recognized as a cause of the nephrotic syndrome. The mechanism which causes this association is not well understood. In experimental animals, partial obstruction of the inferior vena cava proximal to the renal veins or of one or both the renal veins themselves leads to persistent proteinuria (Rowntree *et al.*, 1913; Morris *et al.*, 1963). Similar proteinuria has been observed after thrombosis of the inferior vena cava or of one or both the renal veins in man (Shattock, 1913; Ashner, 1927; Derow *et al.*, 1939; Vilk, 1940; Hasson *et al.*, 1957; Bayley *et al.*, 1965). The considerable loss of albumin leads to hypoalbuminaemia and to a reduced colloid osmotic pressure (Squire, 1953). These factors, and other consequences of protein loss, lead to oedema (Pollak *et al.*, 1956).

Renal vein thrombosis may be found in association with hypercoagulable states, renal amyloidosis (Vilk, 1940; Harrison *et al.*, 1956; Hasson *et al.*, 1957), diabetic glomerulosclerosis (Hasson *et al.*, 1957), or malignancy (Harrison *et al.*, 1956). In many cases, however, the cause is not detected. Except during infancy, primary renal vein thrombosis is rare, probably because of rapidity of blood flow. In a review of

228 cases of renal vein thrombosis by Abeshouse (1945), 40% of the patients were less than 2 months old. In infancy it occurs especially as a complication of acute gastro-enteritis (Harrison *et al.*, 1956).

Clinically, the diagnosis of renal vein thrombosis in a case of the nephrotic syndrome may be suspected when (1) lumbar pains precede the onset (Abeshouse, 1945; Pollak *et al.*, 1956); (2) collateral circulation is present due to associated inferior vena caval obstruction (Pollak *et al.*, 1956; Morris *et al.*, 1963; Bayley *et al.*, 1965); or (3) there are pulmonary embolic manifestations (Blainey *et al.*, 1954; Pollak *et al.*, 1956).

In the case of the nephrotic syndrome described here there were no clinical findings suggestive of renal vein thrombosis. As we have already described the diagnosis was suspected from a disproportion between the gross renal tubular and interstitial changes compared with the relatively minor changes in the glomeruli, an observation already made by Kark in 1955. The radio-Hippuran renogram strengthened our doubts by demonstrating a functional impairment in the renal vascular flow. The diagnosis was established by inferior vena cavography.

The management of this type of case usually includes anticoagulants to prevent extension of the thrombus and pulmonary embolism (Pollak *et al.*, 1956; Morris *et al.*, 1963; Bayley *et al.*, 1965). In the present case anticoagulants were not thought necessary, as there was no evidence of those diseases already noted with which renal vein thrombosis may be associated.

We would like to thank Dr. S. V. Joglekar, Dean of the K.E.M. Hospital and Seth G.S. Medical College, Bombay, for permission to publish an account of this case, Dr. (Mrs.) S. M. Deshmukh for carrying out the inferior vena cavography, and Professor N. M. Purandare for suggestions.

R. C. KHOKHANI,* M.D.,

Senior Medical Registrar and Research Associate.

P. RAGHAVAN,* M.D.,

Professor/Director of Medicine.

S. M. SANT,* M.D.,

Assistant Professor of Pathology.

R. D. GANATRA,† M.B., B.S., F.C.P.S., M.SC.,

Scientific Officer.

S. M. SHARMA,† D.M.R.T.,

Scientific Officer.

M. J. SHAH,* M.R.C.P., D.T.M.&H.,

Honorary Physician and Honorary Professor of Medicine.

* K.E.M. Hospital and Seth G.S. Medical College, Bombay 12.

† Radiation Medicine Centre, Medical Division, Atomic Energy Establishment, Trombay, Bombay.

REFERENCES

- Abeshouse, B. S. (1945). *Urol. cutan. Rev.*, **49**, 661.
 Ashner, P. W. (1927). *J. Urol. (Baltimore)*, **17**, 309.
 Bayley, T. J., Hardwicke, J., Heath, D., and Whitfield, A. G. W. (1965). *Postgrad. med. J.*, **41**, 88.
 Blainey, J. D., Hardwicke, J., and Whitfield, A. G. W. (1954). *Lancet*, **2**, 1208.
 Derow, H. A., Schlesinger, M. J., and Savitz, H. A. (1939). *Arch. intern. Med.*, **63**, 626.
 Harrison, C. V., Milne, M. D., and Steiner, R. E. (1956). *Quart. J. Med.*, **25**, 285.
 Hasson, J., Berkman, J. I., Parker, J. G., and Rifkin, H. (1957). *Ann. intern. Med.*, **47**, 493.
 Kark, R. M. (1955). In *Proceedings 7th Annual Conference Nephrotic Syndrome*, p. 141. National Kidney Disease Foundation, New York.
 Morris, J. F., Ginn, H. E., and Thompson, D. D. (1963). *Amer. J. Med.*, **34**, 867.
 Pollak, V. E., Kark, R. M., Pirani, C. L., Shafter, H. A., and Muehrcke, R. C. (1956). *Ibid.*, **21**, 496.
 Rowntree, L. G., Fitz, R., and Geraghty, J. T. (1913). *Arch. intern. Med.*, **11**, 121.
 Shattock, S. G. (1913). *Proc. roy. Soc. Med.*, **6**, pt. 3, Pathological Section, 126.
 Squire, J. R. (1953). *Brit. med. J.*, **2**, 1389.
 Vilk, N. L. (1940). *Klin. Med. (Mosk.)*, **18**, 91.