signs of any abnormality in the central nervous system. The blood-pressure was known to have been within normal limits previously. The patient was put to bed and sedated, but it was later found that she could not be roused; when seen again by her doctor two hours after the first examination she was comatose, with a blood-pressure of 170/100 mm. Hg; the patient was sent to hospital, but died very shortly after admission.

Post-mortem examination showed a haemorrhage in the left cerebral hemisphere disrupting the internal capsule and adjacent areas of the left corpus striatum. No other abnormality was found—in particular, no intracranial aneurysms—and no evidence of long-standing hypertension.

—We are, etc.,

J. T. ALBAN LLOYD.
D. R. H. WALKER.

Department of Pathology,
Royal Alexandra Hospital,
Rhyll, Flintshire.

Anaesthesia and Monoamine-oxidase Inhibitors

SIR,—I read with interest the letters of Dr. H. C. Churchill-Davidson (20 February, p. 520) and of Dr. M. D. A. Vickers (24 April, p. 1126). I should like to comment that indeed many physicians believe that if any major side-effect occurs in response to a newer drug it is due to the "sensitivity," and somehow an allergic reaction is implied.

As a psychiatrist I have investigated most tranquillizers and energizers, and while we have, fortunately, relatively few severe and fatal reactions to these drugs, such reactions do occur, sometimes due to avoidable circumstances. One of the side-effects from these psychotropic drugs which has been neglected much too often is secondary hypotension. Fatal reactions to shock occurring from hypotension may be avoided at times if the "predisposition" of the patient is known. It has become the belief of many researchers in this field that side-effects from these drugs occur more frequently in the presence of a diffuse impairment of the autonomic nervous system. The observation of the blood-pressure pattern may be helpful. In a recent survey of the blood-pressure patterns in a 600-bed mental-patient hospital we found that only 57.44% had normal blood-pressure patterns; 8.48% were classified as hypertensive; hypotension was present in 18.86%; and labile-borderline patterns were found in 15.22%. This means that hypotension is more frequent than is in general believed (at least three times as frequent as in a general population).

It is quite possible that hypotension is present to a marked degree in patients who take psychotropic drugs for some time. When these patients are in need of anaesthesia and surgery and are given analgesic drugs or additional phenothiazines as a sedative and/or anti-emetic, shock may be aggravated and may result in death; nausea in itself is considered a shock-producing agent.¹ It would probably be best to avoid giving monoamine-oxidase inhibitors during anaesthesia and immediately following surgery, because of their severe hypotensive action, especially in the elderly. This problem may easily be overlooked because of a latent cardiovascular lability frequently present and the unpredictable response of the autonomic nervous system to the monoamine-oxidase inhibitors. The physician and surgeon are quite often

unaware of the status of the autonomic nervous system of his patient, which is not easily established.

While the blood-pressure is a good measure of the autonomic nervous system, a frequent mistake of many researchers, as well as the practising physician and surgeon, has been to use only one or two blood-pressure measures taken at random. By taking systematic blood-pressure measures at regular intervals—for instance, four times a day for one week, blood-pressure patterns may be established which permit the physician to ascertain the presence of any abnormality such as hypertension, orthostatic hypotension, or a labile-borderline condition, any one of which may be a warning in the use of psychotropic drugs.²

There are many factors in drug administration about which little is known. Doubtless an important one is the occurrence of metabolic by-products, which may act as toxins. In the recent discussions of tranlycypromine (Parnate), an effective monoamine-oxidase inhibitor, Himwich³ concluded that imipramine (Tofranil) therapy promotes the accumulation of some substance in the brain which is normally destroyed along a pathway blocked by tranlycypromine. While it is impossible to generalize, any toxic reaction to drug substitutes (e.g., pethidine for morphine) may occur on a similar hypothesis—that is, the accumulation of toxic metabolites which may be fatal.—I am, etc.,

KURT WITTON.

Veterans Administration Hospital,
Fort Meade,
S. Dakota, U.S.A.

REFERENCES

- 1 Selzer, A., and Rytand, D. A., *J. Amer. med. Ass.*, 1958, **168**, 762.
- 2 Witton, K., and Goldman, A. R., *J. nerv. ment. Dis.*, 1965, **140**, 58.
- 3 Himwich, W. A., *Rec. Adv. biol. Psychiat.*, 1961, **4**, 257.

Imipramine in Parkinsonism

SIR,—I would like to comment on the trial of imipramine against a placebo in the treatment of Parkinsonism, reported by Dr. R. R. Strang (3 July, p. 32).

Twenty-seven out of 35 of his placebo group were removed from the trial because of clinical deterioration or failure to respond. None of his imipramine group was removed for this reason. We are not told whether, in the case of any of the patients receiving imipramine, the code was broken because of failure to respond and the patient kept on the trial because he was found to be receiving imipramine. If this did not occur, the difference in failure rate between the two groups is very large, χ^2 being over 39 with one degree of freedom, surely the highest degree of improbability in any of the many reported trials of imipramine.

If, on the other hand, patients receiving imipramine were selectively retained in the trial, and thus given a further chance to improve, a bias has been introduced in favour of imipramine. We cannot therefore conclude that "a most satisfactory degree of improvement in 16 patients (53%) confirms the value of imipramine in Parkinsonism."

Incidentally, Dr. Strang need not be surprised to find imipramine of value in Parkinsonism in spite of its chemical simi-

larity to the phenothiazines. Of the anti-Parkinsonism drugs listed in the 1958 edition of Martindale's *Extra Pharmacopoeia* two are phenothiazine derivatives (ethopropazine hydrochloride and diethazine hydrochloride). Two others are reported to alleviate mental depression in patients with Parkinsonism (benzhexol hydrochloride and cyrcimine hydrochloride).—I am, etc.,

London S.E.19.

J. S. PRICE.

Undescended Testicle

SIR,—Your leading article "Undescended Testicle" (15 May, p. 1259) called to mind a diagnostic test differentiating pseudocryptorchidism from cryptorchidism, described by Dr. Paul L. Bunce,¹ the original description being attributed by him to Mr. Louis Orr. I have found the test most useful in examining large numbers of boys at school examination. It prevents many needless over-referrals to urologists, a source of annoyance to urologist and parents.

In essence, the test consists of having the boy sit on a straight-backed chair and draw up his legs so that the soles of his feet rest on the chair seat. By hugging his knees to his chest pressure is directed to the inguinal canal in such a way that a freely movable testis will appear in the upper scrotum, where it can be examined without difficulty.—I am, etc.,

Utica,
New York, U.S.A.

JOHN H. POWELL.

REFERENCE

- 1 Bunce, P. L., *Pediatrics*, 1961, **27**, 165.

SIR,—The increased risk of malignancy in the ectopic or abdominal testis is, as Dr. R. T. H. Shepherd (26 June, p. 1670) says, a powerful argument in favour of treatment of the maldescended testis. I suggest, however, that in advising orchidectomy Dr. Shepherd may have overlooked one important point. Whilst many studies, including my own,¹ have noted a marked tendency in the undescended testis to undergo malignant change this has only been found in testes remaining in the undescended position through puberty, whether or not they are subsequently brought down by surgery; it appears to be exceedingly rare for the testis undergoing orchidectomy before puberty to develop malignant change. This is an important reason for deciding well in advance whether the testis can descend to the scrotum with puberty (i.e., the retractile variety) and undertaking orchidectomy without delay if spontaneous descent does not seem possible. This offers protection against malignancy, and at this age sacrificing an otherwise well-formed testis would not seem justifiable. I do agree with Dr. Shepherd, however, that when puberty has overtaken a testis unable to descend there is a strong case for orchidectomy rather than accepting the enhanced risk (probably about one in fifty) that will otherwise remain in spite of orchidectomy. It is worth emphasizing that this dilemma need not arise, since a testis of this sort can be recognized without difficulty at an early age, and of course nowadays few escape our system of school health examinations.

It seems important that our knowledge upon this should be more detailed, and may

I repeat the appeal made through your columns in 1963 (4 May, p. 1232) that those with access to the information should let us know more precisely the frequency of malignancy in testes which have undergone orchid-ectomy before puberty. Perhaps Dr. Shepherd

could re-examine his data upon this point?

—I am, etc.,

The Radcliffe Infirmary,
Oxford.

D. J. TIBBS.

REFERENCE

¹ Tibbs, D. J., *Brit. J. Surg.*, 1961, 48, 601.

Abuse of Digitalis

SIR,—In your leading article on the abuse of digitalis (8 May, p. 1208) you emphasize the frequency of disorders of cardiac rhythm. We wish to report a patient with acute digitalis poisoning who developed a great variety of rhythm disturbances.

In an attempt to commit suicide a 15-year-old girl swallowed the contents of one and a half vials of Nativelle's Digitaline (equivalent to 15 mg. of *Digitalis purpurea*). Three hours later gastric lavage was performed and six hours later she was transferred to our unit. On admission she was in good general condition. Blood-

pressure was 110/70 mm. Hg, the heart rate 68 beats per min., there were no irregularities of cardiac rhythm. For the first six hours after her admission she vomited repeatedly; then this subsided, and for the rest of her stay in hospital she was virtually symptom-free apart from some nausea.

The unusual feature of the successive E.C.G. records that were taken was the great variety of rhythm disturbances encountered (see Fig.), almost all the rhythm changes described in digitalis poisoning^{1,2} being seen at one time or another.

The patient was treated with atropine, potassium chloride, and sodium citrate. Potassium was given orally with extreme caution, with twice-daily serum-electrolyte determinations and careful inspection of the E.C.G. records for development of hyperpotassaemia.

The patient left hospital 24 days after her admission in perfect health; her E.C.G., apart from the ordinary digitalis effects, was normal.

—We are, etc.,

D. THOMOPOULOS.

Second Medical Unit,
Evangelismos Hospital,
Athens, Greece.

REFERENCES

- ¹ Bellet, S., *Clinical Disorders of the Heart Beat*, 2nd ed., 1963. Lea and Febiger, Philadelphia.
- ² Wood, P., *Diseases of the Heart and Circulation*, 2nd edition, 1956. Eyre and Spottiswoode, London.

Successful Surgical Treatment of Heart Stab Wound

SIR,—I wish to report a successful instance of surgical repair of a stab wound in the heart.

On 25 November 1964 a young boy aged 15 years was brought into the General Hospital, Port of Spain, Trinidad, with a history that he was stabbed over his heart. On seeing him in the ward at hospital, he was in a great state of collapse. His pulse was imperceptible; he was sweating, pale, and clammy. On examination there was an incised wound about $\frac{1}{2}$ in. (1.9 cm.) long, about 1 in. (2.5 cm.) from the left border of the sternum, in the fourth left intercostal space. On auscultation the heart sounds could be heard, but very feebly. The size of the heart by percussion appeared to be large. X-ray of the chest showed an enlarged heart shadow. This confirmed the diagnosis of a penetrating wound into the heart causing tamponade. His shock was treated with $\frac{1}{2}$ gr. (10 mg.) of morphine, and a bottle of Intradex (dextran) was given intravenously prior to blood-transfusions, and the operating theatre was notified.

Within 45 minutes he was on the operating table and anaesthetized. By that time I could just feel his pulse and his blood-pressure could just be read. Thoracotomy was performed in the left 4th intercostal space by an anterolateral incision. The pericardium was bulging and blue. On opening the pericardium contained much blood and blood clots were evacuated. The heart, which was now relieved of its tamponade, began to beat more forcefully and to emit blood from the stab wound, which was found in the right ventricle. The wound was about $\frac{1}{2}$ in. (1.25 cm.) long and was closed by interrupted silk sutures. Part of the pericardium was left open and the left pleural cavity was drained by sealed drainage.

After operation his condition was better than it was before. He made an uneventful recovery except for intermittent fever the first four days after operation. He was able to leave hospital 16 days after. He was seen four months later and is keeping in perfect health.

—I am, etc.,

General Hospital,
Port of Spain,
Trinidad.

L. A. H. MCSHINE.

Successive E.C.G. records (the most typical leads only are given). (a) (Date of admission 28 June 1964) Sinus rhythm, rate 68, P-R=0.20 sec., ST segment depression, "dragged" downward T waves. Q-Tc=0.27 sec. (b) (29 June 1964, 9 a.m.) Sinus rhythm, rate 83, P-R=0.22 sec. (first degree A-V block). Q-Tc=0.26 sec. (c) (29 June 1964, 1 p.m.) Atrial rate 96. First and second degree A-V block. (d) 29 June 1964, 8 p.m.) First degree A-V block. Q-Tc=0.25 sec. (e) (1 July 1964) Sinus rhythm, rate 32. S-A heart block (? 2:1 or 3:1). (f) (2 July 1964) wandering pacemaker. (g) (3 July 1964) S-A heart block (4:3). In other leads not given in the Figure there was a great variety of response—for example: 7:6, 5:4, 6:5, 7:6, 8:7, 14:13. (h) (4 July 1964) Sinus bradycardia, rate 55. (i) (8 July 1964) Sinus bradycardia, rate 58. (j) (21 July 1964) Regular sinus rhythm, rate 62, ST segment nearly restored, Q-Tc=0.38 sec.