I repeat the appeal made through your columns in 1963 (4 May, p. 1232) that those with access to the information should let us know more precisely the frequency of malignancy in testes which have undergone orchid-ectomy before puberty. Perhaps Dr. Shepherd

could re-examine his data upon this point?

—I am, etc.,

The Radcliffe Infirmary,
Oxford.

D. J. TIBBS.

REFERENCE

¹ Tibbs, D. J., *Brit. J. Surg.*, 1961, 48, 601.

Abuse of Digitalis

SIR,—In your leading article on the abuse of digitalis (8 May, p. 1208) you emphasize the frequency of disorders of cardiac rhythm. We wish to report a patient with acute digitalis poisoning who developed a great variety of rhythm disturbances.

In an attempt to commit suicide a 15-year-old girl swallowed the contents of one and a half vials of Nativelle's Digitaline (equivalent to 15 mg. of *Digitalis purpurea*). Three hours later gastric lavage was performed and six hours later she was transferred to our unit. On admission she was in good general condition. Blood-

pressure was 110/70 mm. Hg, the heart rate 68 beats per min., there were no irregularities of cardiac rhythm. For the first six hours after her admission she vomited repeatedly; then this subsided, and for the rest of her stay in hospital she was virtually symptom-free apart from some nausea.

The unusual feature of the successive E.C.G. records that were taken was the great variety of rhythm disturbances encountered (see Fig.), almost all the rhythm changes described in digitalis poisoning^{1,2} being seen at one time or another.

The patient was treated with atropine, potassium chloride, and sodium citrate. Potassium was given orally with extreme caution, with twice-daily serum-electrolyte determinations and careful inspection of the E.C.G. records for development of hyperpotassaemia.

The patient left hospital 24 days after her admission in perfect health; her E.C.G., apart from the ordinary digitalis effects, was normal.

—We are, etc.,

D. THOMOPOULOS.

Second Medical Unit,
Evangelismos Hospital,
Athens, Greece.

REFERENCES

- ¹ Bellet, S., *Clinical Disorders of the Heart Beat*, 2nd ed., 1963. Lea and Febiger, Philadelphia.
- ² Wood, P., *Diseases of the Heart and Circulation*, 2nd edition, 1956. Eyre and Spottiswoode, London.

Successful Surgical Treatment of Heart Stab Wound

SIR,—I wish to report a successful instance of surgical repair of a stab wound in the heart.

On 25 November 1964 a young boy aged 15 years was brought into the General Hospital, Port of Spain, Trinidad, with a history that he was stabbed over his heart. On seeing him in the ward at hospital, he was in a great state of collapse. His pulse was imperceptible; he was sweating, pale, and clammy. On examination there was an incised wound about $\frac{1}{2}$ in. (1.9 cm.) long, about 1 in. (2.5 cm.) from the left border of the sternum, in the fourth left intercostal space. On auscultation the heart sounds could be heard, but very feebly. The size of the heart by percussion appeared to be large. X-ray of the chest showed an enlarged heart shadow. This confirmed the diagnosis of a penetrating wound into the heart causing tamponade. His shock was treated with $\frac{1}{2}$ gr. (10 mg.) of morphine, and a bottle of Intradex (dextran) was given intravenously prior to blood-transfusions, and the operating theatre was notified.

Within 45 minutes he was on the operating table and anaesthetized. By that time I could just feel his pulse and his blood-pressure could just be read. Thoracotomy was performed in the left 4th intercostal space by an anterolateral incision. The pericardium was bulging and blue. On opening the pericardium contained much blood and blood clots were evacuated. The heart, which was now relieved of its tamponade, began to beat more forcefully and to emit blood from the stab wound, which was found in the right ventricle. The wound was about $\frac{1}{2}$ in. (1.25 cm.) long and was closed by interrupted silk sutures. Part of the pericardium was left open and the left pleural cavity was drained by sealed drainage.

After operation his condition was better than it was before. He made an uneventful recovery except for intermittent fever the first four days after operation. He was able to leave hospital 16 days after. He was seen four months later and is keeping in perfect health.

—I am, etc.,

General Hospital,
Port of Spain,
Trinidad.

L. A. H. MCSHINE.

Successive E.C.G. records (the most typical leads only are given). (a) (Date of admission 28 June 1964) Sinus rhythm, rate 68, P-R=0.20 sec., ST segment depression, "dragged" downward T waves. Q-Tc=0.27 sec. (b) (29 June 1964, 9 a.m.) Sinus rhythm, rate 83, P-R=0.22 sec. (first degree A-V block). Q-Tc=0.26 sec. (c) (29 June 1964, 1 p.m.) Atrial rate 96. First and second degree A-V block. (d) 29 June 1964, 8 p.m.) First degree A-V block. Q-Tc=0.25 sec. (e) (1 July 1964) Sinus rhythm, rate 32. S-A heart block (? 2:1 or 3:1). (f) (2 July 1964) wandering pacemaker. (g) (3 July 1964) S-A heart block (4:3). In other leads not given in the Figure there was a great variety of response—for example: 7:6, 5:4, 6:5, 7:6, 8:7, 14:13. (h) (4 July 1964) Sinus bradycardia, rate 55. (i) (8 July 1964) Sinus bradycardia, rate 58. (j) (21 July 1964) Regular sinus rhythm, rate 62, ST segment nearly restored, Q-Tc=0.38 sec.