Birefringence is, of course, exhibited by both urate and calcium pyrophosphate crystals. Differentiation between the two requires identification of the sign of birefringence, urate being strongly negative, calcium pyrophosphate weakly positive. Proof of the identity of urate crystals is by uricase digestion.

We have had the opportunity of examining synovial fluids from 22 patients with radiological calcification of joint cartilage. In general our findings have given less clear-cut answers than anticipated, and a number of specimens contained more than one type of crystal. In one patient with associated hyperparathyroidism the synovial fluid on two occasions was found to contain crystals which exhibited strong positive birefringence and had an extinction angle of 30°. Histochemical studies showed the presence of calcium. The other radical has not yet been positively identified, but we are impressed with the resemblance of these crystals to calcium oxalate.

It may well be that gout and pseudogout can co-exist in the same person. One patient in this series is regarded as suffering from classical uric-acid gout, yet on one occasion fluid from his inflamed knee-joint yielded typical calcium pyrophosphate crystals.—We are, etc.,

H. L. F. CURREY.
K. V. SWETTENHAM.

The London Hospital,
London E.1.

REFERENCES

- ¹ *Brit. med. J.*, 1964, 2, 889.
- ² Bywaters, E. G. L., Dixon, A. St. J., and Scott, J. T., *Ann. rheum. Dis.*, 1963, 22, 171.

Hospital at Bath

SIR,—The Royal National Hospital for Rheumatic Diseases at Bath, the third oldest voluntary hospital outside London, has just reopened after almost complete rebuilding. The first hospital in the world to devote itself to the treatment of arthritis, it was bombed in 1942, but it still has many interesting records and incidentally two fine pieces of Roman mosaic in its foundations. We are now anxious to build up a small museum, and if any readers have for disposal such relics as cupping-glasses, leech tubes, bone or wooden stethoscopes, blood-letting sets, etc., I would be very interested to hear from them.—I am, etc.,

Bath, Somerset. GEORGE D. KERSLEY.

Acute Hepatitis

SIR,—Since the investigations of Allen and Sayman¹ and more recently by Mirick *et al.*² in the United States, it has become increasingly clear that the risk of post-transfusion hepatitis, particularly in recipients aged over 40, is considerable. The incidence of post-transfusion hepatitis can be significantly reduced by the judicious selection of patients for transfusion, by concurrent administration of gamma-globulin, or by the use of hepatitis-free plasma infusions. It is evident that the supplies of human gamma-globulin would not allow its free use with blood transfusion, and our need for hepatitis-free plasma for replacement therapy is thus more urgent.

In this country an estimate of the size of the problem of hepatitis cannot be made since hepatitis is not notifiable on a national basis. Before we can begin to undertake any preventive measures the first step should clearly be the notification of hepatitis and the establishment of a follow-up system for patients who have received blood transfusions.—I am, etc.,

A. J. ZUCKERMAN.

Department of Bacteriology and Immunology,
London School of Hygiene and
Tropical Medicine,
London W.C.1.

REFERENCES

- ¹ Allen, J. G., and Sayman, W. A., *J. Amer. med. Ass.*, 1962, 180, 1079.
- ² Mirick, G. S., Ward, R., and McCollum, R. W., *New Engl. J. Med.*, 1965, 273, 59.

Otogenic Tetanus

SIR,—The case of otogenic tetanus described by Drs. I. T. Boyle, W. C. Brown, and E. A. Richardson (10 July, p. 89) prompts us to record a further case.

The patient was a boy aged 9 who was subject to repeated attacks of bronchitis and who had suffered from an intermittent discharge from the right ear from the age of 4. This ear, which had been dry for nine months, became painful and started to discharge eight days before admission to Pembury Hospital. Three days before admission he fell into a duck pond, and on the day before admission he complained of a stiff neck and difficulty in opening his mouth. Spasms developed on the day of admission and rapidly became frequent and severe. In the past he had probably received antitetanus serum for a laceration of the wrist, but there was no record of the administration of toxoid either then or later.

On examination at Pembury Hospital the patient was obviously suffering from severe tetanus, with marked trismus and opisthotonos, and severe spasms. His temperature was 100° F. (37.8° C.) and the pulse rate was 120/min. Pus was discharging from the right ear, but there was no tenderness, swelling, or oedema over the mastoid. A bacterial swab taken at this time subsequently grew *Clostridium tetani*.

The patient was given 2 mega-units of crystalline penicillin and sedated with bromethol. In view of the short period of onset (12 hours) and the probable shortness of the incubation period (three days), he was anaesthetized and paralysed with D-tubocurarine, and a cuffed endotracheal tube was inserted. Ventilation was controlled by hand during the journey to Hammersmith Hospital and subsequently was maintained by an East-Radcliffe ventilator. Shortly after transfer a tracheostomy and right mastoidectomy were performed. 100,000 units of A.T.S. were given intravenously before the surgical intervention, and the spasms were controlled with intermittent injections of D-tubocurarine (30 mg. intramuscularly). Sedation was given at night, but not during the day. Total paralysis was maintained for 16 days, but during this time the patient managed to communicate with the nurses by squeezes of the hand. The progress of the disease was checked at intervals by withholding curare until definite spasms occurred. A right middle lobe collapse, which was present on admission, cleared rapidly with physiotherapy and there were no other complications. Weaning from the ventilator was commenced on the 17th day (sedation being maintained with promazine and quinalbarbitone) and the tracheostomy tube was removed seven days after discontinuing the curare. Ten days later he was able to start walking and the residual muscle stiffness had disappeared. Apart from complete deafness in the right ear, the patient made an uninterrupted recovery.

Immunization with tetanus toxoid was commenced before leaving hospital.

The interest of this case lies in the possibility of contamination of the ear by submersion in the duck pond. If infection entered at this time the shortness of the incubation period may be related to the shortness of the route between the middle ear and the central nervous system.

We are grateful to Dr. N. M. Jacoby, consultant paediatrician, Pembury Hospital, for details of this case.

—We are, etc.,

M. K. SYKES.

J. A. DAVIS.

Departments of Anaesthesia
and Child Health,
Hammersmith Hospital,
London W.12.

Poisons Information Centre

SIR,—Earlier this year the 7505 United States Air Force Hospital at Burderop Park, near Swindon, completely closed down, so the Poisons Information Centre, to which you refer in your leading article (24 July, p. 185), is no longer operational. This centre readily and freely offered its services to British doctors, and in this context it is opportune to mention that this was just one of many examples of the excellent liaison and friendly co-operation which existed between American and British doctors in this particular area.

With the closure of American air bases and the 7505 U.S.A.F. Hospital, the Oxford Group of the Anglo-American Medical Society has, regrettably, had to be dissolved, but the Society continues to meet in the Cambridge and London area. The Anglo-American Medical Society was founded in 1961 to "meet the great need that obviously existed for an organization through which British and American doctors could obtain a clearer knowledge and better understanding of each other." Contact was made through regular scientific programmes and social meetings, and I know from discussions I have had with my colleagues, and from letters I receive from American doctors who have now returned to the United States, that the aims and ideals of the Anglo-American Society were attained to the mutual and, I believe, lasting benefit of both communities.—I am, etc.,

A. G. FREEMAN,
Past Chairman,
Anglo-American Medical Society
(Oxford Group).

Swindon,
Wiltshire.

Thoracic Actinomycosis

SIR,—I wish to report a further interesting case of actinomycotic infection, having read the letter of Dr. B. Bissu (24 July, p. 231) attracting attention to this disease.

A labourer aged 20 years was admitted to Whipps Cross Hospital on 22 April 1965 with a history of an injury to his right loin sustained eight weeks previously in a fight. He had had pain in his right chest on breathing for three weeks before admission and had lost about 3 st. (19.1 kg.) in weight since his injury; he also complained of anorexia and had been sweating at night but had no cough. On examination he was febrile (101° F. (38.3° C.)) and looked ill, showed no obvious dental caries, had a dorsal scoliosis to the left, and there was a tender